

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
3 January 2002 (03.01.2002)

PCT

(10) International Publication Number
WO 02/00690 A2

- (51) International Patent Classification⁷: C07K 14/00 09/854,208 10 May 2001 (10.05.2001) US
09/854,280 10 May 2001 (10.05.2001) US
(21) International Application Number: PCT/US01/19692 09/866,028 25 May 2001 (25.05.2001) US
09/866,034 25 May 2001 (25.05.2001) US
(22) International Filing Date: 20 June 2001 (20.06.2001) PCT/US01/17092 25 May 2001 (25.05.2001) US
09/870,574 30 May 2001 (30.05.2001) US
(25) Filing Language: English PCT/US01/17443 30 May 2001 (30.05.2001) US
PCT/US01/17800 1 June 2001 (01.06.2001) US
(26) Publication Language: English
(30) Priority Data:
60/213,637 23 June 2000 (23.06.2000) US
60/219,556 20 July 2000 (20.07.2000) US
60/220,624 25 July 2000 (25.07.2000) US
60/220,664 25 July 2000 (25.07.2000) US
PCT/US00/20710 28 July 2000 (28.07.2000) US
60/222,695 2 August 2000 (02.08.2000) US
09/643,657 17 August 2000 (17.08.2000) US
PCT/US00/23522 23 August 2000 (23.08.2000) US
PCT/US00/23328 24 August 2000 (24.08.2000) US
60/230,978 7 September 2000 (07.09.2000) US
60/000,000 15 September 2000 (15.09.2000) US
09/664,610 18 September 2000 (18.09.2000) US
09/665,350 18 September 2000 (18.09.2000) US
60/242,922 24 October 2000 (24.10.2000) US
09/709,238 8 November 2000 (08.11.2000) US
PCT/US00/30952 8 November 2000 (08.11.2000) US
PCT/US00/30873 10 November 2000 (10.11.2000) US
PCT/US00/32678 1 December 2000 (01.12.2000) US
09/747,259 20 December 2000 (20.12.2000) US
PCT/US00/34956 20 December 2000 (20.12.2000) US
09/767,609 22 January 2001 (22.01.2001) US
09/796,498 28 February 2001 (28.02.2001) US
PCT/US01/06520 28 February 2001 (28.02.2001) US
PCT/US01/06666 1 March 2001 (01.03.2001) US
09/802,706 9 March 2001 (09.03.2001) US
09/808,689 14 March 2001 (14.03.2001) US
09/816,744 22 March 2001 (22.03.2001) US
09/828,366 5 April 2001 (05.04.2001) US
(71) Applicant (for all designated States except US): GENENTECH, INC. [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US).
(72) Inventors; and
(75) Inventors/Applicants (for US only): BAKER, Kevin, P. [GB/US]; 14006 Indian Run Drive, Darnestown, MD 20878 (US). FERRARA, Napoleone [US/US]; #704, 2090 Pacific Avenue, San Francisco, CA 94109 (US). GERBER, Hanspeter [CH/US]; #5, 1121 Tennessee Street, San Francisco, CA 94107 (US). GERRITSEN, Mary, E. [CA/US]; 541 Parrott Drive, San Mateo, CA 94402 (US). GODDARD, Audrey [CA/US]; 110 Congo Street, San Francisco, CA 94131 (US). GODOWSKI, Paul, J. [US/US]; 2627 Easton Drive, Burlingame, CA 94010 (US). GURNEY, Austin, L. [US/US]; 1 Debbie Lane, Belmont, CA 94002 (US). HILLAN, Kenneth, J. [GB/US]; 64 Seward Street, San Francisco, CA 94114 (US). MARSTERS, Scot, A. [US/US]; 990 Cherry Street, San Carlos, CA 94070 (US). PAN, James [CA/US]; 2705 Coronet Boulevard, Belmont, CA 94002 (US). PAONI, Nicholas, F. [US/US]; 1756 Terrace Drive, Belmont, CA 94002 (US). STEPHAN, Jean-Philippe, F. [FR/US]; 320 C Lansdale Avenue, Millbrae, CA 94030 (US). WATANABE, Colin, K. [US/US]; 128 Corliss Drive, Moraga, CA 94556 (US). WILLIAMS, P., Mickey [US/US]; 509 Alto Avenue, Half Moon Bay, CA 94019 (US). WOOD, William, I. [US/US]; 35 Southdown Court, Hillsborough, CA 94010 (US). YE, Weilan [CN/US]; 119 Barkentine Street, Foster City, CA 94404 (US).
(74) Agents: AGARWAL, Atulya, R. et al.; c/o GENENTECH, INC., MS49, 1 DNA Way, South San Francisco, CA 94080-4990 (US).

[Continued on next page]

(54) Title: COMPOSITIONS AND METHODS FOR THE DIAGNOSIS AND TREATMENT OF DISORDERS INVOLVING ANGIOGENESIS

(57) Abstract: Compositions and methods are disclosed for stimulating or inhibiting angiogenesis and/or cardiovascularization in mammals, including humans. Pharmaceutical compositions are based on polypeptides or antagonists thereto that have been identified for one or more of these uses. Disorders that can be diagnosed, prevented, or treated by the compositions herein include trauma such as wounds, various cancers, and disorders of the vessels including atherosclerosis and cardiac hypertrophy. In addition, the present invention is directed to novel polypeptides and to nucleic acid molecules encoding those polypeptides. Also provided herein are vectors and host cells comprising those nucleic acid sequences, chimeric polypeptide molecules comprising the polypeptides of the present invention fused to heterologous polypeptide sequences, antibodies which bind to the polypeptides of the present invention and to methods for producing the polypeptides of the present invention.

BEST AVAILABLE COPY

WO 02/00690 A2



(81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

COMPOSITIONS AND METHODS FOR THE DIAGNOSIS AND TREATMENT OF DISORDERS INVOLVING ANGIOGENESIS

1. Field of the Invention

The present invention relates to compositions and methods useful for the modulation (e.g., promotion or inhibition) of angiogenesis and/or cardiovascularization in mammals in need of such biological effect. The present invention further relates to the diagnosis and treatment of disorders involving angiogenesis (e.g., cardiovascular as well as oncological disorders).

2. Background of the Invention

2.1. Angiogenesis

Angiogenesis, defined as the growth or sprouting of new blood vessels from existing vessels, is a complex process that primarily occurs during embryonic development. Under normal physiological conditions in adults, angiogenesis takes place only in very restricted situations such as hair growth and wounding healing (Auerbach, W. and Auerbach, R., 1994, *Pharmacol Ther* 63(3):265-311; Ribatti et al., 1991, *Haematologica* 76(4):311-20; Risau, 1997, *Nature* 386(6626):671-4). Unregulated angiogenesis has gradually been recognized to be responsible for a wide range of disorders, including, but not limited to cardiovascular disease, cancer, rheumatoid arthritis, psoriasis and diabetic retinopathy (Folkman, 1995, *Nat Med* 1(1):27-31; Isner, 1999, *Circulation* 99(13):1653-5; Koch, 1998, *Arthritis Rheum* 41(6):951-62; Walsh, 1999, *Rheumatology (Oxford)* 38(2):103-12; Ware and Simons, 1997, *Nat Med* 3(2):158-64).

2.2. Cardiac Disorders and Factors

Heart failure affects approximately five million Americans, and new cases of heart failure number about 400,000 each year. It is the single most frequent cause of hospitalization for people age 65 and older in the United States. Recent advances in the management of acute cardiac diseases, including acute myocardial infarction, are resulting in an expanding patient population that will eventually develop chronic heart failure. From 1979 to 1995, hospitalizations for congestive heart failure (CHF) rose from 377,000 to 872,000 (a 130 percent increase) and CHF deaths increased 116 percent.

CHF is a syndrome characterized by left ventricular dysfunction, reduced exercise tolerance, impaired quality of life, and markedly shortened life expectancy. The sine qua non of heart failure is an inability of the heart to pump blood at a rate sufficient to meet the metabolic needs of the body's tissues (in other words, there is insufficient cardiac output).

At least four major compensatory mechanisms are activated in the setting of heart failure to boost cardiac output, including peripheral vasoconstriction, increased heart rate, increased cardiac contractility, and increased plasma volume. These effects are mediated primarily by the sympathetic nervous system and the renin-angiotensin system. See, Eichhorn, American Journal of Medicine, 104: 163-169 (1998). Increased output from the

sympathetic nervous system increases vascular tone, heart rate, and contractility. Angiotensin II elevates blood pressure by 1) directly stimulating vascular smooth muscle contraction, 2) promoting plasma volume expansion by stimulating aldosterone and antidiuretic hormone secretion, 3) stimulating sympathetic-mediated vascular tone, and 4) catalyzing the degradation of bradykinin, which has vasodilatory and natriuretic activity. *See*, review by Brown and Vaughan, Circulation, 97: 1411-1420 (1998). As noted below, angiotensin II may also have directly deleterious effects on the heart by promoting myocyte necrosis (impairing systolic function) and intracardiac fibrosis (impairing diastolic and in some cases systolic function). *See*, Weber, Circulation, 96: 4065-4082 (1998).

A consistent feature of congestive heart failure (CHF) is cardiac hypertrophy, an enlargement of the heart that is activated by both mechanical and hormonal stimuli and enables the heart to adapt to demands for increased cardiac output. Morgan and Baker, Circulation, 83: 13-25 (1991). This hypertrophic response is frequently associated with a variety of distinct pathological conditions such as hypertension, aortic stenosis, myocardial infarction, cardiomyopathy, valvular regurgitation, and intracardiac shunt, all of which result in chronic hemodynamic overload.

Hypertrophy is generally defined as an increase in size of an organ or structure independent of natural growth that does not involve tumor formation. Hypertrophy of the heart is due either to an increase in the mass of the individual cells (myocytes), or to an increase in the number of cells making up the tissue (hyperplasia), or both. While the enlargement of an embryonic heart is largely dependent on an increase in myocyte number (which continues until shortly after birth), post-natal cardiac myocytes lose their proliferative capacity. Further growth occurs through hypertrophy of the individual cells.

Adult myocyte hypertrophy is initially beneficial as a short term response to impaired cardiac function by permitting a decrease in the load on individual muscle fibers. With severe, long-standing overload, however, the hypertrophied cells begin to deteriorate and die. Katz, "Heart Failure", in: Katz A.M. ed., Physiology of the Heart (New York: Raven Press, 1992) pp. 638-668. Cardiac hypertrophy is a significant risk factor for both mortality and morbidity in the clinical course of heart failure. Katz, Trends Cardiovasc. Med., 5: 37-44 (1995). For further details of the causes and pathology of cardiac hypertrophy *see, e.g., Heart Disease. A Textbook of Cardiovascular Medicine*, Braunwald, E. ed. (W.B. Saunders Co., 1988), Chapter 14, "Pathophysiology of Heart Failure."

On a cellular level, the heart is composed of myocytes and surrounding support cells, generically called non-myocytes. While non-myocytes are primarily fibroblast/mesenchymal cells, they also include endothelial and smooth muscle cells. Indeed, although myocytes make up most of the adult myocardial mass, they represent only about 30% of the total cell numbers present in heart. In response to hormonal, physiological, hemodynamic, and pathological stimuli, adult ventricular muscle cells can adapt to increased workloads through the activation of a hypertrophic process. This response is characterized by an increase in myocyte cell size and contractile protein content of individual cardiac muscle cells, without concomitant cell division and activation of embryonic genes, including the gene for atrial natriuretic peptide (ANP). Chien *et al.*, FASEB J., 5: 3037-3046 (1991); Chien *et al.*, Annu. Rev. Physiol., 55: 77-95 (1993). An increment in myocardial mass as a result of an increase in myocyte size that is associated with an accumulation of interstitial collagen within the extracellular matrix and around intramyocardial coronary arteries has been described in left ventricular hypertrophy secondary to pressure overload

in humans. Caspari *et al.*, Cardiovasc. Res., **11**: 554-558 (1977); Schwarz *et al.*, Am. J. Cardiol., **42**: 895-903 (1978); Hess *et al.*, Circulation, **63**: 360-371 (1981); Pearlman *et al.*, Lab. Invest., **46**: 158-164 (1982).

It has also been suggested that paracrine factors produced by non-myocyte supporting cells may additionally be involved in the development of cardiac hypertrophy, and various non-myocyte derived hypertrophic factors, such as, leukocyte inhibitory factor (LIF) and endothelin, have been identified. Metcalf, Growth Factors, **7**: 169-173 (1992); Kurzrock *et al.*, Endocrine Reviews, **12**: 208-217 (1991); Inoue *et al.*, Proc. Natl. Acad. Sci. USA, **86**: 2863-2867 (1989); Yanagisawa and Masaki, Trends Pharm. Sci., **10**: 374-378 (1989); U.S. Patent No. 5,573,762 (issued November 12, 1996). Further exemplary factors that have been identified as potential mediators of cardiac hypertrophy include cardiotrophin-1 (CT-1) (Pennica *et al.*, Proc. Nat. Acad. Sci. USA, **92**: 1142-1146 (1995)), catecholamines, adrenocorticosteroids, angiotensin, and prostaglandins.

At present, the treatment of cardiac hypertrophy varies depending on the underlying cardiac disease. Catecholamines, adrenocorticosteroids, angiotensin, prostaglandins, LIF, endothelin (including endothelin-1, -2, and -3 and big endothelin), and CT-1 are among the factors identified as potential mediators of hypertrophy. For example, beta-adrenergic receptor blocking drugs (beta-blockers, *e.g.*, propranolol, timolol, tertalolol, carteolol, nadolol, betaxolol, penbutolol, acetobutolol, atenolol, metoprolol, carvedilol, etc.) and verapamil have been used extensively in the treatment of hypertrophic cardiomyopathy. The beneficial effects of beta-blockers on symptoms (*e.g.*, chest pain) and exercise tolerance are largely due to a decrease in the heart rate with a consequent prolongation of diastole and increased passive ventricular filling. Thompson *et al.*, Br. Heart J., **44**: 488-98 (1980); Harrison *et al.*, Circulation, **29**: 84-98 (1964). Verapamil has been described to improve ventricular filling and probably reducing myocardial ischemia. Bonow *et al.*, Circulation, **72**: 853-64 (1985).

Nifedipine and diltiazem have also been used occasionally in the treatment of hypertrophic cardiomyopathy. Lorell *et al.*, Circulation, **65**: 499-507 (1982); Betocchi *et al.*, Am. J. Cardiol., **78**: 451-457 (1996). However, because of its potent vasodilating properties, nifedipine may be harmful, especially in patients with outflow obstruction. Disopyramide has been used to relieve symptoms by virtue of its negative inotropic properties. Pollick, N. Engl. J. Med., **307**: 997-999 (1982). In many patients, however, the initial benefits decrease with time. Wigle *et al.*, Circulation, **92**: 1680-1692 (1995). Antihypertensive drug therapy has been reported to have beneficial effects on cardiac hypertrophy associated with elevated blood pressure. Examples of drugs used in antihypertensive therapy, alone or in combination, are calcium antagonists, *e.g.*, nitrendipine; adrenergic receptor blocking agents, *e.g.*, those listed above; angiotensin converting enzyme (ACE) inhibitors such as quinapril, captopril, enalapril, ramipril, benazepril, fosinopril, and lisinopril; diuretics, *e.g.*, chlorothiazide, hydrochlorothiazide, hydroflumethazide, methylchlorothiazide, benzthiazide, dichlorphenamide, acetazolamide, and indapamide; and calcium channel blockers, *e.g.*, diltiazem, nifedipine, verapamil, and nicardipine.

For example, treatment of hypertension with diltiazem and captopril showed a decrease in left ventricular muscle mass, but the Doppler indices of diastolic function did not normalize. Szlachcic *et al.*, Am. J. Cardiol., **63**: 198-201 (1989); Shahi *et al.*, Lancet, **336**: 458-461 (1990). These findings were interpreted to indicate that excessive amounts of interstitial collagen may remain after regression of left ventricular hypertrophy. Rossi *et al.*, Am. Heart J., **124**: 700-709 (1992). Rossi *et al.*, *supra*, investigated the effect of captopril on the prevention and

regression of myocardial cell hypertrophy and interstitial fibrosis in pressure overload cardiac hypertrophy, in experimental rats.

Agents that increase cardiac contractility directly (ionotropic agents) were initially thought to benefit patients with heart failure because they improved cardiac output in the short term. However, all positive inotropic agents except digoxigenin have been found to result in increased long-term mortality, in spite of short-term improvements in cardiac performance. Massie, Curr. Op. in Cardiology, 12: 209-217 (1997); Reddy *et al.*, Curr. Opin. Cardiol., 12: 233-241 (1997). Beta-adrenergic receptor blockers have recently been advocated for use in heart failure. Evidence from clinical trials suggests that improvements in cardiac function can be achieved without increased mortality, though documented improvements of patient survival have not yet been demonstrated. See also, U.S. Pat. Nos. 5,935,924, 5,624,806; 5,661,122; and 5,610,134 and WO 95/28173 regarding the use of cardiotropin-1 or antagonists thereof, or growth hormone and/or insulin-like growth factor-I in the treatment of CHF. Another treatment modality is heart transplantation, but this is limited by the availability of donor hearts.

Endothelin is a vasoconstricting peptide comprising 21 amino acids, isolated from swine arterial endothelial culture supernatant and structurally determined. Yanagisawa *et al.*, Nature, 332: 411-415 (1988). Endothelin was later found to exhibit various actions, and endothelin antibodies as endothelin antagonists have proven effective in the treatment of myocardial infarction, renal failure, and other diseases. Since endothelin is present in live bodies and exhibits vasoconstricting action, it is expected to be an endogenous factor involved in the regulation of the circulatory system, and may be associated with hypertension, cardiovascular diseases such as myocardial infarction, and renal diseases such as acute renal failure. Endothelin antagonists are described, for example, in U.S. Pat. No. 5,773,414; JP Pat. Publ. 3130299/1991, EP 457,195; EP 460,679; and EP 552,489. A new endothelin B receptor for identifying endothelin receptor antagonists is described in U.S. Pat. No. 5,773,223.

Current therapy for heart failure is primarily directed to using angiotensin-converting enzyme (ACE) inhibitors, such as captopril, and diuretics. These drugs improve hemodynamic profile and exercise tolerance and reduce the incidence of morbidity and mortality in patients with CHF. Kramer *et al.*, Circulation, 67(4): 807-816 (1983); Captopril Multicenter Research Group, J.A.C.C., 2(4): 755-763 (1983); The CONSENSUS Trial Study Group, N. Engl. J. Med., 316(23): 1429-1435 (1987); The SOLVD Investigators, N. Engl. J. Med., 325(5): 293-302 (1991). Further, they are useful in treating hypertension, left ventricular dysfunction, atherosclerotic vascular disease, and diabetic nephropathy. Brown and Vaughan, *supra*. However, despite proven efficacy, response to ACE inhibitors has been limited. For example, while prolonging survival in the setting of heart failure, ACE inhibitors appear to slow the progression towards end-stage heart failure, and substantial numbers of patients on ACE inhibitors have functional class III heart failure.

Moreover, improvement of functional capacity and exercise time is only small and mortality, although reduced, continues to be high. The CONSENSUS Trial Study Group, N. Engl. J. Med., 316(23): 1429-1453 (1987); The SOLVD Investigators, N. Engl. J. Med., 325(5): 293-302 (1991); Cohn *et al.*, N. Engl. J. Med., 325(5): 303-310 (1991); The Captopril-Digoxin Multicenter Research Group, JAMA, 259(4): 539-544 (1988). Hence, ACE inhibitors consistently appear unable to relieve symptoms in more than 60% of heart failure patients and reduce mortality of heart failure only by approximately 15-20%. For further adverse effects, see Brown and Vaughan,

supra.

An alternative to ACE inhibitors is represented by specific AT1 receptor antagonists. Clinical studies are planned to compare the efficacy of these two modalities in the treatment of cardiovascular and renal disease. However, animal model data suggests that the ACE/Ang II pathway, while clearly involved in cardiac hypertrophy, is not the only, or even the primary pathway active in this role. Mouse genetic "knockout" models have been made to test individual components of the pathway. In one such model, the primary cardiac receptor for Ang II, AT sub 1A, has been genetically deleted; these mice do not develop hypertrophy when Ang II is given experimentally (confirming the basic success of the model in eliminating hypertrophy secondary to Ang II). However, when the aorta is constricted in these animals (a model of hypertensive cardiac stress), the hearts still become hypertrophic. This suggests that alternative signaling pathways, not depending on this receptor (AT sub 1A), are activated in hypertension. ACE inhibitors would presumably not be able to inhibit these pathways. See, Harada *et al.*, Circulation, 97: 1952-1959 (1998). See also, Homcy, Circulation, 97: 1890-1892 (1998) regarding the enigma associated with the process and mechanism of cardiac hypertrophy.

About 750,000 patients suffer from acute myocardial infarction (AMI) annually, and approximately one-fourth of all deaths in the United States are due to AMI. In recent years, thrombolytic agents, *e.g.*, streptokinase, urokinase, and in particular tissue plasminogen activator (t-PA) have significantly increased the survival of patients who suffered myocardial infarction. When administered as a continuous intravenous infusion over 1.5 to 4 hours, t-PA produces coronary patency at 90 minutes in 69% to 90% of the treated patients. Topol *et al.*, Am. J. Cardiol., 61: 723-728 (1988); Neuhaus *et al.*, J. Am. Coll. Cardiol., 12: 581-587 (1988); Neuhaus *et al.*, J. Am. Coll. Cardiol., 14: 1566-1569 (1989). The highest patency rates have been reported with high dose or accelerated dosing regimens. Topol, J. Am. Coll. Cardiol., 15: 922-924 (1990). t-PA may also be administered as a single bolus, although due to its relatively short half-life, it is better suited for infusion therapy. Tebbe *et al.*, Am. J. Cardiol., 64: 448-453 (1989). A t-PA variant, specifically designed to have longer half-life and very high fibrin specificity, TNK t-PA (a T103N, N117Q, KHRR(296-299)AAAA t-PA variant, Keyt *et al.*, Proc. Natl. Acad. Sci. USA, 91: 3670-3674 (1994)) is particularly suitable for bolus administration. However, despite all these advances, the long-term prognosis of patient survival depends greatly on the post-infarction monitoring and treatment of the patients, which should include monitoring and treatment of cardiac hypertrophy.

2.3. Growth Factors

Various naturally occurring polypeptides reportedly induce the proliferation of endothelial cells. Among those polypeptides are the basic and acidic fibroblast growth factors (FGF) (Burgess and Maciag, Annual Rev. Biochem., 58: 575 (1989)), platelet-derived endothelial cell growth factor (PD-ECGF) (Ishikawa *et al.*, Nature, 338: 557 (1989)), and vascular endothelial growth factor (VEGF). Leung *et al.*, Science, 246: 1306 (1989); Ferrara and Henzel, Biochem. Biophys. Res. Commun., 161: 851 (1989); Tischer *et al.*, Biochem. Biophys. Res. Commun., 165: 1198 (1989); EP 471,754B granted July 31, 1996.

Media conditioned by cells transfected with the human VEGF (hVEGF) cDNA promoted the proliferation of capillary endothelial cells, whereas control cells did not. Leung *et al.*, Science, 246: 1306 (1989). Several

additional cDNAs were identified in human cDNA libraries that encode 121-, 189-, and 206-amino acid isoforms of hVEGF (also collectively referred to as hVEGF-related proteins). The 121-amino acid protein differs from hVEGF by virtue of the deletion of the 44 amino acids between residues 116 and 159 in hVEGF. The 189-amino acid protein differs from hVEGF by virtue of the insertion of 24 amino acids at residue 116 in hVEGF, and apparently is identical to human vascular permeability factor (hVPF). The 206-amino acid protein differs from hVEGF by virtue of an insertion of 41 amino acids at residue 116 in hVEGF. Houck *et al.*, Mol. Endocrin., **5**: 1806 (1991); Ferrara *et al.*, J. Cell. Biochem., **47**: 211 (1991); Ferrara *et al.*, Endocrine Reviews, **13**: 18 (1992); Keck *et al.*, Science, **246**: 1309 (1989); Connolly *et al.*, J. Biol. Chem., **264**: 20017 (1989); EP 370,989 published May 30, 1990.

It is now well established that angiogenesis, which involves the formation of new blood vessels from preexisting endothelium, is implicated in the pathogenesis of a variety of disorders. These include solid tumors and metastasis, atherosclerosis, retrolental fibroplasia, hemangiomas, chronic inflammation, intraocular neovascular syndromes such as proliferative retinopathies, *e.g.*, diabetic retinopathy, age-related macular degeneration (AMD), neovascular glaucoma, immune rejection of transplanted corneal tissue and other tissues, rheumatoid arthritis, and psoriasis. Folkman *et al.*, J. Biol. Chem., **267**: 10931-10934 (1992); Klagsbrun *et al.*, Annu. Rev. Physiol., **53**: 217-239 (1991); and Garner A., "Vascular diseases", In: Pathobiology of Ocular Disease. A Dynamic Approach, Garner A., Klintworth GK, eds., 2nd Edition (Marcel Dekker, NY, 1994), pp 1625-1710.

In the case of tumor growth, angiogenesis appears to be crucial for the transition from hyperplasia to neoplasia, and for providing nourishment for the growth and metastasis of the tumor. Folkman *et al.*, Nature, **339**: 58 (1989). The neovascularization allows the tumor cells to acquire a growth advantage and proliferative autonomy compared to the normal cells. A tumor usually begins as a single aberrant cell which can proliferate only to a size of a few cubic millimeters due to the distance from available capillary beds, and it can stay 'dormant' without further growth and dissemination for a long period of time. Some tumor cells then switch to the angiogenic phenotype to activate endothelial cells, which proliferate and mature into new capillary blood vessels. These newly formed blood vessels not only allow for continued growth of the primary tumor, but also for the dissemination and recolonization of metastatic tumor cells. Accordingly, a correlation has been observed between density of microvessels in tumor sections and patient survival in breast cancer as well as in several other tumors. Weidner *et al.*, N. Engl. J. Med., **324**: 1-6 (1991); Horak *et al.*, Lancet, **340**: 1120-1124 (1992); Macchiarini *et al.*, Lancet, **340**: 145-146 (1992). The precise mechanisms that control the angiogenic switch is not well understood, but it is believed that neovascularization of tumor mass results from the net balance of a multitude of angiogenesis stimulators and inhibitors (Folkman, 1995, Nat Med **1**(1):27-31).

The search for positive regulators of angiogenesis has yielded many candidates, including aFGF, bFGF, TGF- α , TGF- β , HGF, TNF- α , angiogenin, IL-8, etc. Folkman *et al.*, J.B.C., *supra*, and Klagsbrun *et al.*, *supra*. The negative regulators so far identified include thrombospondin (Good *et al.*, Proc. Natl. Acad. Sci. USA., **87**: 6624-6628 (1990)), the 16-kilodalton N-terminal fragment of prolactin (Clapp *et al.*, Endocrinology, **133**: 1292-1299 (1993)), angiostatin (O'Reilly *et al.*, Cell, **79**: 315-328 (1994)), and endostatin. O'Reilly *et al.*, Cell, **88**: 277-285 (1996).

Work done over the last several years has established the key role of VEGF, not only in stimulating vascular endothelial cell proliferation, but also in inducing vascular permeability and angiogenesis. Ferrara *et al.*, Endocr. Rev., **18**: 4-25 (1997). The finding that the loss of even a single VEGF allele results in embryonic lethality points to an irreplaceable role played by this factor in the development and differentiation of the vascular system. Furthermore, VEGF has been shown to be a key mediator of neovascularization associated with tumors and intraocular disorders. Ferrara *et al.*, Endocr. Rev., *supra*. The VEGF mRNA is overexpressed by the majority of human tumors examined. Berkman *et al.*, J. Clin. Invest., **91**: 153-159 (1993); Brown *et al.*, Human Pathol., **26**: 86-91 (1995); Brown *et al.*, Cancer Res., **53**: 4727-4735 (1993); Mattern *et al.*, Brit. J. Cancer, **73**: 931-934 (1996); Dvorak *et al.*, Am. J. Pathol., **146**: 1029-1039 (1995).

Also, the concentration levels of VEGF in eye fluids are highly correlated to the presence of active proliferation of blood vessels in patients with diabetic and other ischemia-related retinopathies. Aiello *et al.*, N. Engl. J. Med., **331**: 1480-1487 (1994). Furthermore, recent studies have demonstrated the localization of VEGF in choroidal neovascular membranes in patients affected by AMD. Lopez *et al.*, Invest. Ophthalmol. Vis. Sci., **37**: 855-868 (1996).

Anti-VEGF neutralizing antibodies suppress the growth of a variety of human tumor cell lines in nude mice (Kim *et al.*, Nature, **362**: 841-844 (1993); Warren *et al.*, J. Clin. Invest., **95**: 1789-1797 (1995); Borgström *et al.*, Cancer Res., **56**: 4032-4039 (1996); Melnyk *et al.*, Cancer Res., **56**: 921-924 (1996)) and also inhibit intraocular angiogenesis in models of ischemic retinal disorders. Adamis *et al.*, Arch. Ophthalmol., **114**: 66-71 (1996). Therefore, anti-VEGF monoclonal antibodies or other inhibitors of VEGF action are promising candidates for the treatment of solid tumors and various intraocular neovascular disorders. Such antibodies are described, for example, in EP 817,648 published January 14, 1998 and in WO98/45331 and WO98/45332 both published October 15, 1998.

There exist several other growth factors and mitogens, including transforming oncogenes, that are capable of rapidly inducing a complex set of genes to be expressed by certain cells. Lau and Nathans, Molecular Aspects of Cellular Regulation, **6**: 165-202 (1991). These genes, which have been named immediate-early- or early-response genes, are transcriptionally activated within minutes after contact with a growth factor or mitogen, independent of *de novo* protein synthesis. A group of these intermediate-early genes encodes secreted, extracellular proteins that are needed for coordination of complex biological processes such as differentiation and proliferation, regeneration, and wound healing. Ryseck *et al.*, Cell Growth Differ., **2**: 235-233 (1991).

Highly-related proteins that belong to this group include *cef10* (Simmons *et al.*, Proc. Natl. Acad. Sci. USA, **86**: 1178-1182 (1989)), *cyr61*, which is rapidly activated by serum- or platelet-derived growth factor (PDGF) (O'Brien *et al.*, Mol. Cell Biol., **10**: 3569-3577 (1990)), human connective tissue growth factor (CTGF) (Bradham *et al.*, J. Cell. Biol., **114**: 1285-1294 (1991)), which is secreted by human vascular endothelial cells in high levels after activation with transforming growth factor beta (TGF- β), exhibits PDGF-like biological and immunological activities, and competes with PDGF for a particular cell surface receptor, *fisp-12* (Ryseck *et al.*, Cell Growth Differ., **2**: 235-233 (1991)), human vascular IBP-like growth factor (VIGF) (WO 96/17931), and *nov*, normally arrested in adult kidney cells, which was found to be overexpressed in myeloblastosis-associated-virus-type-1-induced nephroblastomas. Joliet *et al.*, Mol. Cell. Biol., **12**: 10-21 (1992).

The expression of these immediate-early genes acts as "third messengers" in the cascade of events triggered by growth factors. It is also thought that they are needed to integrate and coordinate complex biological processes, such as differentiation and wound healing in which cell proliferation is a common event.

As additional mitogens, insulin-like growth factor binding proteins (IGFBPs) have been shown, in complex with insulin-like growth factor (IGF), to stimulate increased binding of IGF to fibroblast and smooth muscle cell surface receptors. Clemmons *et al.*, *J. Clin. Invest.*, **77**: 1548 (1986). Inhibitory effects of IGFBP on various IGF actions *in vitro* include stimulation of glucose transport by adipocytes, sulfate incorporation by chondrocytes, and thymidine incorporation in fibroblast. Zapf *et al.*, *J. Clin. Invest.*, **63**: 1077 (1979). In addition, inhibitory effects of IGFBPs on growth factor-mediated mitogen activity in normal cells have been shown.

2.4. Need for Further Treatments

In view of the role of vascular endothelial cell growth and angiogenesis in many diseases and disorders, it is desirable to have a means of reducing or inhibiting one or more of the biological effects causing these processes. It is also desirable to have a means of assaying for the presence of pathogenic polypeptides in normal and diseased conditions, and especially cancer. Further, in a specific aspect, as there is no generally applicable therapy for the treatment of cardiac hypertrophy, the identification of factors that can prevent or reduce cardiac myocyte hypertrophy is of primary importance in the development of new therapeutic strategies to inhibit pathophysiological cardiac growth. While there are several treatment modalities for various cardiovascular and oncologic disorders, there is still a need for additional therapeutic approaches.

3. Summary of the Invention

The present invention provides compositions and methods for modulating (*e.g.*, promoting or inhibiting) angiogenesis and/or cardiovascularization in mammals. The present invention is based on the identification of compounds (*i.e.*, proteins) that test positive in various cardiovascular assays that test modulation (*e.g.*, promotion or inhibition) of certain biological activities. Accordingly, the compounds are believed to be useful drugs and/or drug components for the diagnosis and/or treatment (including prevention and amelioration) of disorders where such effects are desired, such as the promotion or inhibition of angiogenesis, inhibition or stimulation of vascular endothelial cell growth, stimulation of growth or proliferation of vascular endothelial cells, inhibition of tumor growth, inhibition of angiogenesis-dependent tissue growth, stimulation of angiogenesis-dependent tissue growth, inhibition of cardiac hypertrophy and stimulation of cardiac hypertrophy, *e.g.*, for the treatment of congestive heart failure. In addition, the compositions and methods of the invention provide for the diagnostic monitoring of patients undergoing clinical evaluation for the treatment of angiogenesis-related disorders, for monitoring the efficacy of compounds in clinical trials and for identifying subjects who may be predisposed to such angiogenic-related disorders.

In one embodiment, the present invention provides a composition comprising a PRO polypeptide, an agonist or antagonist thereof, or an anti-PRO antibody in admixture with a pharmaceutically acceptable carrier. In one aspect, the composition comprises a therapeutically effective amount of the polypeptide, agonist, antagonist

or antibody. In another aspect, the composition comprises a further active ingredient, namely, a cardiovascular, endothelial or angiogenic agent or an angiostatic agent, preferably an angiogenic or angiostatic agent. Preferably, the composition is sterile. The PRO polypeptide, agonist, antagonist or antibody may be administered in the form of a liquid pharmaceutical formulation, which may be preserved to achieve extended storage stability. Preserved liquid pharmaceutical formulations might contain multiple doses of PRO polypeptide, agonist, antagonist or antibody, and might, therefore, be suitable for repeated use. In a preferred embodiment, where the composition comprises an antibody, the antibody is a monoclonal antibody, an antibody fragment, a humanized antibody, or a single-chain antibody.

In a further embodiment, the present invention provides a method for preparing such a composition useful for the treatment of a cardiovascular, endothelial or angiogenic disorder comprising admixing a therapeutically effective amount of a PRO polypeptide, agonist, antagonist or antibody with a pharmaceutically acceptable carrier.

In a still further aspect, the present invention provides an article of manufacture comprising:

- (a) a composition of matter comprising a PRO polypeptide or agonist or antagonist thereof;
- (b) a container containing said composition; and
- (c) a label affixed to said container, or a package insert included in said container referring to the use of said PRO polypeptide or agonist or antagonist thereof in the treatment of a cardiovascular, endothelial or angiogenic disorder, wherein the agonist or antagonist may be an antibody which binds to the PRO polypeptide. The composition may comprise a therapeutically effective amount of the PRO polypeptide or the agonist or antagonist thereof.

In another embodiment, the present invention provides a method for identifying an agonist of a PRO polypeptide comprising:

- (a) contacting cells and a test compound to be screened under conditions suitable for the induction of a cellular response normally induced by a PRO polypeptide; and
- (b) determining the induction of said cellular response to determine if the test compound is an effective agonist, wherein the induction of said cellular response is indicative of said test compound being an effective agonist.

In another embodiment, the present invention provides a method for identifying an agonist of a PRO polypeptide comprising:

- (a) contacting cells and a test compound to be screened under conditions suitable for the stimulation of cell proliferation by a PRO polypeptide; and
- (b) measuring the proliferation of said cells to determine if the test compound is an effective agonist, wherein the stimulation of cell proliferation is indicative of said test compound being an effective agonist.

In another embodiment, the invention provides a method for identifying a compound that inhibits the activity of a PRO polypeptide comprising contacting a test compound with a PRO polypeptide under conditions and for a time sufficient to allow the test compound and polypeptide to interact and determining whether the activity of the PRO polypeptide is inhibited. In a specific preferred aspect, either the test compound or the PRO polypeptide is immobilized on a solid support. In another preferred aspect, the non-immobilized component carries a detectable

label. In a preferred aspect, this method comprises the steps of:

(a) contacting cells and a test compound to be screened in the presence of a PRO polypeptide under conditions suitable for the induction of a cellular response normally induced by a PRO polypeptide; and

5 (b) determining the induction of said cellular response to determine if the test compound is an effective antagonist.

In another preferred aspect, this process comprises the steps of:

(a) contacting cells and a test compound to be screened in the presence of a PRO polypeptide under conditions suitable for the stimulation of cell proliferation by a PRO polypeptide; and

(b) measuring the proliferation of the cells to determine if the test compound is an effective antagonist.

10 In another embodiment, the invention provides a method for identifying a compound that inhibits the expression of a PRO polypeptide in cells that normally expresses the polypeptide, wherein the method comprises contacting the cells with a test compound and determining whether the expression of the PRO polypeptide is inhibited. In a preferred aspect, this method comprises the steps of:

15 (a) contacting cells and a test compound to be screened under conditions suitable for allowing expression of the PRO polypeptide; and

(b) determining the inhibition of expression of said polypeptide.

In a still further embodiment, the invention provides a compound that inhibits the expression of a PRO polypeptide, such as a compound that is identified by the methods set forth above.

20 Another aspect of the present invention is directed to an agonist or an antagonist of a PRO polypeptide which may optionally be identified by the methods described above.

One type of antagonist of a PRO polypeptide that inhibits one or more of the functions or activities of the PRO polypeptide is an antibody. Hence, in another aspect, the invention provides an isolated antibody that binds a PRO polypeptide. In a preferred aspect, the antibody is a monoclonal antibody, which preferably has non-human complementarity-determining-region (CDR) residues and human framework-region (FR) residues. The antibody
25 may be labeled and may be immobilized on a solid support. In a further aspect, the antibody is an antibody fragment, a single-chain antibody, or a humanized antibody. Preferably, the antibody specifically binds to the polypeptide.

In a still further aspect, the present invention provides a method for diagnosing a disease or susceptibility to a disease which is related to a mutation in a PRO polypeptide-encoding nucleic acid sequence comprising
30 determining the presence or absence of said mutation in the PRO polypeptide nucleic acid sequence, wherein the presence or absence of said mutation is indicative of the presence of said disease or susceptibility to said disease.

In a still further aspect, the invention provides a method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal which comprises analyzing the level of expression of a gene encoding a PRO polypeptide (a) in a test sample of tissue cells obtained from said mammal, and (b) in a control sample of known
35 normal tissue cells of the same cell type, wherein a higher or lower expression level in the test sample as compared to the control sample is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in said mammal. The expression of a gene encoding a PRO polypeptide may optionally be accomplished by measuring

the level of mRNA or the polypeptide in the test sample as compared to the control sample.

In a still further aspect, the present invention provides a method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal which comprises detecting the presence or absence of a PRO polypeptide in a test sample of tissue cells obtained from said mammal, wherein the presence or absence of said PRO polypeptide in said test sample is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in said mammal.

In a still further embodiment, the invention provides a method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal comprising (a) contacting an anti-PRO antibody with a test sample of tissue cells obtained from the mammal, and (b) detecting the formation of a complex between the antibody and the PRO polypeptide in the test sample, wherein the formation of said complex is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in the mammal. The detection may be qualitative or quantitative, and may be performed in comparison with monitoring the complex formation in a control sample of known normal tissue cells of the same cell type. A larger or smaller quantity of complexes formed in the test sample indicates the presence of a cardiovascular, endothelial or angiogenic dysfunction in the mammal from which the test tissue cells were obtained. The antibody preferably carries a detectable label. Complex formation can be monitored, for example, by light microscopy, flow cytometry, fluorimetry, or other techniques known in the art. The test sample is usually obtained from an individual suspected to have a cardiovascular, endothelial or angiogenic disorder.

In another embodiment, the invention provides a method for determining the presence of a PRO polypeptide in a sample comprising exposing a sample suspected of containing the PRO polypeptide to an anti-PRO antibody and determining binding of said antibody to a component of said sample. In a specific aspect, the sample comprises a cell suspected of containing the PRO polypeptide and the antibody binds to the cell. The antibody is preferably detectably labeled and/or bound to a solid support.

In further aspects, the invention provides a cardiovascular, endothelial or angiogenic disorder diagnostic kit comprising an anti-PRO antibody and a carrier in suitable packaging. Preferably, such kit further comprises instructions for using said antibody to detect the presence of the PRO polypeptide. Preferably, the carrier is a buffer, for example. Preferably, the cardiovascular, endothelial or angiogenic disorder is cancer.

In yet another embodiment, the present invention provides a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal an effective amount of a PRO polypeptide. Preferably, the disorder is cardiac hypertrophy, trauma such as wounds or burns, or a type of cancer. In a further aspect, the mammal is further exposed to angioplasty or a drug that treats cardiovascular, endothelial or angiogenic disorders such as ACE inhibitors or chemotherapeutic agents if the cardiovascular, endothelial or angiogenic disorder is a type of cancer. Preferably, the mammal is human, preferably one who is at risk of developing cardiac hypertrophy and more preferably has suffered myocardial infarction.

In another preferred aspect, the cardiac hypertrophy is characterized by the presence of an elevated level of $\text{PGF}_{2\alpha}$. Alternatively, the cardiac hypertrophy may be induced by myocardial infarction, wherein preferably the administration of the PRO polypeptide is initiated within 48 hours, more preferably within 24 hours, following myocardial infarction.

In another preferred embodiment, the cardiovascular, endothelial or angiogenic disorder is cardiac hypertrophy and said PRO polypeptide is administered together with a cardiovascular, endothelial or angiogenic agent. The preferred cardiovascular, endothelial or angiogenic agent for this purpose is selected from the group consisting of an antihypertensive drug, an ACE inhibitor, an endothelin receptor antagonist and a thrombolytic agent. If a thrombolytic agent is administered, preferably the PRO polypeptide is administered following administration of such agent. More preferably, the thrombolytic agent is recombinant human tissue plasminogen activator.

In another preferred aspect, the cardiovascular, endothelial or angiogenic disorder is cardiac hypertrophy and the PRO polypeptide is administered following primary angioplasty for the treatment of acute myocardial infarction, preferably wherein the mammal is further exposed to angioplasty or a cardiovascular, endothelial, or angiogenic agent.

In another preferred embodiment, the cardiovascular, endothelial or angiogenic disorder is a cancer and the PRO polypeptide is administered in combination with a chemotherapeutic agent, a growth inhibitory agent or a cytotoxic agent.

In a further embodiment, the invention provides a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal an effective amount of a PRO polypeptide agonist, antagonist or anti-PRO antibody. Preferably, the cardiovascular, endothelial or angiogenic disorder is cardiac hypertrophy, trauma, a cancer, or age-related macular degeneration. Also preferred is where the mammal is human, and where an effective amount of an angiogenic or angiostatic agent is administered in conjunction with the agonist, antagonist or anti-PRO antibody.

In still further embodiments, the invention provides a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal that suffers therefrom comprising administering to the mammal a nucleic acid molecule that codes for either (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide or (c) an antagonist of a PRO polypeptide, wherein said agonist or antagonist may be an anti-PRO antibody. In a preferred embodiment, the mammal is human. In another preferred embodiment, the gene is administered via *ex vivo* gene therapy. In a further preferred embodiment, the gene is comprised within a vector, more preferably an adenoviral, adeno-associated viral, lentiviral, or retroviral vector.

In yet another aspect, the invention provides a recombinant retroviral particle comprising a retroviral vector consisting essentially of a promoter, nucleic acid encoding (a) a PRO polypeptide, (b) an agonist polypeptide of a PRO polypeptide, or (c) an antagonist polypeptide of a PRO polypeptide, and a signal sequence for cellular secretion of the polypeptide, wherein the retroviral vector is in association with retroviral structural proteins. Preferably, the signal sequence is from a mammal, such as from a native PRO polypeptide.

In a still further embodiment, the invention supplies an *ex vivo* producer cell comprising a nucleic acid construct that expresses retroviral structural proteins and also comprises a retroviral vector consisting essentially of a promoter, nucleic acid encoding (a) a PRO polypeptide, (b) an agonist polypeptide of a PRO polypeptide or (c) an antagonist polypeptide of a PRO polypeptide, and a signal sequence for cellular secretion of the polypeptide, wherein said producer cell packages the retroviral vector in association with the structural proteins to produce

recombinant retroviral particles.

5 In yet another embodiment, the invention provides a method for inhibiting endothelial cell growth in a mammal comprising administering to the mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein endothelial cell growth in said mammal is inhibited, and wherein said agonist or antagonist may be an anti-PRO antibody. Preferably, the mammal is human and the endothelial cell growth is associated with a tumor or a retinal disorder.

10 In yet another embodiment, the invention provides a method for stimulating endothelial cell growth in a mammal comprising administering to the mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein endothelial cell growth in said mammal is stimulated, and wherein said agonist or antagonist may be an anti-PRO antibody. Preferably, the mammal is human.

15 In yet another embodiment, the invention provides a method for inhibiting cardiac hypertrophy in a mammal comprising administering to the mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein cardiac hypertrophy in said mammal is inhibited, and wherein said agonist or antagonist may be an anti-PRO antibody. Preferably, the mammal is human and the cardiac hypertrophy has been induced by myocardial infarction.

20 In yet another embodiment, the invention provides a method for stimulating cardiac hypertrophy in a mammal comprising administering to the mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein cardiac hypertrophy in said mammal is stimulated, and wherein said agonist or antagonist may be an anti-PRO antibody. Preferably, the mammal is human who suffers from congestive heart failure.

In yet another embodiment, the invention provides a method for inhibiting angiogenesis induced by a PRO polypeptide in a mammal comprising administering a therapeutically effective amount of an anti-PRO antibody to the mammal. Preferably, the mammal is a human, and more preferably the mammal has a tumor or a retinal disorder.

25 In yet another embodiment, the invention provides a method for stimulating angiogenesis induced by a PRO polypeptide in a mammal comprising administering a therapeutically effective amount of a PRO polypeptide to the mammal. Preferably, the mammal is a human, and more preferably angiogenesis would promote tissue regeneration or wound healing.

30 In yet another embodiment, the invention provides a method for modulating (*e.g.*, inhibiting or stimulating) endothelial cell growth in a mammal comprising administering to the mammal a PRO21, PRO181, PRO205, PRO214, PRO221, PRO229, PRO231, PRO238, PRO241, PRO247, PRO256, PRO258, PRO263, PRO265, PRO295, PRO321, PRO322, PRO337, PRO363, PRO365, PRO444, PRO533, PRO697, PRO720, PRO725, PRO771, PRO788, PRO791, PRO819, PRO827, PRO828, PRO836, PRO846, PRO865, PRO1005, PRO1006, PRO1007, PRO1025, PRO1029, PRO1054, PRO1071, PRO1075, PRO1079, PRO1080, PRO1114, PRO1131, 35 PRO1155, PRO1160, PRO1184, PRO1186, PRO1190, PRO1192, PRO1195, PRO1244, PRO1272, PRO1273, PRO1274, PRO1279, PRO1283, PRO1286, PRO1306, PRO1309, PRO1325, PRO1329, PRO1347, PRO1356, PRO1376, PRO1382, PRO1411, PRO1412, PRO1419, PRO1474, PRO1477, PRO1488, PRO1508, PRO1550,

PRO1556, PRO1760, PRO1782, PRO1787, PRO1801, PRO1868, PRO1887, PRO1890, PRO3438, PRO3444, PRO4302, PRO4324, PRO4333, PRO4341, PRO4342, PRO4353, PRO4354, PRO4356, PRO4371, PRO4405, PRO4408, PRO4422, PRO4425, PRO4499, PRO5723, PRO5725, PRO5737, PRO5776, PRO6006, PRO6029, PRO6071, PRO7436, PRO9771, PRO9821, PRO9873, PRO10008, PRO10096, PRO19670, PRO20040, PRO20044, PRO21055, PRO21384 or PRO28631 polypeptide, agonist or antagonist thereof, wherein endothelial cell growth in said mammal is modulated.

In yet another embodiment, the invention provides a method for modulating (*e.g.*, inhibiting or stimulating) smooth muscle cell growth in a mammal comprising administering to the mammal a PRO162, PRO181, PRO182, PRO195, PRO204, PRO221, PRO230, PRO256, PRO258, PRO533, PRO697, PRO725, PRO738, PRO826, PRO836, PRO840, PRO846, PRO865, PRO982, PRO1025, PRO1029, PRO1071, PRO1080, PRO1083, PRO1134, PRO1160, PRO1182, PRO1184, PRO1186, PRO1192, PRO1265, PRO1274, PRO1279, PRO1283, PRO1306, PRO1308, PRO1309, PRO1325, PRO1337, PRO1338, PRO1343, PRO1376, PRO1387, PRO1411, PRO1412, PRO1415, PRO1434, PRO1474, PRO1488, PRO1550, PRO1556, PRO1567, PRO1600, PRO1754, PRO1758, PRO1760, PRO1787, PRO1865, PRO1868, PRO1917, PRO1928, PRO3438, PRO3562, PRO4302, PRO4333, PRO4345, PRO4353, PRO4354, PRO4405, PRO4408, PRO4430, PRO4503, PRO5725, PRO6714, PRO9771, PRO9820, PRO9940, PRO10096, PRO21055, PRO21184 or PRO21366 polypeptide, agonist or antagonist thereof, wherein endothelial cell growth in said mammal is modulated.

In yet another embodiment, the invention provides a method for modulating (*e.g.*, inducing or reducing) cardiac hypertrophy in a mammal comprising administering to the mammal a PRO21 polypeptide, agonist or antagonist thereof, wherein cardiac hypertrophy in said mammal is modulated.

In yet another embodiment, the invention provides a method for modulating (*e.g.*, inducing or reducing) endothelial cell apoptosis in a mammal comprising administering to the mammal a PRO4302 polypeptide, agonist or antagonist thereof, wherein cardiac hypertrophy in said mammal is modulated.

In yet another embodiment, the invention provides a method for modulating (*e.g.*, stimulating or inhibiting) angiogenesis in a mammal comprising administering a therapeutically effective amount of a PRO1376 or PRO1449 polypeptide, agonist or antagonist thereof to the mammal, wherein said angiogenesis is modulated.

In yet another embodiment, the invention provides a method for modulating (*e.g.*, inducing or reducing) angiogenesis by modulating (*e.g.*, inducing or reducing) endothelial cell tube formation in a mammal comprising administering to the mammal a PRO178, PRO195, PRO228, PRO301, PRO302, PRO532, PRO724, PRO730, PRO734, PRO793, PRO871, PRO938, PRO1012, PRO1120, PRO1139, PRO1198, PRO1287, PRO1361, PRO1864, PRO1873, PRO2010, PRO3579, PRO4313, PRO4527, PRO4538, PRO4553, PRO4995, PRO5730, PRO6008, PRO7223, PRO7248 or PRO7261 polypeptide, agonist or antagonist thereof, wherein endothelial cell tube formation in said mammal is modulated.

In other embodiments of the present invention, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence that encodes a PRO polypeptide.

In one aspect, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98%

nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule encoding a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

In other aspects, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule comprising the coding sequence of a full-length PRO polypeptide cDNA as disclosed herein, the coding sequence of a PRO polypeptide lacking the signal peptide as disclosed herein, the coding sequence of an extracellular domain of a transmembrane PRO polypeptide, with or without the signal peptide, as disclosed herein or the coding sequence of any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

In a further aspect, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence having at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule that encodes the same mature polypeptide encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein, or (b) the complement of the DNA molecule of (a).

Another aspect of the present invention provides an isolated nucleic acid molecule comprising a nucleotide sequence encoding a PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated, or is complementary to such encoding nucleotide sequence, wherein the transmembrane domain(s) of such polypeptide are disclosed herein. Therefore, soluble extracellular domains of the herein described PRO polypeptides are contemplated.

Another embodiment is directed to fragments of a PRO polypeptide coding sequence, or the complement thereof, that may find use as, for example, hybridization probes, for encoding fragments of a PRO polypeptide that may optionally encode a polypeptide comprising a binding site for an anti-PRO antibody or as antisense oligonucleotide probes. Such nucleic acid fragments are usually at least about 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 350, 400, 450, 500, 600, 700 or 800 nucleotides in length and alternatively at least about 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length. It is noted that novel fragments of a PRO polypeptide-encoding nucleotide sequence may be determined in a routine manner by aligning the PRO polypeptide-encoding nucleotide sequence with other known nucleotide sequences using any of a number of well known sequence alignment programs and determining which PRO polypeptide-encoding nucleotide sequence fragment(s) are novel. All of such PRO polypeptide-encoding nucleotide sequences are contemplated herein. Also contemplated are the PRO polypeptide fragments encoded by these nucleotide molecule fragments, preferably those PRO polypeptide fragments that comprise a binding site for an anti-PRO antibody.

In another embodiment, the invention provides an isolated PRO polypeptide encoded by any of the isolated

nucleic acid sequences hereinabove identified.

5 In a certain aspect, the invention provides an isolated PRO polypeptide comprising an amino acid sequence having at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein.

10 In a further aspect, the invention provides an isolated PRO polypeptide comprising an amino acid sequence having at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to an amino acid sequence encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein.

15 In a specific aspect, the invention provides an isolated PRO polypeptide without the N-terminal signal sequence and/or the initiating methionine and that is encoded by a nucleotide sequence that encodes such an amino acid sequence as hereinbefore described. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

20 Another aspect of the invention provides an isolated PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

25 In yet another embodiment, the invention provides agonists and antagonists of a native PRO polypeptide as defined herein. In a particular embodiment, the agonist or antagonist is an anti-PRO antibody or a small molecule.

30 In a further embodiment, the invention provides a method of identifying agonists or antagonists to a PRO polypeptide which comprise contacting the PRO polypeptide with a candidate molecule and monitoring a biological activity mediated by said PRO polypeptide. Preferably, the PRO polypeptide is a native PRO polypeptide.

In a still further embodiment, the invention provides a composition of matter comprising a PRO polypeptide, or an agonist or antagonist of a PRO polypeptide as herein described, or an anti-PRO antibody, in combination with a carrier. Optionally, the carrier is a pharmaceutically acceptable carrier.

35 Another embodiment of the present invention is directed to the use of a PRO polypeptide, or an agonist or antagonist thereof as hereinbefore described, or an anti-PRO antibody, for the preparation of a medicament useful in the treatment of a condition which is responsive to the PRO polypeptide, an agonist or antagonist thereof or an anti-PRO antibody.

In additional embodiments of the present invention, the invention provides vectors comprising DNA encoding any of the herein described polypeptides. Host cells comprising any such vector are also provided. By way of example, the host cells may be CHO cells, *E. coli*, yeast, or Baculovirus-infected insect cells. A process for producing any of the herein described polypeptides is further provided and comprises culturing host cells under conditions suitable for expression of the desired polypeptide and recovering the desired polypeptide from the cell culture.

In other embodiments, the invention provides chimeric molecules comprising any of the herein described polypeptides fused to a heterologous polypeptide or amino acid sequence. Example of such chimeric molecules comprise any of the herein described polypeptides fused to an epitope tag sequence or a Fc region of an immunoglobulin.

In yet another embodiment, the invention provides an antibody which specifically binds to any of the above or below described polypeptides. Optionally, the antibody is a monoclonal antibody, humanized antibody, antibody fragment or single-chain antibody.

In yet other embodiments, the invention provides oligonucleotide probes useful for isolating genomic and cDNA nucleotide sequences or as antisense probes, wherein those probes may be derived from any of the above or below described nucleotide sequences.

4. Brief Description of the Drawings

Figure 1 shows a nucleotide sequence (SEQ ID NO:1) of a native sequence PRO181 cDNA, wherein SEQ ID NO:1 is a clone designated herein as "DNA23330-1390".

Figure 2 shows the amino acid sequence (SEQ ID NO:2) derived from the coding sequence of SEQ ID NO:1 shown in Figure 1.

Figure 3 shows a nucleotide sequence (SEQ ID NO:3) of a native sequence PRO178 cDNA, wherein SEQ ID NO:3 is a clone designated herein as "DNA23339-1130".

Figure 4 shows the amino acid sequence (SEQ ID NO:4) derived from the coding sequence of SEQ ID NO:3 shown in Figure 3.

Figure 5 shows a nucleotide sequence (SEQ ID NO:5) of a native sequence PRO444 cDNA, wherein SEQ ID NO:5 is a clone designated herein as "DNA26846-1397".

Figure 6 shows the amino acid sequence (SEQ ID NO:6) derived from the coding sequence of SEQ ID NO:5 shown in Figure 5.

Figure 7 shows a nucleotide sequence (SEQ ID NO:7) of a native sequence PRO195 cDNA, wherein SEQ ID NO:7 is a clone designated herein as "DNA26847-1395".

Figure 8 shows the amino acid sequence (SEQ ID NO:8) derived from the coding sequence of SEQ ID NO:7 shown in Figure 7.

Figure 9 shows a nucleotide sequence (SEQ ID NO:9) of a native sequence PRO182 cDNA, wherein SEQ ID NO:9 is a clone designated herein as "DNA27865-1091".

Figure 10 shows the amino acid sequence (SEQ ID NO:10) derived from the coding sequence of SEQ ID

NO:9 shown in Figure 9.

Figure 11 shows a nucleotide sequence (SEQ ID NO:11) of a native sequence PRO205 cDNA, wherein SEQ ID NO:11 is a clone designated herein as "DNA30868-1156".

5 Figure 12 shows the amino acid sequence (SEQ ID NO:12) derived from the coding sequence of SEQ ID NO:11 shown in Figure 11.

Figure 13 shows a nucleotide sequence (SEQ ID NO:13) of a native sequence PRO204 cDNA, wherein SEQ ID NO:13 is a clone designated herein as "DNA30871-1157".

Figure 14 shows the amino acid sequence (SEQ ID NO:14) derived from the coding sequence of SEQ ID NO:13 shown in Figure 13.

10 Figure 15 shows a nucleotide sequence (SEQ ID NO:15) of a native sequence PRO1873 cDNA, wherein SEQ ID NO:15 is a clone designated herein as "DNA30880".

Figure 16 shows the amino acid sequence (SEQ ID NO:16) derived from the coding sequence of SEQ ID NO:15 shown in Figure 15.

15 Figure 17 shows a nucleotide sequence (SEQ ID NO:17) of a native sequence PRO214 cDNA, wherein SEQ ID NO:17 is a clone designated herein as "DNA32286-1191".

Figure 18 shows the amino acid sequence (SEQ ID NO:18) derived from the coding sequence of SEQ ID NO:17 shown in Figure 17.

Figure 19 shows a nucleotide sequence (SEQ ID NO:19) of a native sequence PRO221 cDNA, wherein SEQ ID NO:19 is a clone designated herein as "DNA33089-1132".

20 Figure 20 shows the amino acid sequence (SEQ ID NO:20) derived from the coding sequence of SEQ ID NO:19 shown in Figure 19.

Figure 21 shows a nucleotide sequence (SEQ ID NO:21) of a native sequence PRO228 cDNA, wherein SEQ ID NO:21 is a clone designated herein as "DNA33092-1202".

25 Figure 22 shows the amino acid sequence (SEQ ID NO:22) derived from the coding sequence of SEQ ID NO:21 shown in Figure 21.

Figure 23 shows a nucleotide sequence (SEQ ID NO:23) of a native sequence PRO229 cDNA, wherein SEQ ID NO:23 is a clone designated herein as "DNA33100-1159".

Figure 24 shows the amino acid sequence (SEQ ID NO:24) derived from the coding sequence of SEQ ID NO:23 shown in Figure 23.

30 Figure 25 shows a nucleotide sequence (SEQ ID NO:25) of a native sequence PRO230 cDNA, wherein SEQ ID NO:25 is a clone designated herein as "DNA33223-1136".

Figure 26 shows the amino acid sequence (SEQ ID NO:26) derived from the coding sequence of SEQ ID NO:25 shown in Figure 25.

35 Figure 27 shows a nucleotide sequence (SEQ ID NO:27) of a native sequence PRO7223 cDNA, wherein SEQ ID NO:27 is a clone designated herein as "DNA34385".

Figure 28 shows the amino acid sequence (SEQ ID NO:28) derived from the coding sequence of SEQ ID NO:27 shown in Figure 27.

Figure 29 shows a nucleotide sequence (SEQ ID NO:29) of a native sequence PRO241 cDNA, wherein SEQ ID NO:29 is a clone designated herein as "DNA34392-1170".

Figure 30 shows the amino acid sequence (SEQ ID NO:30) derived from the coding sequence of SEQ ID NO:29 shown in Figure 29.

5 Figure 31 shows a nucleotide sequence (SEQ ID NO:31) of a native sequence PRO263 cDNA, wherein SEQ ID NO:31 is a clone designated herein as "DNA34431-1177".

Figure 32 shows the amino acid sequence (SEQ ID NO:32) derived from the coding sequence of SEQ ID NO:31 shown in Figure 31.

10 Figure 33 shows a nucleotide sequence (SEQ ID NO:33) of a native sequence PRO321 cDNA, wherein SEQ ID NO:33 is a clone designated herein as "DNA34433-1308".

Figure 34 shows the amino acid sequence (SEQ ID NO:34) derived from the coding sequence of SEQ ID NO:33 shown in Figure 33.

Figure 35 shows a nucleotide sequence (SEQ ID NO:35) of a native sequence PRO231 cDNA, wherein SEQ ID NO:35 is a clone designated herein as "DNA34434-1139".

15 Figure 36 shows the amino acid sequence (SEQ ID NO:36) derived from the coding sequence of SEQ ID NO:35 shown in Figure 35.

Figure 37 shows a nucleotide sequence (SEQ ID NO:37) of a native sequence PRO238 cDNA, wherein SEQ ID NO:37 is a clone designated herein as "DNA35600-1162".

20 Figure 38 shows the amino acid sequence (SEQ ID NO:38) derived from the coding sequence of SEQ ID NO:37 shown in Figure 37.

Figure 39 shows a nucleotide sequence (SEQ ID NO:39) of a native sequence PRO247 cDNA, wherein SEQ ID NO:39 is a clone designated herein as "DNA35673-1201".

Figure 40 shows the amino acid sequence (SEQ ID NO:40) derived from the coding sequence of SEQ ID NO:39 shown in Figure 39.

25 Figure 41 shows a nucleotide sequence (SEQ ID NO:41) of a native sequence PRO256 cDNA, wherein SEQ ID NO:41 is a clone designated herein as "DNA35880-1160".

Figure 42 shows the amino acid sequence (SEQ ID NO:42) derived from the coding sequence of SEQ ID NO:41 shown in Figure 41.

30 Figure 43 shows a nucleotide sequence (SEQ ID NO:43) of a native sequence PRO258 cDNA, wherein SEQ ID NO:43 is a clone designated herein as "DNA35918-1174".

Figure 44 shows the amino acid sequence (SEQ ID NO:44) derived from the coding sequence of SEQ ID NO:43 shown in Figure 43.

Figure 45 shows a nucleotide sequence (SEQ ID NO:45) of a native sequence PRO265 cDNA, wherein SEQ ID NO:45 is a clone designated herein as "DNA36350-1158".

35 Figure 46 shows the amino acid sequence (SEQ ID NO:46) derived from the coding sequence of SEQ ID NO:45 shown in Figure 45.

Figure 47 shows a nucleotide sequence (SEQ ID NO:47) of a native sequence PRO21 cDNA, wherein SEQ

ID NO:47 is a clone designated herein as "DNA36638-1056".

Figure 48 shows the amino acid sequence (SEQ ID NO:48) derived from the coding sequence of SEQ ID NO:47 shown in Figure 47.

5 Figure 49 shows a nucleotide sequence (SEQ ID NO:49) of a native sequence PRO295 cDNA, wherein SEQ ID NO:49 is a clone designated herein as "DNA38268-1188".

Figure 50 shows the amino acid sequence (SEQ ID NO:50) derived from the coding sequence of SEQ ID NO:49 shown in Figure 49.

Figure 51 shows a nucleotide sequence (SEQ ID NO:51) of a native sequence PRO302 cDNA, wherein SEQ ID NO:51 is a clone designated herein as "DNA40370-1217".

10 Figure 52 shows the amino acid sequence (SEQ ID NO:52) derived from the coding sequence of SEQ ID NO:51 shown in Figure 51.

Figure 53 shows a nucleotide sequence (SEQ ID NO:53) of a native sequence PRO301 cDNA, wherein SEQ ID NO:53 is a clone designated herein as "DNA40628-1216".

15 Figure 54 shows the amino acid sequence (SEQ ID NO:54) derived from the coding sequence of SEQ ID NO:53 shown in Figure 53.

Figure 55 shows a nucleotide sequence (SEQ ID NO:55) of a native sequence PRO337 cDNA, wherein SEQ ID NO:55 is a clone designated herein as "DNA43316-1237".

Figure 56 shows the amino acid sequence (SEQ ID NO:56) derived from the coding sequence of SEQ ID NO:55 shown in Figure 55.

20 Figure 57 shows a nucleotide sequence (SEQ ID NO:57) of a native sequence PRO7248 cDNA, wherein SEQ ID NO:57 is a clone designated herein as "DNA44195".

Figure 58 shows the amino acid sequence (SEQ ID NO:58) derived from the coding sequence of SEQ ID NO:57 shown in Figure 57.

25 Figure 59 shows a nucleotide sequence (SEQ ID NO:59) of a native sequence PRO846 cDNA, wherein SEQ ID NO:59 is a clone designated herein as "DNA44196-1353".

Figure 60 shows the amino acid sequence (SEQ ID NO:60) derived from the coding sequence of SEQ ID NO:59 shown in Figure 59.

Figure 61 shows a nucleotide sequence (SEQ ID NO:61) of a native sequence PRO1864 cDNA, wherein SEQ ID NO:61 is a clone designated herein as "DNA45409-2511".

30 Figure 62 shows the amino acid sequence (SEQ ID NO:62) derived from the coding sequence of SEQ ID NO:61 shown in Figure 61.

Figure 63 shows a nucleotide sequence (SEQ ID NO:63) of a native sequence PRO363 cDNA, wherein SEQ ID NO:63 is a clone designated herein as "DNA45419-1252".

35 Figure 64 shows the amino acid sequence (SEQ ID NO:64) derived from the coding sequence of SEQ ID NO:63 shown in Figure 63.

Figure 65 shows a nucleotide sequence (SEQ ID NO:65) of a native sequence PRO730 cDNA, wherein SEQ ID NO:65 is a clone designated herein as "DNA45624-1400".

Figure 66 shows the amino acid sequence (SEQ ID NO:66) derived from the coding sequence of SEQ ID NO:65 shown in Figure 65.

Figure 67 shows a nucleotide sequence (SEQ ID NO:67) of a native sequence PRO365 cDNA, wherein SEQ ID NO:67 is a clone designated herein as "DNA46777-1253".

5 Figure 68 shows the amino acid sequence (SEQ ID NO:68) derived from the coding sequence of SEQ ID NO:67 shown in Figure 67.

Figure 69 shows a nucleotide sequence (SEQ ID NO:69) of a native sequence PRO532 cDNA, wherein SEQ ID NO:69 is a clone designated herein as "DNA48335".

10 Figure 70 shows the amino acid sequence (SEQ ID NO:70) derived from the coding sequence of SEQ ID NO:69 shown in Figure 69.

Figure 71 shows a nucleotide sequence (SEQ ID NO:71) of a native sequence PRO322 cDNA, wherein SEQ ID NO:71 is a clone designated herein as "DNA48336-1309".

Figure 72 shows the amino acid sequence (SEQ ID NO:72) derived from the coding sequence of SEQ ID NO:71 shown in Figure 71.

15 Figure 73 shows a nucleotide sequence (SEQ ID NO:73) of a native sequence PRO1120 cDNA, wherein SEQ ID NO:73 is a clone designated herein as "DNA48606-1479".

Figure 74 shows the amino acid sequence (SEQ ID NO:74) derived from the coding sequence of SEQ ID NO:73 shown in Figure 73.

20 Figure 75 shows a nucleotide sequence (SEQ ID NO:75) of a native sequence PRO7261 cDNA, wherein SEQ ID NO:75 is a clone designated herein as "DNA49149".

Figure 76 shows the amino acid sequence (SEQ ID NO:76) derived from the coding sequence of SEQ ID NO:75 shown in Figure 75.

Figure 77 shows a nucleotide sequence (SEQ ID NO:77) of a native sequence PRO533 cDNA, wherein SEQ ID NO:77 is a clone designated herein as "DNA49435-1219".

25 Figure 78 shows the amino acid sequence (SEQ ID NO:78) derived from the coding sequence of SEQ ID NO:77 shown in Figure 77.

Figure 79 shows a nucleotide sequence (SEQ ID NO:79) of a native sequence PRO724 cDNA, wherein SEQ ID NO:79 is a clone designated herein as "DNA49631-1328".

30 Figure 80 shows the amino acid sequence (SEQ ID NO:80) derived from the coding sequence of SEQ ID NO:79 shown in Figure 79.

Figure 81 shows a nucleotide sequence (SEQ ID NO:81) of a native sequence PRO734 cDNA, wherein SEQ ID NO:81 is a clone designated herein as "DNA49817".

Figure 82 shows the amino acid sequence (SEQ ID NO:82) derived from the coding sequence of SEQ ID NO:81 shown in Figure 81.

35 Figure 83 shows a nucleotide sequence (SEQ ID NO:83) of a native sequence PRO771 cDNA, wherein SEQ ID NO:83 is a clone designated herein as "DNA49829-1346".

Figure 84 shows the amino acid sequence (SEQ ID NO:84) derived from the coding sequence of SEQ ID

NO:83 shown in Figure 83.

Figure 85 shows a nucleotide sequence (SEQ ID NO:85) of a native sequence PRO2010 cDNA, wherein SEQ ID NO:85 is a clone designated herein as "DNA50792".

5 Figure 86 shows the amino acid sequence (SEQ ID NO:86) derived from the coding sequence of SEQ ID NO:85 shown in Figure 85.

Figure 87 shows a nucleotide sequence (SEQ ID NO:87) of a native sequence PRO871 cDNA, wherein SEQ ID NO:87 is a clone designated herein as "DNA50919-1361".

Figure 88 shows the amino acid sequence (SEQ ID NO:88) derived from the coding sequence of SEQ ID NO:87 shown in Figure 87.

10 Figure 89 shows a nucleotide sequence (SEQ ID NO:89) of a native sequence PRO697 cDNA, wherein SEQ ID NO:89 is a clone designated herein as "DNA50920-1325".

Figure 90 shows the amino acid sequence (SEQ ID NO:90) derived from the coding sequence of SEQ ID NO:89 shown in Figure 89.

15 Figure 91 shows a nucleotide sequence (SEQ ID NO:91) of a native sequence PRO1083 cDNA, wherein SEQ ID NO:91 is a clone designated herein as "DNA50921-1458".

Figure 92 shows the amino acid sequence (SEQ ID NO:92) derived from the coding sequence of SEQ ID NO:91 shown in Figure 91.

Figure 93 shows a nucleotide sequence (SEQ ID NO:93) of a native sequence PRO725 cDNA, wherein SEQ ID NO:93 is a clone designated herein as "DNA52758-1399".

20 Figure 94 shows the amino acid sequence (SEQ ID NO:94) derived from the coding sequence of SEQ ID NO:93 shown in Figure 93.

Figure 95 shows a nucleotide sequence (SEQ ID NO:95) of a native sequence PRO720 cDNA, wherein SEQ ID NO:95 is a clone designated herein as "DNA53517-1366-1".

25 Figure 96 shows the amino acid sequence (SEQ ID NO:96) derived from the coding sequence of SEQ ID NO:95 shown in Figure 95.

Figure 97 shows a nucleotide sequence (SEQ ID NO:97) of a native sequence PRO738 cDNA, wherein SEQ ID NO:97 is a clone designated herein as "DNA53915-1258".

Figure 98 shows the amino acid sequence (SEQ ID NO:98) derived from the coding sequence of SEQ ID NO:97 shown in Figure 97.

30 Figure 99 shows a nucleotide sequence (SEQ ID NO:99) of a native sequence PRO865 cDNA, wherein SEQ ID NO:99 is a clone designated herein as "DNA53974-1401".

Figure 100 shows the amino acid sequence (SEQ ID NO:100) derived from the coding sequence of SEQ ID NO:99 shown in Figure 99.

35 Figure 101 shows a nucleotide sequence (SEQ ID NO:101) of a native sequence PRO840 cDNA, wherein SEQ ID NO:101 is a clone designated herein as "DNA53987-1438".

Figure 102 shows the amino acid sequence (SEQ ID NO:102) derived from the coding sequence of SEQ ID NO:101 shown in Figure 101.

Figure 103 shows a nucleotide sequence (SEQ ID NO:103) of a native sequence PRO1080 cDNA, wherein SEQ ID NO:103 is a clone designated herein as "DNA56047-1456".

Figure 104 shows the amino acid sequence (SEQ ID NO:104) derived from the coding sequence of SEQ ID NO:103 shown in Figure 103.

5 Figure 105 shows a nucleotide sequence (SEQ ID NO:105) of a native sequence PRO1079 cDNA, wherein SEQ ID NO:105 is a clone designated herein as "DNA56050-1455".

Figure 106 shows the amino acid sequence (SEQ ID NO:106) derived from the coding sequence of SEQ ID NO:105 shown in Figure 105.

10 Figure 107 shows a nucleotide sequence (SEQ ID NO:107) of a native sequence PRO793 cDNA, wherein SEQ ID NO:107 is a clone designated herein as "DNA56110-1437".

Figure 108 shows the amino acid sequence (SEQ ID NO:108) derived from the coding sequence of SEQ ID NO:107 shown in Figure 107.

Figure 109 shows a nucleotide sequence (SEQ ID NO:109) of a native sequence PRO788 cDNA, wherein SEQ ID NO:109 is a clone designated herein as "DNA56405-1357".

15 Figure 110 shows the amino acid sequence (SEQ ID NO:110) derived from the coding sequence of SEQ ID NO:109 shown in Figure 109.

Figure 111 shows a nucleotide sequence (SEQ ID NO:111) of a native sequence PRO938 cDNA, wherein SEQ ID NO:111 is a clone designated herein as "DNA56433-1406".

20 Figure 112 shows the amino acid sequence (SEQ ID NO:112) derived from the coding sequence of SEQ ID NO:111 shown in Figure 111.

Figure 113 shows a nucleotide sequence (SEQ ID NO:113) of a native sequence PRO1012 cDNA, wherein SEQ ID NO:113 is a clone designated herein as "DNA56439-1376".

Figure 114 shows the amino acid sequence (SEQ ID NO:114) derived from the coding sequence of SEQ ID NO:113 shown in Figure 113.

25 Figure 115 shows a nucleotide sequence (SEQ ID NO:115) of a native sequence PRO1477 cDNA, wherein SEQ ID NO:115 is a clone designated herein as "DNA56529-1647".

Figure 116 shows the amino acid sequence (SEQ ID NO:116) derived from the coding sequence of SEQ ID NO:115 shown in Figure 115.

30 Figure 117 shows a nucleotide sequence (SEQ ID NO:117) of a native sequence PRO1134 cDNA, wherein SEQ ID NO:117 is a clone designated herein as "DNA56865-1491".

Figure 118 shows the amino acid sequence (SEQ ID NO:118) derived from the coding sequence of SEQ ID NO:117 shown in Figure 117.

Figure 119 shows a nucleotide sequence (SEQ ID NO:119) of a native sequence PRO162 cDNA, wherein SEQ ID NO:119 is a clone designated herein as "DNA56965-1356".

35 Figure 120 shows the amino acid sequence (SEQ ID NO:120) derived from the coding sequence of SEQ ID NO:119 shown in Figure 119.

Figure 121 shows a nucleotide sequence (SEQ ID NO:121) of a native sequence PRO1114 cDNA, wherein

SEQ ID NO:121 is a clone designated herein as "DNA57033-1403-1".

Figure 122 shows the amino acid sequence (SEQ ID NO:122) derived from the coding sequence of SEQ ID NO:121 shown in Figure 121.

5 Figure 123 shows a nucleotide sequence (SEQ ID NO:123) of a native sequence PRO828 cDNA, wherein SEQ ID NO:123 is a clone designated herein as "DNA57037-1444".

Figure 124 shows the amino acid sequence (SEQ ID NO:124) derived from the coding sequence of SEQ ID NO:123 shown in Figure 123.

Figure 125 shows a nucleotide sequence (SEQ ID NO:125) of a native sequence PRO827 cDNA, wherein SEQ ID NO:125 is a clone designated herein as "DNA57039-1402".

10 Figure 126 shows the amino acid sequence (SEQ ID NO:126) derived from the coding sequence of SEQ ID NO:125 shown in Figure 125.

Figure 127 shows a nucleotide sequence (SEQ ID NO:127) of a native sequence PRO1075 cDNA, wherein SEQ ID NO:127 is a clone designated herein as "DNA57689-1385".

15 Figure 128 shows the amino acid sequence (SEQ ID NO:128) derived from the coding sequence of SEQ ID NO:127 shown in Figure 127.

Figure 129 shows a nucleotide sequence (SEQ ID NO:129) of a native sequence PRO1007 cDNA, wherein SEQ ID NO:129 is a clone designated herein as "DNA57690-1374".

Figure 130 shows the amino acid sequence (SEQ ID NO:130) derived from the coding sequence of SEQ ID NO:129 shown in Figure 129.

20 Figure 131 shows a nucleotide sequence (SEQ ID NO:131) of a native sequence PRO826 cDNA, wherein SEQ ID NO:131 is a clone designated herein as "DNA57694-1341".

Figure 132 shows the amino acid sequence (SEQ ID NO:132) derived from the coding sequence of SEQ ID NO:131 shown in Figure 131.

25 Figure 133 shows a nucleotide sequence (SEQ ID NO:133) of a native sequence PRO819 cDNA, wherein SEQ ID NO:132 is a clone designated herein as "DNA57695-1340".

Figure 134 shows the amino acid sequence (SEQ ID NO:134) derived from the coding sequence of SEQ ID NO:133 shown in Figure 133.

Figure 135 shows a nucleotide sequence (SEQ ID NO:135) of a native sequence PRO1006 cDNA, wherein SEQ ID NO:135 is a clone designated herein as "DNA57699-1412".

30 Figure 136 shows the amino acid sequence (SEQ ID NO:136) derived from the coding sequence of SEQ ID NO:135 shown in Figure 135.

Figure 137 shows a nucleotide sequence (SEQ ID NO:137) of a native sequence PRO982 cDNA, wherein SEQ ID NO:137 is a clone designated herein as "DNA57700-1408".

35 Figure 138 shows the amino acid sequence (SEQ ID NO:138) derived from the coding sequence of SEQ ID NO:137 shown in Figure 137.

Figure 139 shows a nucleotide sequence (SEQ ID NO:139) of a native sequence PRO1005 cDNA, wherein SEQ ID NO:139 is a clone designated herein as "DNA57708-1411".

Figure 140 shows the amino acid sequence (SEQ ID NO:140) derived from the coding sequence of SEQ ID NO:139 shown in Figure 139.

Figure 141 shows a nucleotide sequence (SEQ ID NO:141) of a native sequence PRO791 cDNA, wherein SEQ ID NO:141 is a clone designated herein as "DNA57838-1337".

5 Figure 142 shows the amino acid sequence (SEQ ID NO:142) derived from the coding sequence of SEQ ID NO:141 shown in Figure 141.

Figure 143 shows a nucleotide sequence (SEQ ID NO:143) of a native sequence PRO1071 cDNA, wherein SEQ ID NO:143 is a clone designated herein as "DNA58847-1383".

10 Figure 144 shows the amino acid sequence (SEQ ID NO:144) derived from the coding sequence of SEQ ID NO:143 shown in Figure 43.

Figure 145 shows a nucleotide sequence (SEQ ID NO:145) of a native sequence PRO1415 cDNA, wherein SEQ ID NO:145 is a clone designated herein as "DNA58852-1637".

Figure 146 shows the amino acid sequence (SEQ ID NO:146) derived from the coding sequence of SEQ ID NO:145 shown in Figure 145.

15 Figure 147 shows a nucleotide sequence (SEQ ID NO:147) of a native sequence PRO1054 cDNA, wherein SEQ ID NO:147 is a clone designated herein as "DNA58853-1423".

Figure 148 shows the amino acid sequence (SEQ ID NO:148) derived from the coding sequence of SEQ ID NO:147 shown in Figure 147.

20 Figure 149 shows a nucleotide sequence (SEQ ID NO:149) of a native sequence PRO1411 cDNA, wherein SEQ ID NO:149 is a clone designated herein as "DNA59212-1627".

Figure 150 shows the amino acid sequence (SEQ ID NO:150) derived from the coding sequence of SEQ ID NO:149 shown in Figure 149.

Figure 151 shows a nucleotide sequence (SEQ ID NO:151) of a native sequence PRO1184 cDNA, wherein SEQ ID NO:151 is a clone designated herein as "DNA59220-1514".

25 Figure 152 shows the amino acid sequence (SEQ ID NO:152) derived from the coding sequence of SEQ ID NO:151 shown in Figure 151.

Figure 153 shows a nucleotide sequence (SEQ ID NO:153) of a native sequence PRO1029 cDNA, wherein SEQ ID NO:153 is a clone designated herein as "DNA59493-1420".

30 Figure 154 shows the amino acid sequence (SEQ ID NO:154) derived from the coding sequence of SEQ ID NO:153 shown in Figure 153.

Figure 155 shows a nucleotide sequence (SEQ ID NO:155) of a native sequence PRO1139 cDNA, wherein SEQ ID NO:155 is a clone designated herein as "DNA59497-1496".

Figure 156 shows the amino acid sequence (SEQ ID NO:156) derived from the coding sequence of SEQ ID NO:155 shown in Figure 155.

35 Figure 157 shows a nucleotide sequence (SEQ ID NO:157) of a native sequence PRO1190 cDNA, wherein SEQ ID NO:157 is a clone designated herein as "DNA59586-1520".

Figure 158 shows the amino acid sequence (SEQ ID NO:158) derived from the coding sequence of SEQ

ID NO:157 shown in Figure 157.

Figure 159 shows a nucleotide sequence (SEQ ID NO:159) of a native sequence PRO1309 cDNA, wherein SEQ ID NO:159 is a clone designated herein as "DNA59588-1571".

5 Figure 160 shows the amino acid sequence (SEQ ID NO:160) derived from the coding sequence of SEQ ID NO:159 shown in Figure 159.

Figure 161 shows a nucleotide sequence (SEQ ID NO:161) of a native sequence PRO836 cDNA, wherein SEQ ID NO:161 is a clone designated herein as "DNA59620-1463".

Figure 162 shows the amino acid sequence (SEQ ID NO:162) derived from the coding sequence of SEQ ID NO:161 shown in Figure 161.

10 Figure 163 shows a nucleotide sequence (SEQ ID NO:163) of a native sequence PRO1025 cDNA, wherein SEQ ID NO:163 is a clone designated herein as "DNA59622-1334".

Figure 164 shows the amino acid sequence (SEQ ID NO:164) derived from the coding sequence of SEQ ID NO:163 shown in Figure 163.

15 Figure 165 shows a nucleotide sequence (SEQ ID NO:165) of a native sequence PRO1131 cDNA, wherein SEQ ID NO:165 is a clone designated herein as "DNA59777-1480".

Figure 166 shows the amino acid sequence (SEQ ID NO:166) derived from the coding sequence of SEQ ID NO:165 shown in Figure 165.

Figure 167 shows a nucleotide sequence (SEQ ID NO:167) of a native sequence PRO1182 cDNA, wherein SEQ ID NO:167 is a clone designated herein as "DNA59848-1512".

20 Figure 168 shows the amino acid sequence (SEQ ID NO:168) derived from the coding sequence of SEQ ID NO:167 shown in Figure 167.

Figure 169 shows a nucleotide sequence (SEQ ID NO:169) of a native sequence PRO1155 cDNA, wherein SEQ ID NO:169 is a clone designated herein as "DNA59849-1504".

25 Figure 170 shows the amino acid sequence (SEQ ID NO:170) derived from the coding sequence of SEQ ID NO:169 shown in Figure 169.

Figure 171 shows a nucleotide sequence (SEQ ID NO:171) of a native sequence PRO1186 cDNA, wherein SEQ ID NO:171 is a clone designated herein as "DNA60621-1516".

Figure 172 shows the amino acid sequence (SEQ ID NO:172) derived from the coding sequence of SEQ ID NO:171 shown in Figure 171.

30 Figure 173 shows a nucleotide sequence (SEQ ID NO:173) of a native sequence PRO1198 cDNA, wherein SEQ ID NO:173 is a clone designated herein as "DNA60622-1525".

Figure 174 shows the amino acid sequence (SEQ ID NO:174) derived from the coding sequence of SEQ ID NO:173 shown in Figure 173.

35 Figure 175 shows a nucleotide sequence (SEQ ID NO:175) of a native sequence PRO1265 cDNA, wherein SEQ ID NO:175 is a clone designated herein as "DNA60764-1533".

Figure 176 shows the amino acid sequence (SEQ ID NO:176) derived from the coding sequence of SEQ ID NO:175 shown in Figure 175.

Figure 177 shows a nucleotide sequence (SEQ ID NO:177) of a native sequence PRO1361 cDNA, wherein SEQ ID NO:177 is a clone designated herein as "DNA60783-1611".

Figure 178 shows the amino acid sequence (SEQ ID NO:178) derived from the coding sequence of SEQ ID NO:177 shown in Figure 177.

5 Figure 179 shows a nucleotide sequence (SEQ ID NO:179) of a native sequence PRO1287 cDNA, wherein SEQ ID NO:179 is a clone designated herein as "DNA61755-1554".

Figure 180 shows the amino acid sequence (SEQ ID NO:180) derived from the coding sequence of SEQ ID NO:179 shown in Figure 179.

10 Figure 181 shows a nucleotide sequence (SEQ ID NO:181) of a native sequence PRO1308 cDNA, wherein SEQ ID NO:181 is a clone designated herein as "DNA62306-1570".

Figure 182 shows the amino acid sequence (SEQ ID NO:182) derived from the coding sequence of SEQ ID NO:181 shown in Figure 181.

Figure 183 shows a nucleotide sequence (SEQ ID NO:183) of a native sequence PRO4313 cDNA, wherein SEQ ID NO:183 is a clone designated herein as "DNA62312-2558".

15 Figure 184 shows the amino acid sequence (SEQ ID NO:184) derived from the coding sequence of SEQ ID NO:183 shown in Figure 183.

Figure 185 shows a nucleotide sequence (SEQ ID NO:185) of a native sequence PRO1192 cDNA, wherein SEQ ID NO:185 is a clone designated herein as "DNA62814-1521".

20 Figure 186 shows the amino acid sequence (SEQ ID NO:186) derived from the coding sequence of SEQ ID NO:185 shown in Figure 185.

Figure 187 shows a nucleotide sequence (SEQ ID NO:187) of a native sequence PRO1160 cDNA, wherein SEQ ID NO:187 is a clone designated herein as "DNA62872-1509".

Figure 188 shows the amino acid sequence (SEQ ID NO:188) derived from the coding sequence of SEQ ID NO:187 shown in Figure 187.

25 Figure 189 shows a nucleotide sequence (SEQ ID NO:189) of a native sequence PRO1244 cDNA, wherein SEQ ID NO:189 is a clone designated herein as "DNA64883-1526".

Figure 190 shows the amino acid sequence (SEQ ID NO:190) derived from the coding sequence of SEQ ID NO:189 shown in Figure 189.

30 Figure 191 shows a nucleotide sequence (SEQ ID NO:191) of a native sequence PRO1356 cDNA, wherein SEQ ID NO:191 is a clone designated herein as "DNA64886-1601".

Figure 192 shows the amino acid sequence (SEQ ID NO:192) derived from the coding sequence of SEQ ID NO:191 shown in Figure 191.

Figure 193 shows a nucleotide sequence (SEQ ID NO:193) of a native sequence PRO1274 cDNA, wherein SEQ ID NO:193 is a clone designated herein as "DNA64889-1541".

35 Figure 194 shows the amino acid sequence (SEQ ID NO:194) derived from the coding sequence of SEQ ID NO:193 shown in Figure 193.

Figure 195 shows a nucleotide sequence (SEQ ID NO:195) of a native sequence PRO1272 cDNA, wherein

SEQ ID NO:195 is a clone designated herein as "DNA64896-1539".

Figure 196 shows the amino acid sequence (SEQ ID NO:196) derived from the coding sequence of SEQ ID NO:195 shown in Figure 195.

5 Figure 197 shows a nucleotide sequence (SEQ ID NO:197) of a native sequence PRO1412 cDNA, wherein SEQ ID NO:197 is a clone designated herein as "DNA64897-1628".

Figure 198 shows the amino acid sequence (SEQ ID NO:198) derived from the coding sequence of SEQ ID NO:197 shown in Figure 197.

Figure 199 shows a nucleotide sequence (SEQ ID NO:199) of a native sequence PRO1286 cDNA, wherein SEQ ID NO:199 is a clone designated herein as "DNA64903-1553".

10 Figure 200 shows the amino acid sequence (SEQ ID NO:200) derived from the coding sequence of SEQ ID NO:199 shown in Figure 199.

Figure 201 shows a nucleotide sequence (SEQ ID NO:201) of a native sequence PRO1347 cDNA, wherein SEQ ID NO:201 is a clone designated herein as "DNA64950-1590".

15 Figure 202 shows the amino acid sequence (SEQ ID NO:202) derived from the coding sequence of SEQ ID NO:201 shown in Figure 201.

Figure 203 shows a nucleotide sequence (SEQ ID NO:203) of a native sequence PRO1273 cDNA, wherein SEQ ID NO:203 is a clone designated herein as "DNA65402-1540".

Figure 204 shows the amino acid sequence (SEQ ID NO:204) derived from the coding sequence of SEQ ID NO:203 shown in Figure 203.

20 Figure 205 shows a nucleotide sequence (SEQ ID NO:205) of a native sequence PRO1283 cDNA, wherein SEQ ID NO:205 is a clone designated herein as "DNA65404-1551".

Figure 206 shows the amino acid sequence (SEQ ID NO:206) derived from the coding sequence of SEQ ID NO:205 shown in Figure 205.

25 Figure 207 shows a nucleotide sequence (SEQ ID NO:207) of a native sequence PRO1279 cDNA, wherein SEQ ID NO:207 is a clone designated herein as "DNA65405-1547".

Figure 208 shows the amino acid sequence (SEQ ID NO:208) derived from the coding sequence of SEQ ID NO:207 shown in Figure 207.

Figure 209 shows a nucleotide sequence (SEQ ID NO:209) of a native sequence PRO1306 cDNA, wherein SEQ ID NO:209 is a clone designated herein as "DNA65410-1569".

30 Figure 210 shows the amino acid sequence (SEQ ID NO:210) derived from the coding sequence of SEQ ID NO:209 shown in Figure 209.

Figure 211 shows a nucleotide sequence (SEQ ID NO:211) of a native sequence PRO1195 cDNA, wherein SEQ ID NO:211 is a clone designated herein as "DNA65412-1523".

35 Figure 212 shows the amino acid sequence (SEQ ID NO:212) derived from the coding sequence of SEQ ID NO:211 shown in Figure 211.

Figure 213 shows a nucleotide sequence (SEQ ID NO:213) of a native sequence PRO4995 cDNA, wherein SEQ ID NO:213 is a clone designated herein as "DNA66307-2661".

Figure 214 shows the amino acid sequence (SEQ ID NO:214) derived from the coding sequence of SEQ ID NO:213 shown in Figure 213.

Figure 215 shows a nucleotide sequence (SEQ ID NO:215) of a native sequence PRO1382 cDNA, wherein SEQ ID NO:215 is a clone designated herein as "DNA66526-1616".

5 Figure 216 shows the amino acid sequence (SEQ ID NO:216) derived from the coding sequence of SEQ ID NO:215 shown in Figure 215.

Figure 217 shows a nucleotide sequence (SEQ ID NO:217) of a native sequence PRO1325 cDNA, wherein SEQ ID NO:217 is a clone designated herein as "DNA66659-1593".

10 Figure 218 shows the amino acid sequence (SEQ ID NO:218) derived from the coding sequence of SEQ ID NO:217 shown in Figure 217.

Figure 219 shows a nucleotide sequence (SEQ ID NO:219) of a native sequence PRO1329 cDNA, wherein SEQ ID NO:219 is a clone designated herein as "DNA66660-1585".

Figure 220 shows the amino acid sequence (SEQ ID NO:220) derived from the coding sequence of SEQ ID NO:219 shown in Figure 219.

15 Figure 221 shows a nucleotide sequence (SEQ ID NO:221) of a native sequence PRO1338 cDNA, wherein SEQ ID NO:221 is a clone designated herein as "DNA66667-1596".

Figure 222 shows the amino acid sequence (SEQ ID NO:222) derived from the coding sequence of SEQ ID NO:221 shown in Figure 221.

20 Figure 223 shows a nucleotide sequence (SEQ ID NO:223) of a native sequence PRO1337 cDNA, wherein SEQ ID NO:223 is a clone designated herein as "DNA66672-1586".

Figure 224 shows the amino acid sequence (SEQ ID NO:224) derived from the coding sequence of SEQ ID NO:223 shown in Figure 223.

Figure 225 shows a nucleotide sequence (SEQ ID NO:225) of a native sequence PRO1343 cDNA, wherein SEQ ID NO:225 is a clone designated herein as "DNA66675-1587".

25 Figure 226 shows the amino acid sequence (SEQ ID NO:226) derived from the coding sequence of SEQ ID NO:225 shown in Figure 225.

Figure 227 shows a nucleotide sequence (SEQ ID NO:227) of a native sequence PRO1376 cDNA, wherein SEQ ID NO:227 is a clone designated herein as "DNA67300-1605".

30 Figure 228 shows the amino acid sequence (SEQ ID NO:228) derived from the coding sequence of SEQ ID NO:227 shown in Figure 227.

Figure 229 shows a nucleotide sequence (SEQ ID NO:229) of a native sequence PRO1434 cDNA, wherein SEQ ID NO:229 is a clone designated herein as "DNA68818-2536".

Figure 230 shows the amino acid sequence (SEQ ID NO:230) derived from the coding sequence of SEQ ID NO:229 shown in Figure 229.

35 Figure 231 shows a nucleotide sequence (SEQ ID NO:231) of a native sequence PRO3579 cDNA, wherein SEQ ID NO:231 is a clone designated herein as "DNA68862-2546".

Figure 232 shows the amino acid sequence (SEQ ID NO:232) derived from the coding sequence of SEQ

ID NO:231 shown in Figure 231.

Figure 233 shows a nucleotide sequence (SEQ ID NO:233) of a native sequence PRO1387 cDNA, wherein SEQ ID NO:233 is a clone designated herein as "DNA68872-1620".

5 Figure 234 shows the amino acid sequence (SEQ ID NO:234) derived from the coding sequence of SEQ ID NO:233 shown in Figure 233.

Figure 235 shows a nucleotide sequence (SEQ ID NO:235) of a native sequence PRO1419 cDNA, wherein SEQ ID NO:235 is a clone designated herein as "DNA71290-1630".

Figure 236 shows the amino acid sequence (SEQ ID NO:236) derived from the coding sequence of SEQ ID NO:235 shown in Figure 235.

10 Figure 237 shows a nucleotide sequence (SEQ ID NO:237) of a native sequence PRO1488 cDNA, wherein SEQ ID NO:237 is a clone designated herein as "DNA73736-1657".

Figure 238 shows the amino acid sequence (SEQ ID NO:238) derived from the coding sequence of SEQ ID NO:237 shown in Figure 237.

15 Figure 239 shows a nucleotide sequence (SEQ ID NO:239) of a native sequence PRO1474 cDNA, wherein SEQ ID NO:239 is a clone designated herein as "DNA73739-1645".

Figure 240 shows the amino acid sequence (SEQ ID NO:240) derived from the coding sequence of SEQ ID NO:239 shown in Figure 239.

Figure 241 shows a nucleotide sequence (SEQ ID NO:241) of a native sequence PRO1508 cDNA, wherein SEQ ID NO:241 is a clone designated herein as "DNA73742-1662".

20 Figure 242 shows the amino acid sequence (SEQ ID NO:242) derived from the coding sequence of SEQ ID NO:241 shown in Figure 241.

Figure 243 shows a nucleotide sequence (SEQ ID NO:243) of a native sequence PRO1754 cDNA, wherein SEQ ID NO:243 is a clone designated herein as "DNA76385-1692".

25 Figure 244 shows the amino acid sequence (SEQ ID NO:244) derived from the coding sequence of SEQ ID NO:243 shown in Figure 243.

Figure 245 shows a nucleotide sequence (SEQ ID NO:245) of a native sequence PRO1550 cDNA, wherein SEQ ID NO:245 is a clone designated herein as "DNA76393-1664".

Figure 246 shows the amino acid sequence (SEQ ID NO:246) derived from the coding sequence of SEQ ID NO:245 shown in Figure 245.

30 Figure 247 shows a nucleotide sequence (SEQ ID NO:247) of a native sequence PRO1758 cDNA, wherein SEQ ID NO:247 is a clone designated herein as "DNA76399-1700".

Figure 248 shows the amino acid sequence (SEQ ID NO:248) derived from the coding sequence of SEQ ID NO:247 shown in Figure 247.

35 Figure 249 shows a nucleotide sequence (SEQ ID NO:249) of a native sequence PRO1917 cDNA, wherein SEQ ID NO:249 is a clone designated herein as "DNA76400-2528".

Figure 250 shows the amino acid sequence (SEQ ID NO:250) derived from the coding sequence of SEQ ID NO:249 shown in Figure 249.

Figure 251 shows a nucleotide sequence (SEQ ID NO:251) of a native sequence PRO1787 cDNA, wherein SEQ ID NO:251 is a clone designated herein as "DNA76510-2504".

Figure 252 shows the amino acid sequence (SEQ ID NO:252) derived from the coding sequence of SEQ ID NO:251 shown in Figure 251.

5 Figure 253 shows a nucleotide sequence (SEQ ID NO:253) of a native sequence PRO1556 cDNA, wherein SEQ ID NO:253 is a clone designated herein as "DNA76529-1666".

Figure 254 shows the amino acid sequence (SEQ ID NO:254) derived from the coding sequence of SEQ ID NO:253 shown in Figure 253.

10 Figure 255 shows a nucleotide sequence (SEQ ID NO:255) of a native sequence PRO1760 cDNA, wherein SEQ ID NO:255 is a clone designated herein as "DNA76532-1702".

Figure 256 shows the amino acid sequence (SEQ ID NO:256) derived from the coding sequence of SEQ ID NO:255 shown in Figure 255.

Figure 257 shows a nucleotide sequence (SEQ ID NO:257) of a native sequence PRO1567 cDNA, wherein SEQ ID NO:257 is a clone designated herein as "DNA76541-1675".

15 Figure 258 shows the amino acid sequence (SEQ ID NO:258) derived from the coding sequence of SEQ ID NO:257 shown in Figure 257.

Figure 259 shows a nucleotide sequence (SEQ ID NO:259) of a native sequence PRO1600 cDNA, wherein SEQ ID NO:259 is a clone designated herein as "DNA77503-1686".

20 Figure 260 shows the amino acid sequence (SEQ ID NO:260) derived from the coding sequence of SEQ ID NO:259 shown in Figure 259.

Figure 261 shows a nucleotide sequence (SEQ ID NO:261) of a native sequence PRO1868 cDNA, wherein SEQ ID NO:261 is a clone designated herein as "DNA77624-2515".

Figure 262 shows the amino acid sequence (SEQ ID NO:262) derived from the coding sequence of SEQ ID NO:261 shown in Figure 261.

25 Figure 263 shows a nucleotide sequence (SEQ ID NO:263) of a native sequence PRO1890 cDNA, wherein SEQ ID NO:263 is a clone designated herein as "DNA79230-2525".

Figure 264 shows the amino acid sequence (SEQ ID NO:264) derived from the coding sequence of SEQ ID NO:263 shown in Figure 263.

30 Figure 265 shows a nucleotide sequence (SEQ ID NO:265) of a native sequence PRO1887 cDNA, wherein SEQ ID NO:265 is a clone designated herein as "DNA79862-2522".

Figure 266 shows the amino acid sequence (SEQ ID NO:265) derived from the coding sequence of SEQ ID NO:265 shown in Figure 265.

Figure 267 shows a nucleotide sequence (SEQ ID NO:267) of a native sequence PRO4353 cDNA, wherein SEQ ID NO:267 is a clone designated herein as "DNA80145-2594".

35 Figure 268 shows the amino acid sequence (SEQ ID NO:268) derived from the coding sequence of SEQ ID NO:267 shown in Figure 267.

Figure 269 shows a nucleotide sequence (SEQ ID NO:269) of a native sequence PRO1782 cDNA, wherein

SEQ ID NO:269 is a clone designated herein as "DNA80899-2501".

Figure 270 shows the amino acid sequence (SEQ ID NO:270) derived from the coding sequence of SEQ ID NO:269 shown in Figure 269.

5 Figure 271 shows a nucleotide sequence (SEQ ID NO:271) of a native sequence PRO1928 cDNA, wherein SEQ ID NO:271 is a clone designated herein as "DNA81754-2532".

Figure 272 shows the amino acid sequence (SEQ ID NO:272) derived from the coding sequence of SEQ ID NO:271 shown in Figure 271.

Figure 273 shows a nucleotide sequence (SEQ ID NO:273) of a native sequence PRO1865 cDNA, wherein SEQ ID NO:273 is a clone designated herein as "DNA81757-2512".

10 Figure 274 shows the amino acid sequence (SEQ ID NO:274) derived from the coding sequence of SEQ ID NO:273 shown in Figure 273.

Figure 275 shows a nucleotide sequence (SEQ ID NO:275) of a native sequence PRO4341 cDNA, wherein SEQ ID NO:275 is a clone designated herein as "DNA81761-2583".

15 Figure 276 shows the amino acid sequence (SEQ ID NO:276) derived from the coding sequence of SEQ ID NO:275 shown in Figure 275.

Figure 277 shows a nucleotide sequence (SEQ ID NO:277) of a native sequence PRO6714 cDNA, wherein SEQ ID NO:277 is a clone designated herein as "DNA82358-2738".

Figure 278 shows the amino acid sequence (SEQ ID NO:278) derived from the coding sequence of SEQ ID NO:277 shown in Figure 277.

20 Figure 279 shows a nucleotide sequence (SEQ ID NO:279) of a native sequence PRO5723 cDNA, wherein SEQ ID NO:279 is a clone designated herein as "DNA82361".

Figure 280 shows the amino acid sequence (SEQ ID NO:280) derived from the coding sequence of SEQ ID NO:279 shown in Figure 279.

25 Figure 281 shows a nucleotide sequence (SEQ ID NO:281) of a native sequence PRO3438 cDNA, wherein SEQ ID NO:281 is a clone designated herein as "DNA82364-2538".

Figure 282 shows the amino acid sequence (SEQ ID NO:282) derived from the coding sequence of SEQ ID NO:281 shown in Figure 281.

Figure 283 shows a nucleotide sequence (SEQ ID NO:283) of a native sequence PRO6071 cDNA, wherein SEQ ID NO:283 is a clone designated herein as "DNA82403-2959".

30 Figure 284 shows the amino acid sequence (SEQ ID NO:284) derived from the coding sequence of SEQ ID NO:283 shown in Figure 283.

Figure 285 shows a nucleotide sequence (SEQ ID NO:285) of a native sequence PRO1801 cDNA, wherein SEQ ID NO:285 is a clone designated herein as "DNA83500-2506".

35 Figure 286 shows the amino acid sequence (SEQ ID NO:286) derived from the coding sequence of SEQ ID NO:285 shown in Figure 285.

Figure 287 shows a nucleotide sequence (SEQ ID NO:287) of a native sequence PRO4324 cDNA, wherein SEQ ID NO:287 is a clone designated herein as "DNA83560-2569".

Figure 288 shows the amino acid sequence (SEQ ID NO:288) derived from the coding sequence of SEQ ID NO:287 shown in Figure 287.

Figure 289 shows a nucleotide sequence (SEQ ID NO:289) of a native sequence PRO4333 cDNA, wherein SEQ ID NO:289 is a clone designated herein as "DNA84210-2576".

5 Figure 290 shows the amino acid sequence (SEQ ID NO:290) derived from the coding sequence of SEQ ID NO:289 shown in Figure 289.

Figure 291 shows a nucleotide sequence (SEQ ID NO:291) of a native sequence PRO4405 cDNA, wherein SEQ ID NO:291 is a clone designated herein as "DNA84920-2614".

10 Figure 292 shows the amino acid sequence (SEQ ID NO:292) derived from the coding sequence of SEQ ID NO:291 shown in Figure 291.

Figure 293 shows a nucleotide sequence (SEQ ID NO:293) of a native sequence PRO4356 cDNA, wherein SEQ ID NO:293 is a clone designated herein as "DNA86576-2595".

Figure 294 shows the amino acid sequence (SEQ ID NO:294) derived from the coding sequence of SEQ ID NO:293 shown in Figure 293.

15 Figure 295 shows a nucleotide sequence (SEQ ID NO:295) of a native sequence PRO3444 cDNA, wherein SEQ ID NO:295 is a clone designated herein as "DNA87997".

Figure 296 shows the amino acid sequence (SEQ ID NO:296) derived from the coding sequence of SEQ ID NO:295 shown in Figure 295.

20 Figure 297 shows a nucleotide sequence (SEQ ID NO:297) of a native sequence PRO4302 cDNA, wherein SEQ ID NO:297 is a clone designated herein as "DNA92218-2554".

Figure 298 shows the amino acid sequence (SEQ ID NO:298) derived from the coding sequence of SEQ ID NO:297 shown in Figure 297.

Figure 299 shows a nucleotide sequence (SEQ ID NO:299) of a native sequence PRO4371 cDNA, wherein SEQ ID NO:299 is a clone designated herein as "DNA92233-2599".

25 Figure 300 shows the amino acid sequence (SEQ ID NO:300) derived from the coding sequence of SEQ ID NO:299 shown in Figure 299.

Figure 301 shows a nucleotide sequence (SEQ ID NO:301) of a native sequence PRO4354 cDNA, wherein SEQ ID NO:301 is a clone designated herein as "DNA92256-2596".

30 Figure 302 shows the amino acid sequence (SEQ ID NO:302) derived from the coding sequence of SEQ ID NO:301 shown in Figure 301.

Figure 303 shows a nucleotide sequence (SEQ ID NO:303) of a native sequence PRO5725 cDNA, wherein SEQ ID NO:303 is a clone designated herein as "DNA92265-2669".

Figure 304 shows the amino acid sequence (SEQ ID NO:304) derived from the coding sequence of SEQ ID NO:303 shown in Figure 303.

35 Figure 305 shows a nucleotide sequence (SEQ ID NO:305) of a native sequence PRO4408 cDNA, wherein SEQ ID NO:305 is a clone designated herein as "DNA92274-2617".

Figure 306 shows the amino acid sequence (SEQ ID NO:306) derived from the coding sequence of SEQ

ID NO:305 shown in Figure 305.

Figure 307 shows a nucleotide sequence (SEQ ID NO:307) of a native sequence PRO9940 cDNA, wherein SEQ ID NO:307 is a clone designated herein as "DNA92282".

5 Figure 308 shows the amino acid sequence (SEQ ID NO:308) derived from the coding sequence of SEQ ID NO:307 shown in Figure 307.

Figure 309 shows a nucleotide sequence (SEQ ID NO:309) of a native sequence PRO5737 cDNA, wherein SEQ ID NO:309 is a clone designated herein as "DNA92929-2534-1".

Figure 310 shows the amino acid sequence (SEQ ID NO:310) derived from the coding sequence of SEQ ID NO:309 shown in Figure 309.

10 Figure 311 shows a nucleotide sequence (SEQ ID NO:311) of a native sequence PRO4425 cDNA, wherein SEQ ID NO:311 is a clone designated herein as "DNA93011-2637".

Figure 312 shows the amino acid sequence (SEQ ID NO:312) derived from the coding sequence of SEQ ID NO:311 shown in Figure 311.

15 Figure 313 shows a nucleotide sequence (SEQ ID NO:313) of a native sequence PRO4345 cDNA, wherein SEQ ID NO:313 is a clone designated herein as "DNA94854-2586".

Figure 314 shows the amino acid sequence (SEQ ID NO:314) derived from the coding sequence of SEQ ID NO:313 shown in Figure 313.

Figure 315 shows a nucleotide sequence (SEQ ID NO:315) of a native sequence PRO4342 cDNA, wherein SEQ ID NO:315 is a clone designated herein as "DNA96787-2534-1".

20 Figure 316 shows the amino acid sequence (SEQ ID NO:316) derived from the coding sequence of SEQ ID NO:315 shown in Figure 315.

Figure 317 shows a nucleotide sequence (SEQ ID NO:317) of a native sequence PRO3562 cDNA, wherein SEQ ID NO:317 is a clone designated herein as "DNA96791".

25 Figure 318 shows the amino acid sequence (SEQ ID NO:318) derived from the coding sequence of SEQ ID NO:317 shown in Figure 317.

Figure 319 shows a nucleotide sequence (SEQ ID NO:319) of a native sequence PRO4422 cDNA, wherein SEQ ID NO:319 is a clone designated herein as "DNA96867-2620".

Figure 320 shows the amino acid sequence (SEQ ID NO:320) derived from the coding sequence of SEQ ID NO:319 shown in Figure 319.

30 Figure 321 shows a nucleotide sequence (SEQ ID NO:321) of a native sequence PRO5776 cDNA, wherein SEQ ID NO:321 is a clone designated herein as "DNA96872-2674".

Figure 322 shows the amino acid sequence (SEQ ID NO:322) derived from the coding sequence of SEQ ID NO:321 shown in Figure 321.

35 Figure 323 shows a nucleotide sequence (SEQ ID NO:323) of a native sequence PRO4430 cDNA, wherein SEQ ID NO:323 is a clone designated herein as "DNA96878-2626".

Figure 324 shows the amino acid sequence (SEQ ID NO:324) derived from the coding sequence of SEQ ID NO:323 shown in Figure 323.

Figure 325 shows a nucleotide sequence (SEQ ID NO:325) of a native sequence PRO4499 cDNA, wherein SEQ ID NO:325 is a clone designated herein as "DNA96889-2641".

Figure 326 shows the amino acid sequence (SEQ ID NO:326) derived from the coding sequence of SEQ ID NO:325 shown in Figure 325.

5 Figure 327 shows a nucleotide sequence (SEQ ID NO:327) of a native sequence PRO4503 cDNA, wherein SEQ ID NO:327 is a clone designated herein as "DNA100312-2645".

Figure 328 shows the amino acid sequence (SEQ ID NO:328) derived from the coding sequence of SEQ ID NO:327 shown in Figure 327.

10 Figure 329 shows a nucleotide sequence (SEQ ID NO:329) of a native sequence PRO10008 cDNA, wherein SEQ ID NO:329 is a clone designated herein as "DNA101921".

Figure 330 shows the amino acid sequence (SEQ ID NO:330) derived from the coding sequence of SEQ ID NO:329 shown in Figure 329.

Figure 331 shows a nucleotide sequence (SEQ ID NO:331) of a native sequence PRO5730 cDNA, wherein SEQ ID NO:331 is a clone designated herein as "DNA101926".

15 Figure 332 shows the amino acid sequence (SEQ ID NO:332) derived from the coding sequence of SEQ ID NO:331 shown in Figure 331.

Figure 333 shows a nucleotide sequence (SEQ ID NO:333) of a native sequence PRO6008 cDNA, wherein SEQ ID NO:333 is a clone designated herein as "DNA102844".

20 Figure 334 shows the amino acid sequence (SEQ ID NO:334) derived from the coding sequence of SEQ ID NO:333 shown in Figure 333.

Figure 335 shows a nucleotide sequence (SEQ ID NO:335) of a native sequence PRO4527 cDNA, wherein SEQ ID NO:335 is a clone designated herein as "DNA103197".

Figure 336 shows the amino acid sequence (SEQ ID NO:336) derived from the coding sequence of SEQ ID NO:335 shown in Figure 335.

25 Figure 337 shows a nucleotide sequence (SEQ ID NO:337) of a native sequence PRO4538 cDNA, wherein SEQ ID NO:337 is a clone designated herein as "DNA103208".

Figure 338 shows the amino acid sequence (SEQ ID NO:338) derived from the coding sequence of SEQ ID NO:337 shown in Figure 337.

30 Figure 339 shows a nucleotide sequence (SEQ ID NO:339) of a native sequence PRO4553 cDNA, wherein SEQ ID NO:339 is a clone designated herein as "DNA103223".

Figure 340 shows the amino acid sequence (SEQ ID NO:340) derived from the coding sequence of SEQ ID NO:339 shown in Figure 339.

Figure 341 shows a nucleotide sequence (SEQ ID NO:341) of a native sequence PRO6006 cDNA, wherein SEQ ID NO:341 is a clone designated herein as "DNA105782-2693".

35 Figure 342 shows the amino acid sequence (SEQ ID NO:342) derived from the coding sequence of SEQ ID NO:341 shown in Figure 341.

Figure 343 shows a nucleotide sequence (SEQ ID NO:343) of a native sequence PRO6029 cDNA, wherein

SEQ ID NO:343 is a clone designated herein as "DNA105849-2704".

Figure 344 shows the amino acid sequence (SEQ ID NO:344) derived from the coding sequence of SEQ ID NO:343 shown in Figure 343.

5 Figure 345 shows a nucleotide sequence (SEQ ID NO:345) of a native sequence PRO9821 cDNA, wherein SEQ ID NO:345 is a clone designated herein as "DNA108725-2766".

Figure 346 shows the amino acid sequence (SEQ ID NO:346) derived from the coding sequence of SEQ ID NO:345 shown in Figure 345.

Figure 347 shows a nucleotide sequence (SEQ ID NO:347) of a native sequence PRO9820 cDNA, wherein SEQ ID NO:347 is a clone designated herein as "DNA108769-2765".

10 Figure 348 shows the amino acid sequence (SEQ ID NO:348) derived from the coding sequence of SEQ ID NO:347 shown in Figure 347.

Figure 349 shows a nucleotide sequence (SEQ ID NO:349) of a native sequence PRO9771 cDNA, wherein SEQ ID NO:349 is a clone designated herein as "DNA119498-2965".

15 Figure 350 shows the amino acid sequence (SEQ ID NO:350) derived from the coding sequence of SEQ ID NO:349 shown in Figure 349.

Figure 351 shows a nucleotide sequence (SEQ ID NO:351) of a native sequence PRO7436 cDNA, wherein SEQ ID NO:351 is a clone designated herein as "DNA119535-2756".

Figure 352 shows the amino acid sequence (SEQ ID NO:352) derived from the coding sequence of SEQ ID NO:351 shown in Figure 351.

20 Figure 353 shows a nucleotide sequence (SEQ ID NO:353) of a native sequence PRO10096 cDNA, wherein SEQ ID NO:353 is a clone designated herein as "DNA125185-2806".

Figure 354 shows the amino acid sequence (SEQ ID NO:354) derived from the coding sequence of SEQ ID NO:353 shown in Figure 353.

25 Figure 355 shows a nucleotide sequence (SEQ ID NO:355) of a native sequence PRO19670 cDNA, wherein SEQ ID NO:355 is a clone designated herein as "DNA131639-2874".

Figure 356 shows the amino acid sequence (SEQ ID NO:356) derived from the coding sequence of SEQ ID NO:355 shown in Figure 355.

Figure 357 shows a nucleotide sequence (SEQ ID NO:357) of a native sequence PRO20044 cDNA, wherein SEQ ID NO:357 is a clone designated herein as "DNA139623-2893".

30 Figure 358 shows the amino acid sequence (SEQ ID NO:358) derived from the coding sequence of SEQ ID NO:357 shown in Figure 357.

Figure 359 shows a nucleotide sequence (SEQ ID NO:359) of a native sequence PRO9873 cDNA, wherein SEQ ID NO:359 is a clone designated herein as "DNA143076-2787".

35 Figure 360 shows the amino acid sequence (SEQ ID NO:360) derived from the coding sequence of SEQ ID NO:359 shown in Figure 359.

Figure 361 shows a nucleotide sequence (SEQ ID NO:361) of a native sequence PRO21366 cDNA, wherein SEQ ID NO:361 is a clone designated herein as "DNA143276-2975".

Figure 362 shows the amino acid sequence (SEQ ID NO:362) derived from the coding sequence of SEQ ID NO:361 shown in Figure 361.

Figure 363 shows a nucleotide sequence (SEQ ID NO:363) of a native sequence PRO20040 cDNA, wherein SEQ ID NO:363 is a clone designated herein as "DNA164625-2890".

5 Figure 364 shows the amino acid sequence (SEQ ID NO:364) derived from the coding sequence of SEQ ID NO:363 shown in Figure 363.

Figure 365 shows a nucleotide sequence (SEQ ID NO:365) of a native sequence PRO21184 cDNA, wherein SEQ ID NO:365 is a clone designated herein as "DNA167678-2963".

10 Figure 366 shows the amino acid sequence (SEQ ID NO:366) derived from the coding sequence of SEQ ID NO:365 shown in Figure 365.

Figure 367 shows a nucleotide sequence (SEQ ID NO:367) of a native sequence PRO21055 cDNA, wherein SEQ ID NO:367 is a clone designated herein as "DNA170021-2923".

Figure 368 shows the amino acid sequence (SEQ ID NO:368) derived from the coding sequence of SEQ ID NO:367 shown in Figure 367.

15 Figure 369 shows a nucleotide sequence (SEQ ID NO:369) of a native sequence PRO28631 cDNA, wherein SEQ ID NO:369 is a clone designated herein as "DNA170212-3000".

Figure 370 shows the amino acid sequence (SEQ ID NO:370) derived from the coding sequence of SEQ ID NO:369 shown in Figure 369.

20 Figure 371 shows a nucleotide sequence (SEQ ID NO:371) of a native sequence PRO21384 cDNA, wherein SEQ ID NO:371 is a clone designated herein as "DNA177313-2982".

Figure 372 shows the amino acid sequence (SEQ ID NO:372) derived from the coding sequence of SEQ ID NO:371 shown in Figure 371.

Figure 373 shows a nucleotide sequence (SEQ ID NO:373) of a native sequence PRO1449 cDNA, wherein SEQ ID NO:373 is a clone designated herein as "DNA64908-1163-1".

25 Figure 374 shows the amino acid sequence (SEQ ID NO:374) derived from the coding sequence of SEQ ID NO:373 shown in Figure 373.

Figure 375 shows wholemount in situ hybridization results on mouse embryos using a mouse orthologue of PRO1449 which has about 78% amino acid identity with PRO1449. The results show that PRO1449 orthologue is expressed in the developing vasculature. The cross-section further shows expression in endothelial cells and progenitors of endothelial cells.

30

Figure 376 shows that a PRO1449 orthologue having about 78% amino acid identity with PRO1449 is expressed in vasculature of many inflamed and diseased tissues, but is very low, or lacking, in normal adult vessels.

Figure 377 shows that a PRO1449 orthologue having about 78% amino acid identity with PRO1449 induces ectopic vessels in the eyes of chicken embryos.

35

5. Detailed Description of the Invention

5.1. Definitions

The phrases "cardiovascular, endothelial and angiogenic disorder", "cardiovascular, endothelial and angiogenic dysfunction", "cardiovascular, endothelial or angiogenic disorder" and "cardiovascular, endothelial or angiogenic dysfunction" are used interchangeably and refer in part to systemic disorders that affect vessels, such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins, and/or lymphatics. This would include indications that stimulate angiogenesis and/or cardiovascularization, and those that inhibit angiogenesis and/or cardiovascularization. Such disorders include, for example, arterial disease, such as atherosclerosis, hypertension, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, and arterial restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, cancer such as vascular tumors, e.g., hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, Kaposi's sarcoma, lymphangioma, and lymphangiosarcoma, tumor angiogenesis, trauma such as wounds, burns, and other injured tissue, implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. This would also include angina, myocardial infarctions such as acute myocardial infarctions, cardiac hypertrophy, and heart failure such as CHF.

"Hypertrophy", as used herein, is defined as an increase in mass of an organ or structure independent of natural growth that does not involve tumor formation. Hypertrophy of an organ or tissue is due either to an increase in the mass of the individual cells (true hypertrophy), or to an increase in the number of cells making up the tissue (hyperplasia), or both. Certain organs, such as the heart, lose the ability to divide shortly after birth. Accordingly, "cardiac hypertrophy" is defined as an increase in mass of the heart, which, in adults, is characterized by an increase in myocyte cell size and contractile protein content without concomitant cell division. The character of the stress responsible for inciting the hypertrophy, (e.g., increased preload, increased afterload, loss of myocytes, as in myocardial infarction, or primary depression of contractility), appears to play a critical role in determining the nature of the response. The early stage of cardiac hypertrophy is usually characterized morphologically by increases in the size of myofibrils and mitochondria, as well as by enlargement of mitochondria and nuclei. At this stage, while muscle cells are larger than normal, cellular organization is largely preserved. At a more advanced stage of cardiac hypertrophy, there are preferential increases in the size or number of specific organelles, such as mitochondria, and new contractile elements are added in localized areas of the cells, in an irregular manner. Cells subjected to long-standing hypertrophy show more obvious disruptions in cellular organization, including markedly enlarged nuclei with highly lobulated membranes, which displace adjacent myofibrils and cause breakdown of normal Z-band registration. The phrase "cardiac hypertrophy" is used to include all stages of the progression of this condition, characterized by various degrees of structural damage of the heart muscle, regardless of the underlying cardiac disorder. Hence, the term also includes physiological conditions instrumental in the development of cardiac hypertrophy, such as elevated blood pressure, aortic stenosis, or myocardial infarction.

"Heart failure" refers to an abnormality of cardiac function where the heart does not pump blood at the rate

needed for the requirements of metabolizing tissues. The heart failure can be caused by a number of factors, including ischemic, congenital, rheumatic, or idiopathic forms.

“Congestive heart failure” (CHF) is a progressive pathologic state where the heart is increasingly unable to supply adequate cardiac output (the volume of blood pumped by the heart over time) to deliver the oxygenated blood to peripheral tissues. As CHF progresses, structural and hemodynamic damages occur. While these damages have a variety of manifestations, one characteristic symptom is ventricular hypertrophy. CHF is a common end result of a number of various cardiac disorders.

“Myocardial infarction” generally results from atherosclerosis of the coronary arteries, often with superimposed coronary thrombosis. It may be divided into two major types: transmural infarcts, in which myocardial necrosis involves the full thickness of the ventricular wall, and subendocardial (nontransmural) infarcts, in which the necrosis involves the subendocardium, the intramural myocardium, or both, without extending all the way through the ventricular wall to the epicardium. Myocardial infarction is known to cause both a change in hemodynamic effects and an alteration in structure in the damaged and healthy zones of the heart. Thus, for example, myocardial infarction reduces the maximum cardiac output and the stroke volume of the heart. Also associated with myocardial infarction is a stimulation of the DNA synthesis occurring in the interstice as well as an increase in the formation of collagen in the areas of the heart not affected.

As a result of the increased stress or strain placed on the heart in prolonged hypertension due, for example, to the increased total peripheral resistance, cardiac hypertrophy has long been associated with “hypertension”. A characteristic of the ventricle that becomes hypertrophic as a result of chronic pressure overload is an impaired diastolic performance. Fouad *et al.*, J. Am. Coll. Cardiol., 4: 1500-1506 (1984); Smith *et al.*, J. Am. Coll. Cardiol., 5: 869-874 (1985). A prolonged left ventricular relaxation has been detected in early essential hypertension, in spite of normal or supranormal systolic function. Hartford *et al.*, Hypertension, 6: 329-338 (1984). However, there is no close parallelism between blood pressure levels and cardiac hypertrophy. Although improvement in left ventricular function in response to antihypertensive therapy has been reported in humans, patients variously treated with a diuretic (hydrochlorothiazide), a β -blocker (propranolol), or a calcium channel blocker (diltiazem), have shown reversal of left ventricular hypertrophy, without improvement in diastolic function. Inouye *et al.*, Am. J. Cardiol., 53: 1583-7 (1984).

Another complex cardiac disease associated with cardiac hypertrophy is “hypertrophic cardiomyopathy”. This condition is characterized by a great diversity of morphologic, functional, and clinical features (Maron *et al.*, N. Engl. J. Med., 316: 780-789 (1987); Spirito *et al.*, N. Engl. J. Med., 320: 749-755 (1989); Louie and Edwards, Prog. Cardiovasc. Dis., 36: 275-308 (1994); Wigle *et al.*, Circulation, 92: 1680-1692 (1995)), the heterogeneity of which is accentuated by the fact that it afflicts patients of all ages. Spirito *et al.*, N. Engl. J. Med., 336: 775-785 (1997). The causative factors of hypertrophic cardiomyopathy are also diverse and little understood. In general, mutations in genes encoding sarcomeric proteins are associated with hypertrophic cardiomyopathy. Recent data suggest that β -myosin heavy chain mutations may account for approximately 30 to 40 percent of cases of familial hypertrophic cardiomyopathy. Watkins *et al.*, N. Engl. J. Med., 326: 1108-1114 (1992); Schwartz *et al.*, Circulation, 91: 532-540 (1995); Marian and Roberts, Circulation, 92: 1336-1347 (1995); Thierfelder *et al.*, Cell, 77: 701-712

(1994); Watkins *et al.*, Nat. Gen., 11: 434-437 (1995). Besides β -myosin heavy chain, other locations of genetic mutations include cardiac troponin T, alpha topomyosin, cardiac myosin binding protein C, essential myosin light chain, and regulatory myosin light chain. See, Malik and Watkins, Curr. Opin. Cardiol., 12: 295-302 (1997).

Supravalvular "aortic stenosis" is an inherited vascular disorder characterized by narrowing of the ascending aorta, but other arteries, including the pulmonary arteries, may also be affected. Untreated aortic stenosis may lead to increased intracardiac pressure resulting in myocardial hypertrophy and eventually heart failure and death. The pathogenesis of this disorder is not fully understood, but hypertrophy and possibly hyperplasia of medial smooth muscle are prominent features of this disorder. It has been reported that molecular variants of the elastin gene are involved in the development and pathogenesis of aortic stenosis. U.S. Patent No. 5,650,282 issued July 22, 1997.

"Valvular regurgitation" occurs as a result of heart diseases resulting in disorders of the cardiac valves. Various diseases, like rheumatic fever, can cause the shrinking or pulling apart of the valve orifice, while other diseases may result in endocarditis, an inflammation of the endocardium or lining membrane of the atrioventricular orifices and operation of the heart. Defects such as the narrowing of the valve stenosis or the defective closing of the valve result in an accumulation of blood in the heart cavity or regurgitation of blood past the valve. If uncorrected, prolonged valvular stenosis or insufficiency may result in cardiac hypertrophy and associated damage to the heart muscle, which may eventually necessitate valve replacement.

The treatment of all these, and other cardiovascular, endothelial and angiogenic disorders, which may or may not be accompanied by cardiac hypertrophy, is encompassed by the present invention.

The terms "cancer", "cancerous", and "malignant" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include but are not limited to, carcinoma including adenocarcinoma, lymphoma, blastoma, melanoma, sarcoma, and leukemia. More particular examples of such cancers include squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, gastrointestinal cancer, Hodgkin's and non-Hodgkin's lymphoma, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer such as hepatic carcinoma and hepatoma, bladder cancer, breast cancer, colon cancer, colorectal cancer, endometrial carcinoma, salivary gland carcinoma, kidney cancer such as renal cell carcinoma and Wilms' tumors, basal cell carcinoma, melanoma, prostate cancer, vulval cancer, thyroid cancer, testicular cancer, esophageal cancer, and various types of head and neck cancer. The preferred cancers for treatment herein are breast, colon, lung, melanoma, ovarian, and others involving vascular tumors as noted above.

The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (*e.g.*, ^{131}I , ^{125}I , ^{90}Y , and ^{186}Re), chemotherapeutic agents, and toxins such as enzymatically active toxins of bacterial, fungal, plant, or animal origin, or fragments thereof.

A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents, folic acid antagonists, anti-metabolites of nucleic acid metabolism, antibiotics, pyrimidine analogs, 5-fluorouracil, cisplatin, purine nucleosides, amines, amino acids, triazol nucleosides, or corticosteroids. Specific examples include Adriamycin, Doxorubicin, 5-Fluorouracil,

Cytosine arabinoside ("Ara-C"), Cyclophosphamide, Thiotepa, Busulfan, Cytosin, Taxol, Toxotere, Methotrexate, Cisplatin, Melphalan, Vinblastine, Bleomycin, Etoposide, Ifosfamide, Mitomycin C, Mitoxantrone, Vincristine, Vinorelbine, Carboplatin, Teniposide, Daunomycin, Carminomycin, Aminopterin, Dactinomycin, Mitomycins, Esperamicins (*see* U.S. Pat. No. 4,675,187), Melphalan, and other related nitrogen mustards. Also included in this definition are hormonal agents that act to regulate or inhibit hormone action on tumors, such as tamoxifen and onapristone.

A "growth-inhibitory agent" when used herein refers to a compound or composition that inhibits growth of a cell, such as an Wnt-overexpressing cancer cell, either *in vitro* or *in vivo*. Thus, the growth-inhibitory agent is one which significantly reduces the percentage of malignant cells in S phase. Examples of growth-inhibitory agents include agents that block cell cycle progression (at a place other than S phase), such as agents that induce G1 arrest and M-phase arrest. Classical M-phase blockers include the vincas (vincristine and vinblastine), taxol, and topo II inhibitors such as doxorubicin, daunorubicin, etoposide, and bleomycin. Those agents that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such as tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C. Further information can be found in The Molecular Basis of Cancer, Mendelsohn and Israel, eds., Chapter 1, entitled "Cell cycle regulation, oncogenes, and antineoplastic drugs" by Murakami *et al.* (WB Saunders: Philadelphia, 1995), especially p. 13. Additional examples include tumor necrosis factor (TNF), an antibody capable of inhibiting or neutralizing the angiogenic activity of acidic or basic FGF or hepatocyte growth factor (HGF), an antibody capable of inhibiting or neutralizing the coagulant activities of tissue factor, protein C, or protein S (*see*, WO 91/01753, published 21 February 1991), or an antibody capable of binding to HER2 receptor (WO 89/06692), such as the 4D5 antibody (and functional equivalents thereof) (*e.g.*, WO 92/22653).

"Treatment" is an intervention performed with the intention of preventing the development or altering the pathology of a cardiovascular, endothelial, and angiogenic disorder. The concept of treatment is used in the broadest sense, and specifically includes the prevention (prophylaxis), moderation, reduction, and curing of cardiovascular, endothelial, and angiogenic disorders of any stage. Accordingly, "treatment" refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) or ameliorate a cardiovascular, endothelial, and angiogenic disorder such as hypertrophy. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented. The disorder may result from any cause, including idiopathic, cardiotrophic, or myotrophic causes, or ischemia or ischemic insults, such as myocardial infarction.

"Chronic" administration refers to administration of the agent(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial effect, such as an anti-hypertrophic effect, for an extended period of time.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, sheep, pigs, etc. Preferably, the mammal is human.

Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

The phrase "cardiovascular, endothelial or angiogenic agents" refers generically to any drug that acts in treating cardiovascular, endothelial, and angiogenic disorders. Examples of cardiovascular agents are those that promote vascular homeostasis by modulating blood pressure, heart rate, heart contractility, and endothelial and smooth muscle biology, all of which factors have a role in cardiovascular disease. Specific examples of these include angiotensin-II receptor antagonists; endothelin receptor antagonists such as, for example, BOSENTANTM and MOXONODINTM; interferon-gamma (IFN- γ); des-aspartate-angiotensin I; thrombolytic agents, *e.g.*, streptokinase, urokinase, t-PA, and a t-PA variant specifically designed to have longer half-life and very high fibrin specificity, TNK t-PA (a T103N, N117Q, KHRR(296-299)AAAA t-PA variant, Keyt *et al.*, Proc. Natl. Acad. Sci. USA, 91: 3670-3674 (1994)); inotropic or hypertensive agents such as digoxigenin and β -adrenergic receptor blocking agents, *e.g.*, propranolol, timolol, tertalolol, carteolol, nadolol, betaxolol, penbutolol, acetobutolol, atenolol, metoprolol, and carvedilol; angiotensin converting enzyme (ACE) inhibitors, *e.g.*, quinapril, captopril, enalapril, ramipril, benazepril, fosinopril, and lisinopril; diuretics, *e.g.*, chlorothiazide, hydrochlorothiazide, hydroflumethazide, methylchlothiazide, benzthiazide, dichlorphenamide, acetazolamide, and indapamide; and calcium channel blockers, *e.g.*, diltiazem, nifedipine, verapamil, nicardipine. One preferred category of this type is a therapeutic agent used for the treatment of cardiac hypertrophy or of a physiological condition instrumental in the development of cardiac hypertrophy, such as elevated blood pressure, aortic stenosis, or myocardial infarction.

"Angiogenic agents" and "endothelial agents" are active agents that promote angiogenesis and/or endothelial cell growth, or, if applicable, vasculogenesis. This would include factors that accelerate wound healing, such as growth hormone, insulin-like growth factor-I (IGF-I), VEGF, VIGF, PDGF, epidermal growth factor (EGF), CTGF and members of its family, FGF, and TGF- α and TGF- β .

"Angiostatic agents" are active agents that inhibit angiogenesis or vasculogenesis or otherwise inhibit or prevent growth of cancer cells. Examples include antibodies or other antagonists to angiogenic agents as defined above, such as antibodies to VEGF. They additionally include cytotherapeutic agents such as cytotoxic agents, chemotherapeutic agents, growth-inhibitory agents, apoptotic agents, and other agents to treat cancer, such as anti-HER-2, anti-CD20, and other bioactive and organic chemical agents.

In a pharmacological sense, in the context of the present invention, a "therapeutically effective amount" of an active agent such as a PRO polypeptide or agonist or antagonist thereto or an anti-PRO antibody, refers to an amount effective in the treatment of a cardiovascular, endothelial or angiogenic disorder in a mammal and can be determined empirically.

As used herein, an "effective amount" of an active agent such as a PRO polypeptide or agonist or antagonist thereto or an anti-PRO antibody, refers to an amount effective for carrying out a stated purpose, wherein such amounts may be determined empirically for the desired effect.

The terms "PRO polypeptide" and "PRO" as used herein and when immediately followed by a numerical designation refer to various polypeptides, wherein the complete designation (*i.e.*, PRO/number) refers to specific polypeptide sequences as described herein. The terms "PRO/number polypeptide" and "PRO/number" wherein the term "number" is provided as an actual numerical designation as used herein encompass native sequence polypeptides and polypeptide variants (which are further defined herein). The PRO polypeptides described herein

may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods.

A "native sequence PRO polypeptide" comprises a polypeptide having the same amino acid sequence as the corresponding PRO polypeptide derived from nature. Such native sequence PRO polypeptides can be isolated from nature or can be produced by recombinant or synthetic means. The term "native sequence PRO polypeptide" specifically encompasses naturally-occurring truncated or secreted forms of the specific PRO polypeptide (e.g., an extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide. In various embodiments of the invention, the native sequence PRO polypeptides disclosed herein are mature or full-length native sequence polypeptides comprising the full-length amino acids sequences shown in the accompanying figures. Start and stop codons are shown in bold font and underlined in the figures. However, while the PRO polypeptide disclosed in the accompanying figures are shown to begin with methionine residues designated herein as amino acid position 1 in the figures, it is conceivable and possible that other methionine residues located either upstream or downstream from the amino acid position 1 in the figures may be employed as the starting amino acid residue for the PRO polypeptides.

The PRO polypeptide "extracellular domain" or "ECD" refers to a form of the PRO polypeptide which is essentially free of the transmembrane and cytoplasmic domains. Ordinarily, a PRO polypeptide ECD will have less than 1% of such transmembrane and/or cytoplasmic domains and preferably, will have less than 0.5% of such domains. It will be understood that any transmembrane domains identified for the PRO polypeptides of the present invention are identified pursuant to criteria routinely employed in the art for identifying that type of hydrophobic domain. The exact boundaries of a transmembrane domain may vary but most likely by no more than about 5 amino acids at either end of the domain as initially identified herein. Optionally, therefore, an extracellular domain of a PRO polypeptide may contain from about 5 or fewer amino acids on either side of the transmembrane domain/extracellular domain boundary as identified in the Examples or specification and such polypeptides, with or without the associated signal peptide, and nucleic acid encoding them, are contemplated by the present invention.

The approximate location of the "signal peptides" of the various PRO polypeptides disclosed herein are shown in the present specification and/or the accompanying figures. It is noted, however, that the C-terminal boundary of a signal peptide may vary, but most likely by no more than about 5 amino acids on either side of the signal peptide C-terminal boundary as initially identified herein, wherein the C-terminal boundary of the signal peptide may be identified pursuant to criteria routinely employed in the art for identifying that type of amino acid sequence element (e.g., Nielsen *et al.*, Prot. Eng., 10:1-6 (1997) and von Heinje *et al.*, Nucl. Acids Res., 14:4683-4690 (1986)). Moreover, it is also recognized that, in some cases, cleavage of a signal sequence from a secreted polypeptide is not entirely uniform, resulting in more than one secreted species. These mature polypeptides, where the signal peptide is cleaved within no more than about 5 amino acids on either side of the C-terminal boundary of the signal peptide as identified herein, and the polynucleotides encoding them, are contemplated by the present invention.

"PRO polypeptide variant" means an active PRO polypeptide as defined above or below having at least about 80% amino acid sequence identity with a full-length native sequence PRO polypeptide sequence as disclosed

herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Such PRO polypeptide variants include, for instance, PRO polypeptides wherein one or more amino acid residues are added, or deleted, at the N- or C-terminus of the full-length native amino acid sequence. Ordinarily, a PRO polypeptide variant will have at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, PRO variant polypeptides are at least about 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150 or 200 amino acids in length and alternatively at least about 300 amino acids in length, or more.

"Percent (%) amino acid sequence identity" with respect to the PRO polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in a PRO sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are obtained as described below by using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc., and the source code shown in Table 1 has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Table 1. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

For purposes herein, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scored as identical matches by the sequence alignment program

ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. As examples of % amino acid sequence identity calculations, Tables 2-3 demonstrate how to calculate the % amino acid sequence identity of the amino acid sequence designated "Comparison Protein" to the amino acid sequence designated "PRO".

Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described above using the ALIGN-2 sequence comparison computer program. However, % amino acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul *et al.*, Nucleic Acids Res., 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov>, or otherwise obtained from the National Institute of Health, Bethesda, MD. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10, minimum low complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final gapped alignment = 25 and scoring matrix = BLOSUM62.

In situations where NCBI-BLAST2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A.

In addition, % amino acid sequence identity may also be determined using the WU-BLAST-2 computer program (Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, *i.e.*, the adjustable parameters, are set with the following values: overlap span = 1, overlap fraction = 0.125, word threshold (T) = 11, and scoring matrix = BLOSUM62. For purposes herein, a % amino acid sequence identity value is determined by dividing (a) the number of matching identical amino acids residues between the amino acid sequence of the PRO polypeptide of interest having a sequence derived from the native PRO polypeptide and the comparison amino acid sequence of interest (*i.e.*, the sequence against which the PRO polypeptide of interest is being compared which may be a PRO variant polypeptide) as determined by WU-BLAST-2 by (b) the total number of amino acid residues of the PRO polypeptide of interest. For example, in the statement "a polypeptide comprising an amino acid sequence A which

has or having at least 80% amino acid sequence identity to the amino acid sequence B", the amino acid sequence A is the comparison amino acid sequence of interest and the amino acid sequence B is the amino acid sequence of the PRO polypeptide of interest.

"PRO variant polynucleotide" or "PRO variant nucleic acid sequence" means a nucleic acid molecule which encodes an active PRO polypeptide as defined below and which has at least about 80% nucleic acid sequence identity with a nucleotide acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, a PRO variant polynucleotide will have at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity with a nucleic acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal sequence, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Variants do not encompass the native nucleotide sequence.

Ordinarily, PRO variant polynucleotides are at least about 30, 60, 90, 120, 150, 180, 210, 240, 270, 300, 450, or 600 nucleotides in length and alternatively at least about 900 nucleotides in length, or more.

"Percent (%) nucleic acid sequence identity" with respect to the PRO polypeptide-encoding nucleic acid sequences identified herein is defined as the percentage of nucleotides in a candidate sequence that are identical with the nucleotides in a PRO polypeptide-encoding nucleic acid sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared. For purposes herein, however, % nucleic acid sequence identity values are obtained as described below by using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc., and the source code shown in Table 1 has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Table 1. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

For purposes herein, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that

has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

$$100 \text{ times the fraction } W/Z$$

5

where W is the number of nucleotides scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C. As examples of % nucleic acid sequence identity calculations, Tables 4-5 demonstrate how to calculate the % nucleic acid sequence identity of the nucleic acid sequence designated "Comparison DNA" to the nucleic acid sequence designated "PRO-DNA".

10

Unless specifically stated otherwise, all % nucleic acid sequence identity values used herein are obtained as described above using the ALIGN-2 sequence comparison computer program. However, % nucleic acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul *et al.*, Nucleic Acids Res., 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov> or otherwise obtained from the National Institute of Health, Bethesda, MD. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10, minimum low complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final gapped alignment = 25 and scoring matrix = BLOSUM62.

15

20

In situations where NCBI-BLAST2 is employed for sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

25

$$100 \text{ times the fraction } W/Z$$

where W is the number of nucleotides scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C.

30

In addition, % nucleic acid sequence identity values may also be generated using the WU-BLAST-2 computer program (Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, *i.e.*, the adjustable parameters, are set with the following values: overlap span = 1, overlap fraction = 0.125, word threshold (T) = 11, and scoring matrix = BLOSUM62. For purposes herein, a % nucleic acid sequence identity value is determined by dividing (a)

35

the number of matching identical nucleotides between the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest having a sequence derived from the native sequence PRO polypeptide-encoding nucleic acid and the comparison nucleic acid molecule of interest (*i.e.*, the sequence against which the PRO polypeptide-encoding nucleic acid molecule of interest is being compared which may be a variant PRO polynucleotide) as determined by WU-BLAST-2 by (b) the total number of nucleotides of the PRO polypeptide-encoding nucleic acid molecule of interest. For example, in the statement "an isolated nucleic acid molecule comprising a nucleic acid sequence A which has or having at least 80% nucleic acid sequence identity to the nucleic acid sequence B", the nucleic acid sequence A is the comparison nucleic acid molecule of interest and the nucleic acid sequence B is the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest.

In other embodiments, PRO variant polynucleotides are nucleic acid molecules that encode an active PRO polypeptide and which are capable of hybridizing, preferably under stringent hybridization and wash conditions, to nucleotide sequences encoding the full-length PRO polypeptide as shown in the specification herein and accompanying figures. PRO variant polypeptides may be those that are encoded by a PRO variant polynucleotide.

"Isolated", when used to describe the various polypeptides disclosed herein, means a polypeptide that has been identified and separated and/or recovered from a component of its natural environment. Preferably, the isolated polypeptide is free of association with all components with which it is naturally associated. Contaminant components of its natural environment are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes polypeptide *in situ* within recombinant cells, since at least one component of the PRO natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

An "isolated" nucleic acid molecule encoding a PRO polypeptide or an "isolated" nucleic acid molecule encoding an anti-PRO antibody is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the PRO-encoding nucleic acid or the natural source of the anti-PRO-encoding nucleic acid. Preferably, the isolated nucleic acid is free of association with all components with which it is naturally associated. An isolated PRO-encoding nucleic acid molecule or an isolated anti-PRO-encoding nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated nucleic acid molecules therefore are distinguished from the PRO-encoding nucleic acid molecule or from the anti-PRO-encoding nucleic acid molecule as it exists in natural cells. However, an isolated nucleic acid molecule encoding a PRO polypeptide or an isolated nucleic acid molecule encoding an anti-PRO antibody includes PRO-nucleic acid molecules or anti-PRO-nucleic acid molecules contained in cells that ordinarily express PRO polypeptides or anti-PRO antibodies where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked

coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize, for example, promoters, polyadenylation signals, and enhancers.

Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a PRO polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in the same reading frame. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

"Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature that can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, *see*, Ausubel *et al.*, Current Protocols in Molecular Biology (Wiley Interscience Publishers, 1995).

"Stringent conditions" or "high-stringency conditions", as defined herein, may be identified by those that: (1) employ low ionic strength and high temperature for washing, for example, 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42°C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55°C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55°C.

"Moderately-stringent conditions" may be identified as described by Sambrook *et al.*, Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Press, 1989), and include the use of washing solution and hybridization conditions (*e.g.*, temperature, ionic strength, and % SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters

in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

The modifier "epitope-tagged" when used herein refers to a chimeric polypeptide comprising a PRO polypeptide fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, yet is short enough such that it does not interfere with activity of the polypeptide to which it is fused. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues).

"Active" or "activity" in the context of PRO variants refers to form(s) of PRO proteins that retain the biologic and/or immunologic activities of a native or naturally-occurring PRO polypeptide.

"Biological activity" in the context of a molecule that antagonizes a PRO polypeptide that can be identified by the screening assays disclosed herein (*e.g.*, an organic or inorganic small molecule, peptide, etc.) is used to refer to the ability of such molecules to bind or complex with the PRO polypeptide identified herein, or otherwise interfere with the interaction of the PRO polypeptide with other cellular proteins or otherwise inhibits the transcription or translation of the PRO polypeptide. Particularly preferred biological activity includes cardiac hypertrophy, activity that acts on systemic disorders that affect vessels, such as diabetes mellitus, as well as diseases of the arteries, capillaries, veins, and/or lymphatics, and cancer.

The term "antagonist" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes one or more of the biological activities of a native PRO polypeptide disclosed herein, for example, if applicable, its mitogenic or angiogenic activity. Antagonists of a PRO polypeptide may act by interfering with the binding of a PRO polypeptide to a cellular receptor, by incapacitating or killing cells that have been activated by a PRO polypeptide, or by interfering with vascular endothelial cell activation after binding of a PRO polypeptide to a cellular receptor. All such points of intervention by a PRO polypeptide antagonist shall be considered equivalent for purposes of this invention. The antagonists inhibit the mitogenic, angiogenic, or other biological activity of PRO polypeptides, and thus are useful for the treatment of diseases or disorders characterized by undesirable excessive neovascularization, including by way of example tumors, and especially solid malignant tumors, rheumatoid arthritis, psoriasis, atherosclerosis, diabetic and other retinopathies, retrolental fibroplasia, age-related macular degeneration, neovascular glaucoma, hemangiomas, thyroid hyperplasias (including Grave's disease), corneal and other tissue transplantation, and chronic inflammation. The antagonists also are useful for the treatment of diseases or disorders characterized by undesirable excessive vascular permeability, such as edema associated with brain tumors, ascites associated with malignancies, Meigs' syndrome, lung inflammation, nephrotic syndrome, pericardial effusion (such as that associated with pericarditis), and pleural effusion. In a similar manner, the term "agonist" is used in the broadest sense and includes any molecule that mimics a biological activity of a native PRO polypeptide disclosed herein. Suitable agonist or antagonist molecules specifically include agonist or antagonist antibodies or antibody fragments, fragments, or amino acid sequence variants of native PRO polypeptides, peptides, small organic molecules, etc.

A "small molecule" is defined herein to have a molecular weight below about 500 daltons.

The term "PRO polypeptide receptor" as used herein refers to a cellular receptor for a PRO polypeptide, ordinarily a cell-surface receptor found on vascular endothelial cells, as well as variants thereof that retain the ability to bind a PRO polypeptide.

"Antibodies" (Abs) and "immunoglobulins" (Igs) are glycoproteins having the same structural characteristics. While antibodies exhibit binding specificity to a specific antigen, immunoglobulins include both antibodies and other antibody-like molecules that lack antigen specificity. Polypeptides of the latter kind are, for example, produced at low levels by the lymph system and at increased levels by myelomas. The term "antibody" is used in the broadest sense and specifically covers, without limitation, intact monoclonal antibodies, polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies) formed from at least two intact antibodies, and antibody fragments, so long as they exhibit the desired biological activity.

"Native antibodies" and "native immunoglobulins" are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light-chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light- and heavy-chain variable domains.

The term "variable" refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody to and for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called complementarity-determining regions (CDRs) or hypervariable regions both in the light-chain and the heavy-chain variable domains. The more highly conserved portions of variable domains are called the framework regions (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a β -sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the β -sheet structure. The CDRs in each chain are held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of antibodies. See, Kabat *et al.*, NIH Publ. No.91-3242, Vol. I, pages 647-669 (1991). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

"Antibody fragments" comprise a portion of an intact antibody, preferably the antigen-binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies (Zapata *et al.*, Protein Eng., 8(10): 1057-1062 (1995)); single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, whose name reflects its ability to crystallize

readily. Pepsin treatment yields an $F(ab')_2$ fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment that contains a complete antigen-recognition and -binding site. This region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the V_H - V_L dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. $F(ab')_2$ antibody fragments originally were produced as pairs of Fab' fragments that have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM; and several of these may be further divided into subclasses (isotypes), *e.g.*, IgG1, IgG2, IgG3, IgG4, IgA, and IgA2. The heavy-chain constant domains that correspond to the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations that typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma culture, uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler *et al.*, Nature, 256: 495 (1975), or may be made by recombinant DNA methods (*see, e.g.*, U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, Nature, 352: 624-628 (1991) and Marks *et al.*, J. Mol. Biol., 222: 581-597 (1991), for example.

The monoclonal antibodies herein specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity. U.S. Patent No. 4,816,567; Morrison *et al.*, Proc. Natl. Acad. Sci. USA, 81: 6851-6855 (1984).

"Humanized" forms of non-human (*e.g.*, murine) antibodies are chimeric immunoglobulins, immunoglobulin chains, or fragments thereof (such as Fv, Fab, Fab', F(ab')₂, or other antigen-binding subsequences of antibodies) that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a CDR of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv FR residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and maximize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin sequence. The humanized antibody preferably also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, Nature, 321: 522-525 (1986); Reichmann *et al.*, Nature, 332: 323-329 (1988); and Presta, Curr. Op. Struct. Biol., 2: 593-596 (1992). The humanized antibody includes a PRIMATIZED™ antibody wherein the antigen-binding region of the antibody is derived from an antibody produced by immunizing macaque monkeys with the antigen of interest.

"Single-chain Fv" or "sFv" antibody fragments comprise the V_H and V_L domains of an antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains that enables the sFv to form the desired structure for antigen binding. For a review of sFv see, Pluckthun in The Pharmacology of Monoclonal Antibodies, Vol. 113, Rosenberg and Moore, eds. (Springer-Verlag: New York, 1994), pp. 269-315.

The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) in the same polypeptide chain (V_H - V_L). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger *et al.*, Proc. Natl. Acad. Sci. USA, 90: 6444-6448 (1993).

An "isolated" antibody is one that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would interfere

with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody *in situ* within recombinant cells, since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

An antibody that "specifically binds to" or is "specific for" a particular polypeptide or an epitope on a particular polypeptide is one that binds to that particular polypeptide or epitope on a particular polypeptide without substantially binding to any other polypeptide or polypeptide epitope.

The word "label" when used herein refers to a detectable compound or other composition that is conjugated directly or indirectly to the antibody so as to generate a "labeled" antibody. The label may be detectable by itself (e.g., radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition that is detectable. Radionuclides that can serve as detectable labels include, for example, I-131, I-123, I-125, Y-90, Re-188, At-211, Cu-67, Bi-212, and Pd-109. The label may also be a non-detectable entity such as a toxin.

By "solid phase" is meant a non-aqueous matrix to which an antibody of the present invention can adhere. Examples of solid phases encompassed herein include those formed partially or entirely of glass (e.g., controlled pore glass), polysaccharides (e.g., agarose), polyacrylamides, polystyrene, polyvinyl alcohol and silicones. In certain embodiments, depending on the context, the solid phase can comprise the well of an assay plate; in others it is a purification column (e.g., an affinity chromatography column). This term also includes a discontinuous solid phase of discrete particles, such as those described in U.S. Patent No. 4,275,149.

A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant that is useful for delivery of a drug (such as the PRO polypeptide or antibodies thereto disclosed herein) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

As used herein, the term "immunoadhesin" designates antibody-like molecules that combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity that is other than the antigen recognition and binding site of an antibody (i.e., is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD, or IgM.

As shown below, Table 1 provides the complete source code for the ALIGN-2 sequence comparison computer program. This source code may be routinely compiled for use on a UNIX operating system to provide

the ALIGN-2 sequence comparison computer program.

In addition, Tables 2-5 show hypothetical exemplifications for using the below described method to determine % amino acid sequence identity (Tables 2-3) and % nucleic acid sequence identity (Tables 4-5) using the ALIGN-2 sequence comparison computer program, wherein "PRO" represents the amino acid sequence of a hypothetical PRO polypeptide of interest, "Comparison Protein" represents the amino acid sequence of a polypeptide against which the "PRO" polypeptide of interest is being compared, "PRO-DNA" represents a hypothetical PRO-encoding nucleic acid sequence of interest, "Comparison DNA" represents the nucleotide sequence of a nucleic acid molecule against which the "PRO-DNA" nucleic acid molecule of interest is being compared, "X", "Y", and "Z" each represent different hypothetical amino acid residues and "N", "L" and "V" each represent different hypothetical nucleotides.

Table 1

```

/*
 *
 * C-C increased from 12 to 15
 * Z is average of EQ
 * B is average of ND
 * match with stop is _M; stop-stop = 0; J (joker) match = 0
 */
#define _M      -8      /* value of a match with a stop */

int _day[26][26] = {
/* A B C D E F G H I J K L M N O P Q R S T U V W X Y Z */
/* A */ { 2, 0, -2, 0, 0, -4, 1, -1, -1, 0, -1, -2, -1, 0, _M, 1, 0, -2, 1, 1, 0, 0, -6, 0, -3, 0 },
/* B */ { 0, 3, -4, 3, 2, -5, 0, 1, -2, 0, 0, -3, -2, 2, _M, -1, 1, 0, 0, 0, 0, -2, -5, 0, -3, 1 },
/* C */ { -2, -4, 15, -5, -5, -4, -3, -3, -2, 0, -5, -6, -5, -4, _M, -3, -5, -4, 0, -2, 0, -2, -8, 0, 0, -5 },
/* D */ { 0, 3, -5, 4, 3, -6, 1, 1, -2, 0, 0, -4, -3, 2, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 2 },
/* E */ { 0, 2, -5, 3, 4, -5, 0, 1, -2, 0, 0, -3, -2, 1, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 3 },
/* F */ { -4, -5, -4, -6, -5, 9, -5, -2, 1, 0, -5, 2, 0, -4, _M, -5, -5, -4, -3, -3, 0, -1, 0, 0, 7, -5 },
/* G */ { 1, 0, -3, 1, 0, -5, 5, -2, -3, 0, -2, -4, -3, 0, _M, -1, -1, -3, 1, 0, 0, -1, -7, 0, -5, 0 },
/* H */ { -1, 1, -3, 1, 1, -2, -2, 6, -2, 0, 0, -2, -2, 2, _M, 0, 3, 2, -1, -1, 0, -2, -3, 0, 0, 2 },
/* I */ { -1, -2, -2, -2, -2, 1, -3, -2, 5, 0, -2, 2, 2, -2, _M, -2, -2, -2, -1, 0, 0, 4, -5, 0, -1, -2 },
/* J */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0 },
/* K */ { -1, 0, -5, 0, 0, -5, -2, 0, -2, 0, 5, -3, 0, 1, _M, -1, 1, 3, 0, 0, 0, -2, -3, 0, -4, 0 },
/* L */ { -2, -3, -6, -4, -3, 2, -4, -2, 2, 0, -3, 6, 4, -3, _M, -3, -2, -3, -3, -1, 0, 2, -2, 0, -1, -2 },
/* M */ { -1, -2, -5, -3, -2, 0, -3, -2, 2, 0, 0, 4, 6, -2, _M, -2, -1, 0, -2, -1, 0, 2, -4, 0, -2, -1 },
/* N */ { 0, 2, -4, 2, 1, -4, 0, 2, -2, 0, 1, -3, -2, 2, _M, -1, 1, 0, 1, 0, 0, -2, -4, 0, -2, 1 },
/* O */ { _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M },
/* P */ { 1, -1, -3, -1, -1, -5, -1, 0, -2, 0, -1, -3, -2, -1, _M, 6, 0, 0, 1, 0, 0, -1, -6, 0, -5, 0 },
/* Q */ { 0, 1, -5, 2, 2, -5, -1, 3, -2, 0, 1, -2, -1, 1, _M, 0, 4, 1, -1, -1, 0, -2, -5, 0, -4, 3 },
/* R */ { -2, 0, -4, -1, -1, -4, -3, 2, -2, 0, 3, -3, 0, 0, _M, 0, 1, 6, 0, -1, 0, -2, 2, 0, -4, 0 },
/* S */ { 1, 0, 0, 0, 0, -3, 1, -1, -1, 0, 0, -3, -2, 1, _M, 1, -1, 0, 2, 1, 0, -1, -2, 0, -3, 0 },
/* T */ { 1, 0, -2, 0, 0, -3, 0, -1, 0, 0, 0, -1, -1, 0, _M, 0, -1, -1, 1, 3, 0, 0, -5, 0, -3, 0 },
/* U */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0 },
/* V */ { 0, -2, -2, -2, -2, -1, -1, -2, 4, 0, -2, 2, 2, -2, _M, -1, -2, -2, -1, 0, 0, 4, -6, 0, -2, -2 },
/* W */ { -6, -5, -8, -7, -7, 0, -7, -3, -5, 0, -3, -2, -4, -4, _M, -6, -5, 2, -2, -5, 0, -6, 17, 0, 0, -6 },
/* X */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0 },
/* Y */ { -3, -3, 0, -4, -4, 7, -5, 0, -1, 0, -4, -1, -2, -2, _M, -5, -4, -4, -3, -3, 0, -2, 0, 0, 10, -4 },
/* Z */ { 0, 1, -5, 2, 3, -5, 0, 2, -2, 0, 0, -2, -1, 1, _M, 0, 3, 0, 0, 0, 0, -2, -6, 0, -4, 4 }
};

```


Table 1 (cont')

```

/*
*/
#include <stdio.h>
#include <ctype.h>

#define MAXJMP      16      /* max jumps in a diag */
#define MAXGAP      24      /* don't continue to penalize gaps larger than this */
#define JMPS        1024    /* max jmps in an path */
#define MX          4       /* save if there's at least MX-1 bases since last jmp */

#define DMAT        3       /* value of matching bases */
#define DMIS        0       /* penalty for mismatched bases */
#define DINS0       8       /* penalty for a gap */
#define DINS1       1       /* penalty per base */
#define PINS0       8       /* penalty for a gap */
#define PINS1       4       /* penalty per residue */

struct jmp {
    short          n[MAXJMP];    /* size of jmp (neg for dely) */
    unsigned short x[MAXJMP];    /* base no. of jmp in seq x */
};                               /* limits seq to 2^16 -1 */

struct diag {
    int            score;        /* score at last jmp */
    long           offset;       /* offset of prev block */
    short          jmp;         /* current jmp index */
    struct jmp     jp;          /* list of jmps */
};

struct path {
    int            spc;          /* number of leading spaces */
    short          n[JMPS];      /* size of jmp (gap) */
    int            x[JMPS];      /* loc of jmp (last elem before gap) */
};

char             *ofile;        /* output file name */
char             *namex[2];     /* seq names: getseqs( ) */
char             *prog;         /* prog name for err msgs */
char             *seqx[2];      /* seqs: getseqs( ) */
int              dmax;          /* best diag: nw( ) */
int              dmax0;         /* final diag */
int              dna;           /* set if dna: main( ) */
int              endgaps;       /* set if penalizing end gaps */
int              gapx, gapy;     /* total gaps in seqs */
int              len0, len1;     /* seq lens */
int              ngapx, ngapy;   /* total size of gaps */
int              smax;          /* max score: nw( ) */
int              *xbm;          /* bitmap for matching */
long             offset;        /* current offset in jmp file */
struct diag      *dx;           /* holds diagonals */
struct path      pp[2];         /* holds path for seqs */

char             *calloc( ), *malloc( ), *index( ), *strcpy( );
char             *getseq( ), *g_calloc( );

```

Table 1 (cont')

```

/* Needleman-Wunsch alignment program
*
* usage: progs file1 file2
* where file1 and file2 are two dna or two protein sequences.
* The sequences can be in upper- or lower-case and may contain ambiguity
* Any lines beginning with ';', '>' or '<' are ignored
* Max file length is 65535 (limited by unsigned short x in the jmp struct)
* A sequence with 1/3 or more of its elements ACGTU is assumed to be DNA
* Output is in the file "align.out"
*
* The program may create a tmp file in /tmp to hold info about traceback.
* Original version developed under BSD 4.3 on a vax 8650
*/
#include "nw.h"
#include "day.h"

static _dbval[26] = {
    1,14,2,13,0,0,4,11,0,0,12,0,3,15,0,0,0,5,6,8,8,7,9,0,10,0
};

static _pbval[26] = {
    1, 2|(1<<('D'-'A'))|(1<<('N'-'A')), 4, 8, 16, 32, 64,
    128, 256, 0xFFFFFFFF, 1<<10, 1<<11, 1<<12, 1<<13, 1<<14,
    1<<15, 1<<16, 1<<17, 1<<18, 1<<19, 1<<20, 1<<21, 1<<22,
    1<<23, 1<<24, 1<<25|(1<<('E'-'A'))|(1<<('Q'-'A'))
};

main(ac, av)
int    ac;
char   *av[];
{
    prog = av[0];
    if (ac != 3) {
        fprintf(stderr, "usage: %s file1 file2\n", prog);
        fprintf(stderr, "where file1 and file2 are two dna or two protein sequences.\n");
        fprintf(stderr, "The sequences can be in upper- or lower-case\n");
        fprintf(stderr, "Any lines beginning with ';', '>' or '<' are ignored\n");
        fprintf(stderr, "Output is in the file \"align.out\"\n");
        exit(1);
    }
    namex[0] = av[1];
    namex[1] = av[2];
    seqx[0] = getseq(namex[0], &len0);
    seqx[1] = getseq(namex[1], &len1);
    xbm = (dna)? _dbval : _pbval;

    endgaps = 0; /* 1 to penalize endgaps */
    ofile = "align.out"; /* output file */

    nw(); /* fill in the matrix, get the possible jumps */
    readjumps(); /* get the actual jumps */
    print(); /* print stats, alignment */

    cleanup(0); /* unlink any tmp files */
}

```

main

Table 1 (cont')

```

/* do the alignment, return best score: main( )
* dna: values in Fitch and Smith, PNAS, 80, 1382-1386, 1983
* pro: PAM 250 values
* When scores are equal, we prefer mismatches to any gap, prefer
* a new gap to extending an ongoing gap, and prefer a gap in seqx
* to a gap in seq y.
*/
nw( )
{
    char      *px, *py;           /* seqs and ptrs */
    int       *ndely, *dely;      /* keep track of dely */
    int       ndelx, delx;        /* keep track of delx */
    int       *tmp;              /* for swapping row0, row1 */
    int       mis;               /* score for each type */
    int       ins0, ins1;         /* insertion penalties */
    register  id;                /* diagonal index */
    register  ij;                /* jmp index */
    register  *col0, *col1;       /* score for curr, last row */
    register  xx, yy;            /* index into seqs */

    dx = (struct diag *)g_calloc("to get diags", len0+len1+1, sizeof(struct diag));

    ndely = (int *)g_calloc("to get ndely", len1+1, sizeof(int));
    dely = (int *)g_calloc("to get dely", len1+1, sizeof(int));
    col0 = (int *)g_calloc("to get col0", len1+1, sizeof(int));
    col1 = (int *)g_calloc("to get col1", len1+1, sizeof(int));
    ins0 = (dna)? DINS0 : PINS0;
    ins1 = (dna)? DINS1 : PINS1;

    smax = -10000;
    if (endgaps) {
        for (col0[0] = dely[0] = -ins0, yy = 1; yy <= len1; yy++) {
            col0[yy] = dely[yy] = col0[yy-1] - ins1;
            ndely[yy] = yy;
        }
        col0[0] = 0;          /* Waterman Bull Math Biol 84 */
    }
    else
        for (yy = 1; yy <= len1; yy++)
            dely[yy] = -ins0;

    /* fill in match matrix
    */
    for (px = seqx[0], xx = 1; xx <= len0; px++, xx++) {
        /* initialize first entry in col
        */
        if (endgaps) {
            if (xx == 1)
                col1[0] = delx = -(ins0+ins1);
            else
                col1[0] = delx = col0[0] - ins1;
            ndelx = xx;
        }
        else {
            col1[0] = 0;
            delx = -ins0;
            ndelx = 0;
        }
    }
}

```

Table 1 (cont')

...nw

```

for (py = seqx[1], yy = 1; yy <= len1; py++, yy++) {
    mis = col0[yy-1];
    if (dna)
        mis += (xbm[*px-'A']&xbm[*py-'A'])? DMAT : DMIS;
    else
        mis += _day[*px-'A'][*py-'A'];

    /* update penalty for del in x seq;
     * favor new del over ongong del
     * ignore MAXGAP if weighting endgaps
     */
    if (endgaps || ndely[yy] < MAXGAP) {
        if (col0[yy] - ins0 >= dely[yy]) {
            dely[yy] = col0[yy] - (ins0+ins1);
            ndely[yy] = 1;
        } else {
            dely[yy] -= ins1;
            ndely[yy]++;
        }
    } else {
        if (col0[yy] - (ins0+ins1) >= dely[yy]) {
            dely[yy] = col0[yy] - (ins0+ins1);
            ndely[yy] = 1;
        } else
            ndely[yy]++;
    }

    /* update penalty for del in y seq;
     * favor new del over ongong del
     */
    if (endgaps || ndelx < MAXGAP) {
        if (col1[yy-1] - ins0 >= delx) {
            delx = col1[yy-1] - (ins0+ins1);
            ndelx = 1;
        } else {
            delx -= ins1;
            ndelx++;
        }
    } else {
        if (col1[yy-1] - (ins0+ins1) >= delx) {
            delx = col1[yy-1] - (ins0+ins1);
            ndelx = 1;
        } else
            ndelx++;
    }

    /* pick the maximum score; we're favoring
     * mis over any del and delx over dely
     */

```

Table 1 (cont')

...nw

```

id = xx - yy + len1 - 1;
if (mis >= delx && mis >= dely[yy])
    col1[yy] = mis;
else if (delx >= dely[yy]) {
    col1[yy] = delx;
    ij = dx[id].ijmp;
    if (dx[id].jp.n[0] && (!dna || (ndelx >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writeimps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
    }
    dx[id].jp.n[ij] = ndelx;
    dx[id].jp.x[ij] = xx;
    dx[id].score = delx;
}
else {
    col1[yy] = dely[yy];
    ij = dx[id].ijmp;
    if (dx[id].jp.n[0] && (!dna || (ndely[yy] >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writeimps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
    }
    dx[id].jp.n[ij] = -ndely[yy];
    dx[id].jp.x[ij] = xx;
    dx[id].score = dely[yy];
}
if (xx == len0 && yy < len1) {
    /* last col
    */
    if (endgaps)
        col1[yy] -= ins0+ins1*(len1-yy);
    if (col1[yy] > smax) {
        smax = col1[yy];
        dmax = id;
    }
}
}
if (endgaps && xx < len0)
    col1[yy-1] -= ins0+ins1*(len0-xx);
if (col1[yy-1] > smax) {
    smax = col1[yy-1];
    dmax = id;
}
tmp = col0; col0 = col1; col1 = tmp;
}
(void) free((char *)ndely);
(void) free((char *)dely);
(void) free((char *)col0);
(void) free((char *)col1);
}

```

Table 1 (cont')

```

/*
 *
 * print() -- only routine visible outside this module
 *
 * static:
 * getmat() -- trace back best path, count matches: print()
 * pr_align() -- print alignment of described in array p[]: print()
 * dumpblock() -- dump a block of lines with numbers, stars: pr_align()
 * nums() -- put out a number line: dumpblock()
 * putline() -- put out a line (name, [num], seq, [num]): dumpblock()
 * stars() -- put a line of stars: dumpblock()
 * stripname() -- strip any path and prefix from a seqname
 */

#include "nw.h"

#define SPC 3
#define P_LINE 256 /* maximum output line */
#define P_SPC 3 /* space between name or num and seq */

extern _day[26][26];
int olen; /* set output line length */
FILE *fx; /* output file */

print()
{
    int lx, ly, firstgap, lastgap; /* overlap */

    if ((fx = fopen(ofile, "w")) == 0) {
        fprintf(stderr, "%s: can't write %s\n", prog, ofile);
        cleanup(1);
    }
    fprintf(fx, "<first sequence: %s (length = %d)\n", namex[0], len0);
    fprintf(fx, "<second sequence: %s (length = %d)\n", namex[1], len1);
    olen = 60;
    lx = len0;
    ly = len1;
    firstgap = lastgap = 0;
    if (dmax < len1 - 1) { /* leading gap in x */
        pp[0].spc = firstgap = len1 - dmax - 1;
        ly -= pp[0].spc;
    }
    else if (dmax > len1 - 1) { /* leading gap in y */
        pp[1].spc = firstgap = dmax - (len1 - 1);
        lx -= pp[1].spc;
    }
    if (dmax0 < len0 - 1) { /* trailing gap in x */
        lastgap = len0 - dmax0 - 1;
        lx -= lastgap;
    }
    else if (dmax0 > len0 - 1) { /* trailing gap in y */
        lastgap = dmax0 - (len0 - 1);
        ly -= lastgap;
    }
    getmat(lx, ly, firstgap, lastgap);
    pr_align();
}

```

print

Table 1 (cont')

```

/*
 * trace back the best path, count matches
 */
static
getmat(lx, ly, firstgap, lastgap)                                     getmat
{
    int      lx, ly;
    int      firstgap, lastgap;
    /* "core" (minus endgaps) */
    /* leading trailing overlap */

    int      nm, i0, i1, siz0, siz1;
    char      outx[32];
    double    pct;
    register  n0, n1;
    register char *p0, *p1;

    /* get total matches, score
    */
    i0 = i1 = siz0 = siz1 = 0;
    p0 = seqx[0] + pp[1].spc;
    p1 = seqx[1] + pp[0].spc;
    n0 = pp[1].spc + 1;
    n1 = pp[0].spc + 1;

    nm = 0;
    while ( *p0 && *p1 ) {
        if (siz0) {
            p1++;
            n1++;
            siz0--;
        }
        else if (siz1) {
            p0++;
            n0++;
            siz1--;
        }
        else {
            if (xbm[*p0-'A']&xbm[*p1-'A'])
                nm++;
            if (n0++ == pp[0].x[i0])
                siz0 = pp[0].n[i0++];
            if (n1++ == pp[1].x[i1])
                siz1 = pp[1].n[i1++];
            p0++;
            p1++;
        }
    }

    /* pct homology:
    * if penalizing endgaps, base is the shorter seq
    * else, knock off overhangs and take shorter core
    */
    if (endgaps)
        lx = (len0 < len1)? len0 : len1;
    else
        lx = (lx < ly)? lx : ly;
    pct = 100.*(double)nm/(double)lx;
    fprintf(fx, "\n");
    fprintf(fx, "< %d match%s in an overlap of %d: %.2f percent similarity\n",
        nm, (nm == 1)? "" : "es", lx, pct);

```

Table 1 (cont')

```

fprintf(fx, "< gaps in first sequence: %d", gapx);
if (gapx) {
    (void) sprintf(outx, " (%d %s%s)",
        ngapx, (dna)? "base":"residue", (ngapx == 1)? ":" : "s");
    fprintf(fx, "%s", outx);

    fprintf(fx, ", gaps in second sequence: %d", gapy);
    if (gapy) {
        (void) sprintf(outx, " (%d %s%s)",
            ngapy, (dna)? "base":"residue", (ngapy == 1)? ":" : "s");
        fprintf(fx, "%s", outx);
    }
    if (dna)
        fprintf(fx,
            "\n< score: %d (match = %d, mismatch = %d, gap penalty = %d + %d per base)\n",
            smax, DMAT, DMIS, DINS0, DINS1);
    else
        fprintf(fx,
            "\n< score: %d (Dayhoff PAM 250 matrix, gap penalty = %d + %d per residue)\n",
            smax, PINS0, PINS1);
    if (endgaps)
        fprintf(fx,
            "< endgaps penalized. left endgap: %d %s%s, right endgap: %d %s%s\n",
            firstgap, (dna)? "base" : "residue", (firstgap == 1)? ":" : "s",
            lastgap, (dna)? "base" : "residue", (lastgap == 1)? ":" : "s");
    else
        fprintf(fx, "< endgaps not penalized\n");
}

static      nm;          /* matches in core -- for checking */
static      lmax;        /* lengths of stripped file names */
static      ij[2];       /* jmp index for a path */
static      nc[2];       /* number at start of current line */
static      ni[2];       /* current elem number -- for gapping */
static      siz[2];
static char  *ps[2];      /* ptr to current element */
static char  *po[2];      /* ptr to next output char slot */
static char  out[2][P_LINE]; /* output line */
static char  star[P_LINE]; /* set by stars() */

/*
 * print alignment of described in struct path pp[]
 */
static
pr_align()
{
    int      nn;          /* char count */
    int      more;
    register i;

    for (i = 0, lmax = 0; i < 2; i++) {
        nn = stripname(namex[i]);
        if (nn > lmax)
            lmax = nn;

        nc[i] = 1;
        ni[i] = 1;
        siz[i] = ij[i] = 0;
        ps[i] = seqx[i];
        po[i] = out[i];
    }
}

```

...getmat

pr_align

Table 1 (cont')**...pr_align**

```

for (nn = nm = 0, more = 1; more; ) {
    for (i = more = 0; i < 2; i++) {
        /*
         * do we have more of this sequence?
         */
        if (!*ps[i])
            continue;

        more++;

        if (pp[i].spc) { /* leading space */
            *po[i]++ = ' ';
            pp[i].spc--;
        }
        else if (siz[i]) { /* in a gap */
            *po[i]++ = '-';
            siz[i]--;
        }
        else { /* we're putting a seq element
                */
            *po[i] = *ps[i];
            if (islower(*ps[i]))
                *ps[i] = toupper(*ps[i]);

            po[i]++;
            ps[i]++;

            /*
             * are we at next gap for this seq?
             */
            if (ni[i] == pp[i].x[ij[i]]) {
                /*
                 * we need to merge all gaps
                 * at this location
                 */
                siz[i] = pp[i].n[ij[i]] + +;
                while (ni[i] == pp[i].x[ij[i]])
                    siz[i] += pp[i].n[ij[i]] + +;
            }
            ni[i]++;
        }
    }
    if (++nn == olen || !more && nn) {
        dumpblock();
        for (i = 0; i < 2; i++)
            po[i] = out[i];
        nn = 0;
    }
}

/*
 * dump a block of lines, including numbers, stars: pr_align( )
 */
static
dumpblock( )
{
    register i;

    for (i = 0; i < 2; i++)
        *po[i]-- = '\0';
}

```

dumpblock

Table 1 (cont')

...dumpblock

```

(void) putc('\n', fx);
for (i = 0; i < 2; i++) {
    if (*out[i] && (*out[i] != ' ' || *(po[i]) != ' ')) {
        if (i == 0)
            nums(i);
        if (i == 0 && *out[1])
            stars( );
        putline(i);
        if (i == 0 && *out[1])
            fprintf(fx, star);
        if (i == 1)
            nums(i);
    }
}

/*
 * put out a number line: dumpblock( )
 */
static
nums(ix)
int ix; /* index in out[] holding seq line */
{
    char nline[P_LINE];
    register i, j;
    register char *pn, *px, *py;

    for (pn = nline, i = 0; i < lmax+P_SPC; i++, pn++)
        *pn = ' ';
    for (i = nc[ix], py = out[ix]; *py; py++, pn++) {
        if (*py == ' ' || *py == '-')
            *pn = ' ';
        else {
            if (i%10 == 0 || (i == 1 && nc[ix] != 1)) {
                j = (i < 0)? -i : i;
                for (px = pn; j; j /= 10, px--)
                    *px = j%10 + '0';
                if (i < 0)
                    *px = '-';
            }
            else
                *pn = ' ';
            i++;
        }
    }
    *pn = '\0';
    nc[ix] = i;
    for (pn = nline; *pn; pn++)
        (void) putc(*pn, fx);
    (void) putc('\n', fx);
}

/*
 * put out a line (name, [num], seq, [num]): dumpblock( )
 */
static
putline(ix)
int ix;
{

```

nums

putline

Table 1 (cont')

...putline

```

int          i;
register char *px;

for (px = namex[ix], i = 0; *px && *px != ':'; px++, i++)
    (void) putc(*px, fx);
for (; i < lmax+P_SPC; i++)
    (void) putc(' ', fx);

/* these count from 1:
 * ni[] is current element (from 1)
 * nc[] is number at start of current line
 */
for (px = out[ix]; *px; px++)
    (void) putc(*px&0x7F, fx);
(void) putc('\n', fx);
}

/*
 * put a line of stars (seqs always in out[0], out[1]): dumpblock( )
 */
static
stars( )
{
    int          i;
    register char *p0, *p1, cx, *px;

    if (!*out[0] || (*out[0] == ' ' && *(po[0]) == ' ') ||
        !*out[1] || (*out[1] == ' ' && *(po[1]) == ' '))
        return;
    px = star;
    for (i = lmax+P_SPC; i; i--)
        *px++ = ' ';

    for (p0 = out[0], p1 = out[1]; *p0 && *p1; p0++, p1++) {
        if (isalpha(*p0) && isalpha(*p1)) {
            if (xbm[*p0-'A']&xbm[*p1-'A']) {
                cx = '*';
                nm++;
            }
            else if (!dna && _day[*p0-'A'][*p1-'A'] > 0)
                cx = '.';
            else
                cx = ' ';
        }
        else
            cx = ' ';
        *px++ = cx;
    }
    *px++ = '\n';
    *px = '\0';
}

```

stars

Table 1 (cont')

```

/*
 * strip path or prefix from pn, return len: pr_align( )
 */
static
stripname(pn)                                stripname
    char    *pn;    /* file name (may be path) */
{
    register char    *px, *py;

    py = 0;
    for (px = pn; *px; px++)
        if (*px == '/')
            py = px + 1;
    if (py)
        (void) strcpy(pn, py);
    return(strlen(pn));
}

```

Table 1 (cont')

```

/*
 * cleanup( ) -- cleanup any tmp file
 * getseq( ) -- read in seq, set dna, len, maxlen
 * g_calloc( ) -- calloc( ) with error checkin
 * readjumps( ) -- get the good jumps, from tmp file if necessary
 * writejumps( ) -- write a filled array of jumps to a tmp file: nw( )
 */
#include "nw.h"
#include <sys/file.h>

char    *jname = "/tmp/homgXXXXXX";          /* tmp file for jumps */
FILE    *fj;

int      cleanup( );                          /* cleanup tmp file */
long     lseek( );

/*
 * remove any tmp file if we blow
 */
cleanup(i)                                     cleanup
{
    int    i;
    if (fj)
        (void) unlink(jname);
    exit(i);
}

/*
 * read, return ptr to seq, set dna, len, maxlen
 * skip lines starting with ';', '<', or '>'
 * seq in upper or lower case
 */
char    *
getseq(file, len)                             getseq
{
    char    *file;    /* file name */
    int     *len;     /* seq len */

    char    line[1024], *pseq;
    register char *px, *py;
    int     natgc, tlen;
    FILE    *fp;

    if ((fp = fopen(file, "r")) == 0) {
        fprintf(stderr, "%s: can't read %s\n", prog, file);
        exit(1);
    }
    tlen = natgc = 0;
    while (fgets(line, 1024, fp)) {
        if (*line == ';' || *line == '<' || *line == '>')
            continue;
        for (px = line; *px != '\n'; px++)
            if (isupper(*px) || islower(*px))
                tlen++;
    }
    if ((pseq = malloc((unsigned)(tlen+6))) == 0) {
        fprintf(stderr, "%s: malloc( ) failed to get %d bytes for %s\n", prog, tlen+6, file);
        exit(1);
    }
    pseq[0] = pseq[1] = pseq[2] = pseq[3] = '\0';

```

Table 1 (cont')

...getseq

```

py = pseq + 4;
*len = tlen;
rewind(fp);

while (fgets(line, 1024, fp)) {
    if (*line == ';' || *line == '<' || *line == '>')
        continue;
    for (px = line; *px != '\n'; px++) {
        if (isupper(*px))
            *py++ = *px;
        else if (islower(*px))
            *py++ = toupper(*px);
        if (index("ATGCU", *(py-1)))
            natgc++;
    }
    *py++ = '\0';
    *py = '\0';
    (void) fclose(fp);
    dna = natgc > (tlen/3);
    return(pseq+4);
}

char *
g_calloc(msg, nx, sz)
char *msg;          /* program, calling routine */
int nx, sz;          /* number and size of elements */
{
    char *px, *calloc( );

    if ((px = calloc((unsigned)nx, (unsigned)sz)) == 0) {
        if (*msg) {
            fprintf(stderr, "%s: g_calloc( ) failed %s (n=%d, sz=%d)\n", prog, msg, nx, sz);
            exit(1);
        }
    }
    return(px);
}

/*
 * get final jmps from dx[] or tmp file, set pp[], reset dmax: main( )
 */
readjmps( )
{
    int fd = -1;
    int siz, i0, i1;
    register i, j, xx;

    if (fj) {
        (void) fclose(fj);
        if ((fd = open(jname, O_RDONLY, 0)) < 0) {
            fprintf(stderr, "%s: can't open( ) %s\n", prog, jname);
            cleanup(1);
        }
    }

    for (i = i0 = i1 = 0, dmax0 = dmax, xx = len0; ; i++) {
        while (1) {
            for (j = dx[dmax].ijmp; j >= 0 && dx[dmax].jp.x[j] >= xx; j--)
                ;

```

g_calloc

readjmps

Table 1 (cont')

...readjumps

```

    if (j < 0 && dx[dmax].offset && fj) {
        (void) lseek(fd, dx[dmax].offset, 0);
        (void) read(fd, (char *)&dx[dmax].jp, sizeof(struct jmp));
        (void) read(fd, (char *)&dx[dmax].offset, sizeof(dx[dmax].offset));
        dx[dmax].ijmp = MAXJMP-1;
    }
    else
        break;
}
if (i >= JMPS) {
    fprintf(stderr, "%s: too many gaps in alignment\n", prog);
    cleanup(1);
}
if (j >= 0) {
    siz = dx[dmax].jp.n[j];
    xx = dx[dmax].jp.x[j];
    dmax += siz;
    if (siz < 0) { /* gap in second seq */
        pp[1].n[i1] = -siz;
        xx += siz;
        /* id = xx - yy + len1 - 1
        */
        pp[1].x[i1] = xx - dmax + len1 - 1;
        gapy++;
        ngapy -= siz;
        /* ignore MAXGAP when doing endgaps */
        siz = (-siz < MAXGAP || endgaps)? -siz : MAXGAP;
        i1++;
    }
    else if (siz > 0) { /* gap in first seq */
        pp[0].n[i0] = siz;
        pp[0].x[i0] = xx;
        gapx++;
        ngapx += siz;
        /* ignore MAXGAP when doing endgaps */
        siz = (siz < MAXGAP || endgaps)? siz : MAXGAP;
        i0++;
    }
}
else
    break;
}

/* reverse the order of jumps
*/
for (j = 0, i0--; j < i0; j++, i0--) {
    i = pp[0].n[j]; pp[0].n[j] = pp[0].n[i0]; pp[0].n[i0] = i;
    i = pp[0].x[j]; pp[0].x[j] = pp[0].x[i0]; pp[0].x[i0] = i;
}
for (j = 0, i1--; j < i1; j++, i1--) {
    i = pp[1].n[j]; pp[1].n[j] = pp[1].n[i1]; pp[1].n[i1] = i;
    i = pp[1].x[j]; pp[1].x[j] = pp[1].x[i1]; pp[1].x[i1] = i;
}
if (fd >= 0)
    (void) close(fd);
if (fj) {
    (void) unlink(jname);
    fj = 0;
    offset = 0;
}

```

Table 1 (cont')

```

/*
 * write a filled jmp struct offset of the prev one (if any): nw( )
 */
writejumps(ix)                                writejumps
{
    int      ix;
    char      *mktemp( );

    if (!fj) {
        if (mktemp(jname) < 0) {
            fprintf(stderr, "%s: can't mktemp( ) %s\n", prog, jname);
            cleanup(1);
        }
        if ((fj = fopen(jname, "w")) == 0) {
            fprintf(stderr, "%s: can't write %s\n", prog, jname);
            exit(1);
        }
    }
    (void) fwrite((char *)&dx[ix].jp, sizeof(struct jmp), 1, fj);
    (void) fwrite((char *)&dx[ix].offset, sizeof(dx[ix].offset), 1, fj);
}

```


Table 2

PRO	XXXXXXXXXXXXXXXXXX	(Length = 15 amino acids)
Comparison Protein	XXXXXXXXYYYYYYY	(Length = 12 amino acids)

5 % amino acid sequence identity =
 (the number of identically matching amino acid residues between the two polypeptide sequences as determined by
 ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) =
 5 divided by 15 = 33.3%

Table 3

PRO	XXXXXXXXXXXXX	(Length = 10 amino acids)
Comparison Protein	XXXXXXXXYYYYYYZZYZ	(Length = 15 amino acids)

10
 15 % amino acid sequence identity =
 (the number of identically matching amino acid residues between the two polypeptide sequences as determined by
 ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) =
 5 divided by 10 = 50%

Table 4

20 PRO-DNA	NNNNNNNNNNNNNN	(Length = 14 nucleotides)
Comparison DNA	NNNNNNLLLLLLLL	(Length = 16 nucleotides)

25 % nucleic acid sequence identity =
 (the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-
 2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) =
 6 divided by 14 = 42.9%

Table 5

30 PRO-DNA	NNNNNNNNNNNN	(Length = 12 nucleotides)
Comparison DNA	NNNNLLLVV	(Length = 9 nucleotides)

35 % nucleic acid sequence identity =
 (the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-
 2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) =
 4 divided by 12 = 33.3%

5.2. Compositions and Methods of the Invention

5.2.1. PRO Variants

In addition to the full-length native sequence PRO polypeptides described herein, it is contemplated that PRO variants can be prepared. PRO variants can be prepared by introducing appropriate nucleotide changes into the PRO DNA, and/or by synthesis of the desired PRO polypeptide. Those skilled in the art will appreciate that amino acid changes may alter post-translational processes of the PRO polypeptide such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

Variations in the native full-length sequence PRO polypeptide or in various domains of the PRO polypeptide described herein, can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Patent No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the PRO polypeptide that results in a change in the amino acid sequence of the PRO polypeptide as compared with the native sequence PRO polypeptide. Optionally the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the PRO polypeptide. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the PRO polypeptide with that of homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, *i.e.*, conservative amino acid replacements. Insertions or deletions may optionally be in the range of about 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the full-length or mature native sequence.

In particular embodiments, conservative substitutions of interest are shown in Table 6 under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial changes, denominated exemplary substitutions in Table 6, or as further described below in reference to amino acid classes, are introduced and the products screened.

Table 6

	Original Residue	Exemplary Substitutions	Preferred Substitutions
5	Ala (A)	val; leu; ile	val
	Arg (R)	lys; gln; asn	lys
	Asn (N)	gln; his; lys; arg	gln
	Asp (D)	glu	glu
	Cys (C)	ser	ser
10	Gln (Q)	asn	asn
	Glu (E)	asp	asp
	Gly (G)	pro; ala	ala
	His (H)	asn; gln; lys; arg	arg
	Ile (I)	leu; val; met; ala; phe; norleucine	leu
15	Leu (L)	norleucine; ile; val; met; ala; phe	ile
	Lys (K)	arg; gln; asn	arg
	Met (M)	leu; phe; ile	leu
	Phe (F)	leu; val; ile; ala; tyr	leu
	Pro (P)	ala	ala
20	Ser (S)	thr	thr
	Thr (T)	ser	ser
	Trp (W)	tyr; phe	tyr
	Tyr (Y)	trp; phe; thr; ser	phe
	Val (V)	ile; leu; met; phe; ala; norleucine	leu

Substantial modifications in function or immunological identity of the PRO polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophilic: cys, ser, thr;
- (3) acidic: asp, glu;
- (4) basic: asn, gln, his, lys, arg;
- (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic: trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, more preferably, into the remaining (non-conserved) sites.

The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter *et al.*, Nucl. Acids Res., 13:4331 (1986); Zoller *et al.*, Nucl. Acids Res., 10:6487 (1987)], cassette mutagenesis [Wells *et al.*, Gene, 34:315 (1985)], restriction selection mutagenesis [Wells *et al.*, Philos. Trans. R. Soc. London SerA, 317:415

(1986)] or other known techniques can be performed on the cloned DNA to produce the PRO variant DNA.

Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant [Cunningham and Wells, Science, 244: 1081-1085 (1989)]. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, The Proteins, (W.H. Freeman & Co., N.Y.); Chothia, J. Mol. Biol., 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

5.2.2. Modifications of PRO Polypeptides

Covalent modifications of PRO polypeptides are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of a PRO polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C- terminal residues of the PRO polypeptide. Derivatization with bifunctional agents is useful, for instance, for crosslinking the PRO polypeptide to a water-insoluble support matrix or surface for use in the method for purifying anti-PRO antibodies, and vice-versa. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidyl)propionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio]propioimide.

Other modifications include deamidation of glutamyl and asparagyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains [T.E. Creighton, Proteins: Structure and Molecular Properties, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

Another type of covalent modification of the PRO polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in the native sequence PRO polypeptide (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means), and/or adding one or more glycosylation sites that are not present in the native sequence PRO polypeptide. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

Addition of glycosylation sites to the PRO polypeptide may be accomplished by altering the amino acid sequence. The alteration may be made, for example, by the addition of, or substitution by, one or more serine or threonine residues to the native sequence PRO polypeptide (for O-linked glycosylation sites). The PRO amino acid sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the PRO polypeptide at preselected bases such that codons are generated that will translate into the desired amino

acids.

Another means of increasing the number of carbohydrate moieties on the PRO polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, *e.g.*, in WO 87/05330 published 11 September 1987, and in Aplin and Wriston, CRC Crit. Rev. Biochem., pp. 259-306 (1981).

Removal of carbohydrate moieties present on the PRO polypeptide may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, *et al.*, Arch. Biochem. Biophys., 259:52 (1987) and by Edge *et al.*, Anal. Biochem., 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura *et al.*, Meth. Enzymol., 138:350 (1987).

Another type of covalent modification of the PRO polypeptide comprises linking the PRO polypeptide to one of a variety of nonproteinaceous polymers, *e.g.*, polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

The PRO polypeptide of the present invention may also be modified in a way to form a chimeric molecule comprising the PRO polypeptide fused to another, heterologous polypeptide or amino acid sequence.

In one embodiment, such a chimeric molecule comprises a fusion of the PRO polypeptide with a protein transduction domain which targets the PRO polypeptide for delivery to various tissues and more particularly across the brain blood barrier, using, for example, the protein transduction domain of human immunodeficiency virus TAT protein (Schwarze *et al.*, 1999, *Science* 285: 1569-72).

In another embodiment, such a chimeric molecule comprises a fusion of the PRO polypeptide with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl- terminus of the PRO polypeptide. The presence of such epitope-tagged forms of the PRO polypeptide can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the PRO polypeptide to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-His) or poly-histidine-glycine (poly-His-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field *et al.*, Mol. Cell. Biol., 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan *et al.*, Molecular and Cellular Biology, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky *et al.*, Protein Engineering, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp *et al.*, BioTechnology, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin *et al.*, Science, 255:192-194 (1992)]; an α -tubulin epitope peptide [Skinner *et al.*, J. Biol. Chem., 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth *et al.*, Proc. Natl. Acad. Sci. USA, 87:6393-6397 (1990)].

In an alternative embodiment, the chimeric molecule may comprise a fusion of the PRO polypeptide with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an "immunoadhesin"), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions

preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of a PRO polypeptide in place of at least one variable region within an Ig molecule. In a particularly preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions *see* also, U.S. Patent No. 5,428,130 issued June 27, 1995.

5 5.2.3. Preparation of the PRO Polypeptide

The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO polypeptides. In particular, cDNAs encoding PRO polypeptides have been identified and isolated, as disclosed in further detail in the Examples below. It is noted that proteins produced in separate expression rounds may be given different PRO numbers but the UNQ number is unique for any given DNA and the encoded protein, and will not be changed. However, for sake of simplicity, in the present specification the protein encoded by the PRO DNA as well as all further native homologues and variants included in the foregoing definition of PRO polypeptides, will be referred to as "PRO" regardless of their origin or mode of preparation.

10 The description below relates primarily to production of PRO polypeptides by culturing cells transformed or transfected with a vector containing nucleic acid encoding PRO polypeptides. It is, of course, contemplated that alternative methods that are well known in the art may be employed to prepare the PRO polypeptide. For instance, the PRO polypeptide sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques. *See, e.g., Stewart et al., Solid-Phase Peptide Synthesis* (W.H. Freeman Co.: San Francisco, CA, 1969); Merrifield, *J. Am. Chem. Soc.*, **85**: 2149-2154 (1963). *In vitro* protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, with an Applied Biosystems Peptide Synthesizer (Foster City, CA) using manufacturer's instructions. Various portions of the PRO polypeptide may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the full-length PRO polypeptide.

15 5.2.3.1. Isolation of DNA Encoding PRO Polypeptides

25 DNA encoding the PRO polypeptide may be obtained from a cDNA library prepared from tissue believed to possess the mRNA encoding the PRO polypeptide and to express it at a detectable level. Accordingly, DNAs encoding the human PRO polypeptide can be conveniently obtained from cDNA libraries prepared from human tissues, such as described in the Examples. The gene encoding the PRO polypeptide may also be obtained from a genomic library or by oligonucleotide synthesis.

30 Libraries can be screened with probes (such as antibodies to the PRO polypeptide or oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook *et al., supra*. An alternative means to isolate the gene encoding the PRO polypeptide is to use PCR methodology. Sambrook *et al., supra*; Dieffenbach *et al., PCR Primer: A Laboratory Manual* (New York: Cold

Spring Harbor Laboratory Press, 1995).

The Examples below describe techniques for screening a cDNA library. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art, and include the use of radiolabels like ^{32}P -labeled ATP, biotinylation, or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook *et al.*, *supra*.

Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases. Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined through sequence alignment using computer software programs such as ALIGN, DNASTar, and INHERIT, which employ various algorithms to measure homology.

Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook *et al.*, *supra*, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

5.2.3.2. Selection and Transformation of Host Cells

Host cells are transfected or transformed with expression or cloning vectors described herein for PRO polypeptide production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH, and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in Mammalian Cell Biotechnology: A Practical Approach, M. Butler, ed. (IRL Press, 1991) and Sambrook *et al.*, *supra*.

Methods of transfection are known to the ordinarily skilled artisan, for example, CaPO_4 treatment and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook *et al.*, *supra*, or electroporation is generally used for prokaryotes or other cells that contain substantial cell-wall barriers. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw *et al.*, Gene, **23**: 315 (1983) and WO 89/05859 published 29 June 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, Virology, **52**:456-457 (1978) can be employed. General aspects of mammalian cell host system transformations have been described in U.S. Patent No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen *et al.*, J. Bact., **130**: 946 (1977) and Hsiao *et al.*, Proc. Natl. Acad. Sci. (USA), **76**: 3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, *e.g.*, polybrene or polyornithine, may also be used. For various techniques for

transforming mammalian cells, *see*, Keown *et al.*, Methods in Enzymology, 185: 527-537 (1990) and Mansour *et al.*, Nature, 336: 348-352 (1988).

Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include, but are not limited to, eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325); and K5 772 (ATCC 53,635). Other suitable prokaryotic host cells include Enterobacteriaceae such as *Escherichia*, *e.g.*, *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, *e.g.*, *Salmonella typhimurium*, *Serratia*, *e.g.*, *Serratia marcescans*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (*e.g.*, *B. licheniformis* 41P disclosed in DD 266,710 published 12 April 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For example, strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including *E. coli* W3110 strain 1A2, which has the complete genotype *tonA*; *E. coli* W3110 strain 9E4, which has the complete genotype *tonA ptr3*; *E. coli* W3110 strain 27C7 (ATCC 55,244), which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT kan^r*; *E. coli* W3110 strain 37D6, which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT rbs7 ilvG kan^r*; *E. coli* W3110 strain 40B4, which is strain 37D6 with a non-kanamycin resistant *degP* deletion mutation; and an *E. coli* strain having mutant periplasmic protease disclosed in U.S. Patent No. 4,946,783 issued 7 August 1990. Alternatively, *in vitro* methods of cloning, *e.g.*, PCR or other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for vectors encoding the PRO polypeptide. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* (Beach and Nurse, Nature, 290: 140 [1981]; EP 139,383 published 2 May 1985); *Kluyveromyces* hosts (U.S. Patent No. 4,943,529; Fleer *et al.*, Bio/Technology, 9: 968-975 (1991)) such as, *e.g.*, *K. lactis* (MW98-8C, CBS683, CBS4574; Louvencourt *et al.*, J. Bacteriol., 737 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilum* (ATCC 36,906; Van den Berg *et al.*, Bio/Technology, 8: 135 (1990)), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna *et al.*, J. Basic Microbiol., 28: 265-278 [1988]); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa* (Case *et al.*, Proc. Natl. Acad. Sci. USA, 76: 5259-5263 [1979]); *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published 31 October 1990); and filamentous fungi such as, *e.g.*, *Neurospora*, *Penicillium*, *Tolypocladium* (WO 91/00357 published 10 January 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance *et al.*, Biochem. Biophys. Res. Commun., 112: 284-289 [1983]; Tilburn *et al.*, Gene, 26: 205-221 [1983]; Yelton *et al.*, Proc. Natl. Acad. Sci. USA, 81: 1470-1474 [1984]) and *A. niger* (Kelly and Hynes, EMBO J., 4: 475-479 [1985]). Methylophilic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*,

Torulopsis, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts may be found in C. Anthony, The Biochemistry of Methylootrophs, 269 (1982).

Suitable host cells for the expression of nucleic acid encoding glycosylated PRO polypeptides are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells. Examples of useful mammalian host cell lines include Chinese hamster ovary (CHO) and COS cells. More specific examples include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, J. Gen. Virol., 36: 59 (1977)); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. USA, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod., 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

5.2.3.3. Selection and Use of a Replicable Vector

The nucleic acid (*e.g.*, cDNA or genomic DNA) encoding the PRO polypeptide may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally include, but are not limited to, one or more of a signal sequence if the sequence is to be secreted, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques that are known to the skilled artisan.

The PRO polypeptide may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the DNA encoding the PRO polypeptide that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, *e.g.*, the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders, the latter described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published 4 April 1990), or the signal described in WO 90/13646 published 15 November 1990. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 μ plasmid origin

is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV, or BPV) are useful for cloning vectors in mammalian cells.

Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, *e.g.*, ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, *e.g.*, the gene encoding D-alanine racemase for *Bacilli*.

An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the nucleic acid encoding the PRO polypeptide such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub *et al.*, Proc. Natl. Acad. Sci. USA, 77: 4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7. Stinchcomb *et al.*, Nature, 282: 39 (1979); Kingsman *et al.*, Gene, 7: 141 (1979); Tschemper *et al.*, Gene, 10: 157 (1980). The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1. Jones, Genetics, 85: 12 (1977).

Expression and cloning vectors usually contain a promoter operably linked to the nucleic acid sequence encoding the PRO polypeptide to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the β -lactamase and lactose promoter systems (Chang *et al.*, Nature, 275: 615 (1978); Goeddel *et al.*, Nature, 281: 544 (1979)), alkaline phosphatase, a tryptophan (*trp*) promoter system (Goeddel, Nucleic Acids Res., 8: 4057 (1980); EP 36,776), and hybrid promoters such as the *tac* promoter (deBoer *et al.*, Proc. Natl. Acad. Sci. USA, 80: 21-25 (1983)). Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding the PRO polypeptide.

Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase (Hitzeman *et al.*, J. Biol. Chem., 255: 2073 (1980)) or other glycolytic enzymes (Hess *et al.*, J. Adv. Enzyme Reg., 7: 149 (1968); Holland, Biochemistry, 17: 4900 (1978)), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters that are inducible promoters having the additional advantage of transcription controlled by growth conditions are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

PRO nucleic acid transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus, and Simian Virus 40 (SV40); by heterologous mammalian promoters, *e.g.*, the actin

promoter or an immunoglobulin promoter; and by heat-shock promoters, provided such promoters are compatible with the host cell systems.

Transcription of a DNA encoding the PRO polypeptide by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the sequence coding for PRO polypeptides, but is preferably located at a site 5' from the promoter.

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding the PRO polypeptide.

Still other methods, vectors, and host cells suitable for adaptation to the synthesis of the PRO polypeptide in recombinant vertebrate cell culture are described in Gething *et al.*, Nature, 293: 620-625 (1981); Mantei *et al.*, Nature, 281: 40-46 (1979); EP 117,060; and EP 117,058.

5.2.3.4. Detecting Gene Amplification/Expression

Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA (Thomas, Proc. Natl. Acad. Sci. USA, 77:5201-5205 (1980)), dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native-sequence PRO polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to DNA encoding the PRO polypeptide and encoding a specific antibody epitope.

5.2.3.5. Purification of PRO Polypeptides

Forms of PRO polypeptides may be recovered from culture medium or from host cell lysates. If

membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g., TRITON-X™ 100) or by enzymatic cleavage. Cells employed in expression of nucleic acid encoding the PRO polypeptide can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell-lysing agents. It may be desired to purify the PRO polypeptide from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the PRO polypeptide. Various methods of protein purification may be employed and such methods are known in the art and described, for example, in Deutscher, Methods in Enzymology, 182 (1990); Scopes, Protein Purification: Principles and Practice (Springer-Verlag: New York, 1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular PRO polypeptide produced.

5.2.4. Uses of PRO Polypeptides

5.2.4.1. Assays for Cardiovascular, Endothelial, and Angiogenic Activity

Various assays can be used to test the polypeptide herein for cardiovascular, endothelial, and angiogenic activity. Such assays include those provided in the Examples below.

Assays for testing for endothelin antagonist activity, as disclosed in U.S. Pat. No. 5,773,414, include a rat heart ventricle binding assay where the polypeptide is tested for its ability to inhibit iodinated endothelin-1 binding in a receptor assay, an endothelin receptor binding assay testing for intact cell binding of radiolabeled endothelin-1 using rabbit renal artery vascular smooth muscle cells, an inositol phosphate accumulation assay where functional activity is determined in Rat-1 cells by measuring intra-cellular levels of second messengers, an arachidonic acid release assay that measures the ability of added compounds to reduce endothelin-stimulated arachidonic acid release in cultured vascular smooth muscles, *in vitro* (isolated vessel) studies using endothelium from male New Zealand rabbits, and *in vivo* studies using male Sprague-Dawley rats.

Assays for tissue generation activity include, without limitation, those described in WO 95/16035 (bone, cartilage, tendon); WO 95/05846 (nerve, neuronal), and WO 91/07491 (skin, endothelium).

Assays for wound-healing activity include, for example, those described in Winter, Epidermal Wound Healing, Maibach, HI and Rovee, DT, eds. (Year Book Medical Publishers, Inc., Chicago), pp. 71-112, as modified by the article of Eaglstein and Mertz, J. Invest. Dermatol., 71: 382-384 (1978).

An assay to screen for a test molecule relating to a PRO polypeptide that binds an endothelin B₁ (ETB₁) receptor polypeptide and modulates signal transduction activity involves providing a host cell transformed with a DNA encoding endothelin B₁ receptor polypeptide, exposing the cells to the test candidate, and measuring endothelin B₁ receptor signal transduction activity, as described, e.g., in U.S. Pat. No. 5,773,223.

There are several cardiac hypertrophy assays. *In vitro* assays include induction of spreading of adult rat cardiac myocytes. In this assay, ventricular myocytes are isolated from a single (male Sprague-Dawley) rat,

essentially following a modification of the procedure described in detail by Piper *et al.*, "Adult ventricular rat heart muscle cells" in Cell Culture Techniques in Heart and Vessel Research, H.M. Piper, ed. (Berlin: Springer-Verlag, 1990), pp. 36-60. This procedure permits the isolation of adult ventricular myocytes and the long-term culture of these cells in the rod-shaped phenotype. Phenylephrine and Prostaglandin F_{2α} (PGF_{2α}) have been shown to induce a spreading response in these adult cells. The inhibition of myocyte spreading induced by PGF_{2α} or PGF_{2α} analogs (e.g., fluprostenol) and phenylephrine by various potential inhibitors of cardiac hypertrophy is then tested.

One example of an *in vivo* assay is a test for inhibiting cardiac hypertrophy induced by fluprostenol *in vivo*. This pharmacological model tests the ability of the PRO polypeptide to inhibit cardiac hypertrophy induced in rats (e.g., male Wistar or Sprague-Dawley) by subcutaneous injection of fluprostenol (an agonist analog of PGF_{2α}). It is known that rats with pathologic cardiac hypertrophy induced by myocardial infarction have chronically elevated levels of extractable PGF_{2α} in their myocardium. Lai *et al.*, Am. J. Physiol. (Heart Circ. Physiol.), 271: H2197-H2208 (1996). Accordingly, factors that can inhibit the effects of fluprostenol on myocardial growth *in vivo* are potentially useful for treating cardiac hypertrophy. The effects of the PRO polypeptide on cardiac hypertrophy are determined by measuring the weight of heart, ventricles, and left ventricle (normalized by body weight) relative to fluprostenol-treated rats not receiving the PRO polypeptide.

Another example of an *in vivo* assay is the pressure-overload cardiac hypertrophy assay. For *in vivo* testing it is common to induce pressure-overload cardiac hypertrophy by constriction of the abdominal aorta of test animals. In a typical protocol, rats (e.g., male Wistar or Sprague-Dawley) are treated under anesthesia, and the abdominal aorta of each rat is narrowed down just below the diaphragm. Beznak M., Can. J. Biochem. Physiol., 33: 985-94 (1955). The aorta is exposed through a surgical incision, and a blunted needle is placed next to the vessel. The aorta is constricted with a ligature of silk thread around the needle, which is immediately removed and which reduces the lumen of the aorta to the diameter of the needle. This approach is described, for example, in Rossi *et al.*, Am. Heart J., 124: 700-709 (1992) and O'Rourke and Reibel, P.S.E.M.B., 200: 95-100 (1992).

In yet another *in vivo* assay, the effect on cardiac hypertrophy following experimentally induced myocardial infarction (MI) is measured. Acute MI is induced in rats by left coronary artery ligation and confirmed by electrocardiographic examination. A sham-operated group of animals is also prepared as control animals. Earlier data have shown that cardiac hypertrophy is present in the group of animals with MI, as evidenced by an 18% increase in heart weight-to-body weight ratio. Lai *et al.*, *supra*. Treatment of these animals with candidate blockers of cardiac hypertrophy, e.g., the PRO polypeptide, provides valuable information about the therapeutic potential of the candidates tested. One further such assay test for induction of cardiac hypertrophy is disclosed in U.S. Pat. No. 5,773,415, using Sprague-Dawley rats.

For cancer, a variety of well-known animal models can be used to further understand the role of the genes identified herein in the development and pathogenesis of tumors, and to test the efficacy of candidate therapeutic agents, including antibodies and other antagonists of native PRO polypeptides, such as small-molecule antagonists. The *in vivo* nature of such models makes them particularly predictive of responses in human patients. Animal models of tumors and cancers (e.g., breast cancer, colon cancer, prostate cancer, lung cancer, etc.) include both non-recombinant and recombinant (transgenic) animals. Non-recombinant animal models include, for example,

rodent, *e.g.*, murine models. Such models can be generated by introducing tumor cells into syngeneic mice using standard techniques, *e.g.*, subcutaneous injection, tail vein injection, spleen implantation, intraperitoneal implantation, implantation under the renal capsule, or orthotopic implantation, *e.g.*, colon cancer cells implanted in colonic tissue. *See, e.g.*, PCT publication No. WO 97/33551, published September 18, 1997. Probably the most often used animal species in oncological studies are immunodeficient mice and, in particular, nude mice. The observation that the nude mouse with thymic hypo/aplasia could successfully act as a host for human tumor xenografts has led to its widespread use for this purpose. The autosomal recessive *nu* gene has been introduced into a very large number of distinct congenic strains of nude mouse, including, for example, ASW, A/He, AKR, BALB/c, B10.LP, C17, C3H, C57BL, C57, CBA, DBA, DDD, I/st, NC, NFR, NFS, NFS/N, NZB, NZC, NZW, P, RIII, and SJL. In addition, a wide variety of other animals with inherited immunological defects other than the nude mouse have been bred and used as recipients of tumor xenografts. For further details *see, e.g.*, The Nude Mouse in Oncology Research, E. Boven and B. Winograd, eds. (CRC Press, Inc., 1991).

The cells introduced into such animals can be derived from known tumor/cancer cell lines, such as any of the above-listed tumor cell lines, and, for example, the B104-1-1 cell line (stable NIH-3T3 cell line transfected with the *neu* protooncogene); *ras*-transfected NIH-3T3 cells; Caco-2 (ATCC HTB-37); or a moderately well-differentiated grade II human colon adenocarcinoma cell line, HT-29 (ATCC HTB-38); or from tumors and cancers. Samples of tumor or cancer cells can be obtained from patients undergoing surgery, using standard conditions involving freezing and storing in liquid nitrogen. Karmali *et al.*, Br. J. Cancer, 48: 689-696 (1983).

Tumor cells can be introduced into animals such as nude mice by a variety of procedures. The subcutaneous (s.c.) space in mice is very suitable for tumor implantation. Tumors can be transplanted s.c. as solid blocks, as needle biopsies by use of a trochar, or as cell suspensions. For solid-block or trochar implantation, tumor tissue fragments of suitable size are introduced into the s.c. space. Cell suspensions are freshly prepared from primary tumors or stable tumor cell lines, and injected subcutaneously. Tumor cells can also be injected as subdermal implants. In this location, the inoculum is deposited between the lower part of the dermal connective tissue and the s.c. tissue.

Animal models of breast cancer can be generated, for example, by implanting rat neuroblastoma cells (from which the *neu* oncogene was initially isolated), or *neu*-transformed NIH-3T3 cells into nude mice, essentially as described by Drebin *et al.* Proc. Nat. Acad. Sci. USA, 83: 9129-9133 (1986).

Similarly, animal models of colon cancer can be generated by passaging colon cancer cells in animals, *e.g.*, nude mice, leading to the appearance of tumors in these animals. An orthotopic transplant model of human colon cancer in nude mice has been described, for example, by Wang *et al.*, Cancer Research, 54: 4726-4728 (1994) and Too *et al.*, Cancer Research, 55: 681-684 (1995). This model is based on the so-called "METAMOUSE™" sold by AntiCancer, Inc., (San Diego, California).

Tumors that arise in animals can be removed and cultured *in vitro*. Cells from the *in vitro* cultures can then be passaged to animals. Such tumors can serve as targets for further testing or drug screening. Alternatively, the tumors resulting from the passage can be isolated and RNA from pre-passage cells and cells isolated after one or more rounds of passage analyzed for differential expression of genes of interest. Such passaging techniques can

be performed with any known tumor or cancer cell lines.

For example, Meth A, CMS4, CMS5, CMS21, and WEHI-164 are chemically induced fibrosarcomas of BALB/c female mice (DeLeo *et al.*, J. Exp. Med., **146**: 720 (1977)), which provide a highly controllable model system for studying the anti-tumor activities of various agents. Palladino *et al.*, J. Immunol., **138**: 4023-4032 (1987). Briefly, tumor cells are propagated *in vitro* in cell culture. Prior to injection into the animals, the cell lines are washed and suspended in buffer, at a cell density of about 10×10^6 to 10×10^7 cells/ml. The animals are then infected subcutaneously with 10 to 100 μ l of the cell suspension, allowing one to three weeks for a tumor to appear.

In addition, the Lewis lung (3LL) carcinoma of mice, which is one of the most thoroughly studied experimental tumors, can be used as an investigational tumor model. Efficacy in this tumor model has been correlated with beneficial effects in the treatment of human patients diagnosed with small-cell carcinoma of the lung (SCCL). This tumor can be introduced in normal mice upon injection of tumor fragments from an affected mouse or of cells maintained in culture. Zupi *et al.*, Br. J. Cancer, **41**: suppl. 4, 30 (1980). Evidence indicates that tumors can be started from injection of even a single cell and that a very high proportion of infected tumor cells survive. For further information about this tumor model *see*, Zacharski, Haemostasis, **16**: 300-320 (1986).

One way of evaluating the efficacy of a test compound in an animal model with an implanted tumor is to measure the size of the tumor before and after treatment. Traditionally, the size of implanted tumors has been measured with a slide caliper in two or three dimensions. The measure limited to two dimensions does not accurately reflect the size of the tumor; therefore, it is usually converted into the corresponding volume by using a mathematical formula. However, the measurement of tumor size is very inaccurate. The therapeutic effects of a drug candidate can be better described as treatment-induced growth delay and specific growth delay. Another important variable in the description of tumor growth is the tumor volume doubling time. Computer programs for the calculation and description of tumor growth are also available, such as the program reported by Rygaard and Spang-Thomsen, Proc. 6th Int. Workshop on Immune-Deficient Animals, Wu and Sheng eds. (Basel, 1989), p. 301. It is noted, however, that necrosis and inflammatory responses following treatment may actually result in an increase in tumor size, at least initially. Therefore, these changes need to be carefully monitored, by a combination of a morphometric method and flow cytometric analysis.

Further, recombinant (transgenic) animal models can be engineered by introducing the coding portion of the PRO gene identified herein into the genome of animals of interest, using standard techniques for producing transgenic animals. Animals that can serve as a target for transgenic manipulation include, without limitation, mice, rats, rabbits, guinea pigs, sheep, goats, pigs, and non-human primates, *e.g.*, baboons, chimpanzees and monkeys. Techniques known in the art to introduce a transgene into such animals include pronucleic microinjection (U.S. Patent No. 4,873,191); retrovirus-mediated gene transfer into germ lines (*e.g.*, Van der Putten *et al.*, Proc. Natl. Acad. Sci. USA, **82**: 6148-615 (1985)); gene targeting in embryonic stem cells (Thompson *et al.*, Cell, **56**: 313-321 (1989)); electroporation of embryos (Lo, Mol. Cell. Biol., **3**: 1803-1814 (1983)); and sperm-mediated gene transfer. Lavitrano *et al.*, Cell, **57**: 717-73 (1989). For a review, *see* for example, U.S. Patent No. 4,736,866.

For the purpose of the present invention, transgenic animals include those that carry the transgene only in part of their cells ("mosaic animals"). The transgene can be integrated either as a single transgene, or in

concatamers, *e.g.*, head-to-head or head-to-tail tandems. Selective introduction of a transgene into a particular cell type is also possible by following, for example, the technique of Lasko *et al.*, Proc. Natl. Acad. Sci. USA, **89**: 6232-636 (1992). The expression of the transgene in transgenic animals can be monitored by standard techniques. For example, Southern blot analysis or PCR amplification can be used to verify the integration of the transgene. The level of mRNA expression can then be analyzed using techniques such as *in situ* hybridization, Northern blot analysis, PCR, or immunocytochemistry. The animals are further examined for signs of tumor or cancer development.

Alternatively, "knock-out" animals can be constructed that have a defective or altered gene encoding a PRO polypeptide identified herein, as a result of homologous recombination between the endogenous gene encoding the PRO polypeptide and altered genomic DNA encoding the same polypeptide introduced into an embryonic cell of the animal. For example, cDNA encoding a particular PRO polypeptide can be used to clone genomic DNA encoding that polypeptide in accordance with established techniques. A portion of the genomic DNA encoding a particular PRO polypeptide can be deleted or replaced with another gene, such as a gene encoding a selectable marker that can be used to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector. *See, e.g.*, Thomas and Capecchi, Cell, **51**: 503 (1987) for a description of homologous recombination vectors. The vector is introduced into an embryonic stem cell line (*e.g.*, by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected. *See, e.g.*, Li *et al.*, Cell, **69**: 915 (1992). The selected cells are then injected into a blastocyst of an animal (*e.g.*, a mouse or rat) to form aggregation chimeras. *See, e.g.*, Bradley, in Teratocarcinomas and Embryonic Stem Cells: A Practical Approach, E. J. Robertson, ed. (IRL: Oxford, 1987), pp. 113-152. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a "knock-out" animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knockout animals can be characterized, for instance, by their ability to defend against certain pathological conditions and by their development of pathological conditions due to absence of the PRO polypeptide.

The efficacy of antibodies specifically binding the PRO polypeptides identified herein, and other drug candidates, can be tested also in the treatment of spontaneous animal tumors. A suitable target for such studies is the feline oral squamous cell carcinoma (SCC). Feline oral SCC is a highly invasive, malignant tumor that is the most common oral malignancy of cats, accounting for over 60% of the oral tumors reported in this species. It rarely metastasizes to distant sites, although this low incidence of metastasis may merely be a reflection of the short survival times for cats with this tumor. These tumors are usually not amenable to surgery, primarily because of the anatomy of the feline oral cavity. At present, there is no effective treatment for this tumor. Prior to entry into the study, each cat undergoes complete clinical examination and biopsy, and is scanned by computed tomography (CT). Cats diagnosed with sublingual oral squamous cell tumors are excluded from the study. The tongue can become paralyzed as a result of such tumor, and even if the treatment kills the tumor, the animals may not be able to feed themselves. Each cat is treated repeatedly, over a longer period of time. Photographs of the tumors will be taken daily during the treatment period, and at each subsequent recheck. After treatment, each cat undergoes another CT

scan. CT scans and thoracic radiograms are evaluated every 8 weeks thereafter. The data are evaluated for differences in survival, response, and toxicity as compared to control groups. Positive response may require evidence of tumor regression, preferably with improvement of quality of life and/or increased life span.

In addition, other spontaneous animal tumors, such as fibrosarcoma, adenocarcinoma, lymphoma, chondroma, or leiomyosarcoma of dogs, cats, and baboons can also be tested. Of these, mammary adenocarcinoma in dogs and cats is a preferred model as its appearance and behavior are very similar to those in humans. However, the use of this model is limited by the rare occurrence of this type of tumor in animals.

Other *in vitro* and *in vivo* cardiovascular, endothelial, and angiogenic tests known in the art are also suitable herein.

5.2.4.2. Tissue Distribution

The results of the cardiovascular, endothelial, and angiogenic assays herein can be verified by further studies, such as by determining mRNA expression in various human tissues.

As noted before, gene amplification and/or gene expression in various tissues may be measured by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA (Thomas, Proc. Natl. Acad. Sci. USA, 77:5201-5205 (1980)), dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes.

Gene expression in various tissues, alternatively, may be measured by immunological methods, such as immunohistochemical staining of tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native-sequence PRO polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to PRO DNA and encoding a specific antibody epitope. General techniques for generating antibodies, and special protocols for *in situ* hybridization are provided hereinbelow.

5.2.4.3. Antibody Binding Studies

The results of the cardiovascular, endothelial, and angiogenic study can be further verified by antibody binding studies, in which the ability of anti-PRO antibodies to inhibit the effect of the PRO polypeptides on endothelial cells or other cells used in the cardiovascular, endothelial, and angiogenic assays is tested. Exemplary antibodies include polyclonal, monoclonal, humanized, bispecific, and heteroconjugate antibodies, the preparation of which will be described hereinbelow.

Antibody binding studies may be carried out in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays. Zola, Monoclonal Antibodies: A Manual of Techniques (CRC Press, Inc., 1987), pp.147-158.

Competitive binding assays rely on the ability of a labeled standard to compete with the test sample analyte for binding with a limited amount of antibody. The amount of target protein in the test sample is inversely proportional to the amount of standard that becomes bound to the antibodies. To facilitate determining the amount of standard that becomes bound, the antibodies preferably are insolubilized before or after the competition, so that the standard and analyte that are bound to the antibodies may conveniently be separated from the standard and analyte that remain unbound.

Sandwich assays involve the use of two antibodies, each capable of binding to a different immunogenic portion, or epitope, of the protein to be detected. In a sandwich assay, the test sample analyte is bound by a first antibody that is immobilized on a solid support, and thereafter a second antibody binds to the analyte, thus forming an insoluble three-part complex. *See, e.g.*, U.S. Pat. No. 4,376,110. The second antibody may itself be labeled with a detectable moiety (direct sandwich assays) or may be measured using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich assay). For example, one type of sandwich assay is an ELISA assay, in which case the detectable moiety is an enzyme.

For immunohistochemistry, the tissue sample may be fresh or frozen or may be embedded in paraffin and fixed with a preservative such as formalin, for example.

5.2.4.4. Cell-Based Tumor Assays

Cell-based assays and animal models for cardiovascular, endothelial, and angiogenic disorders, such as tumors, can be used to verify the findings of a cardiovascular, endothelial, and angiogenic assay herein, and further to understand the relationship between the genes identified herein and the development and pathogenesis of undesirable cardiovascular, endothelial, and angiogenic cell growth. The role of gene products identified herein in the development and pathology of undesirable cardiovascular, endothelial, and angiogenic cell growth, *e.g.*, tumor cells, can be tested by using cells or cells lines that have been identified as being stimulated or inhibited by the PRO polypeptide herein. Such cells include, for example, those set forth in the Examples below.

In a different approach, cells of a cell type known to be involved in a particular cardiovascular, endothelial, and angiogenic disorder are transfected with the cDNAs herein, and the ability of these cDNAs to induce excessive growth or inhibit growth is analyzed. If the cardiovascular, endothelial, and angiogenic disorder is cancer, suitable tumor cells include, for example, stable tumor cell lines such as the B104-1-1 cell line (stable NIH-3T3 cell line transfected with the *neu* protooncogene) and *ras*-transfected NIH-3T3 cells, which can be transfected with the desired gene and monitored for tumorigenic growth. Such transfected cell lines can then be used to test the ability of poly- or monoclonal antibodies or antibody compositions to inhibit tumorigenic cell growth by exerting cytostatic or cytotoxic activity on the growth of the transformed cells, or by mediating antibody-dependent cellular cytotoxicity (ADCC). Cells transfected with the coding sequences of the genes identified herein can further be used to identify drug candidates for the treatment of cardiovascular, endothelial, and angiogenic disorders such as cancer.

In addition, primary cultures derived from tumors in transgenic animals (as described above) can be used in the cell-based assays herein, although stable cell lines are preferred. Techniques to derive continuous cell lines from transgenic animals are well known in the art. *See, e.g.*, Small *et al.*, Mol. Cell. Biol., 5: 642-648 (1985).

5.2.4.5. Gene Therapy

Described below are methods and compositions whereby disease symptoms may be ameliorated. Certain diseases are brought about, at least in part, by an excessive level of gene product, or by the presence of a gene product exhibiting an abnormal or excessive activity. As such, the reduction in the level and/or activity of such gene products would bring about the amelioration of such disease symptoms.

Alternatively, certain other diseases are brought about, at least in part, by the absence or reduction of the level of gene expression, or a reduction in the level of a gene product's activity. As such, an increase in the level of gene expression and/or the activity of such gene products would bring about the amelioration of such disease symptoms.

In some cases, the up-regulation of a gene in a disease state reflects a protective role for that gene product in responding to the disease condition. Enhancement of such a target gene's expression, or the activity of the target gene product, will reinforce the protective effect it exerts. Some disease states may result from an abnormally low level of activity of such a protective gene. In these cases also, an increase in the level of gene expression and/or the activity of such gene products would bring about the amelioration of such disease symptoms.

The PRO polypeptides described herein and polypeptidyl agonists and antagonists may be employed in accordance with the present invention by expression of such polypeptides *in vivo*, which is often referred to as gene therapy.

There are two major approaches to getting the nucleic acid (optionally contained in a vector) into the patient's cells: *in vivo* and *ex vivo*. For *in vivo* delivery the nucleic acid is injected directly into the patient, usually at the sites where the PRO polypeptide is required, *i.e.*, the site of synthesis of the PRO polypeptide, if known, and the site (*e.g.*, wound) where biological activity of the PRO polypeptide is needed. For *ex vivo* treatment, the patient's cells are removed, the nucleic acid is introduced into these isolated cells, and the modified cells are administered to the patient either directly or, for example, encapsulated within porous membranes that are implanted into the patient (*see, e.g.*, U.S. Pat. Nos. 4,892,538 and 5,283,187). There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells *in vitro*, or transferred *in vivo* in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells *in vitro* include the use of liposomes, electroporation, microinjection, transduction, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc. Transduction involves the association of a replication-defective, recombinant viral (preferably retroviral) particle with a cellular receptor, followed by introduction of the nucleic acids contained by the particle into the cell. A commonly used vector for *ex vivo* delivery of the gene is a retrovirus.

The currently preferred *in vivo* nucleic acid transfer techniques include transfection with viral or non-viral vectors (such as adenovirus, lentivirus, Herpes simplex I virus, or adeno-associated virus (AAV)) and lipid-based systems (useful lipids for lipid-mediated transfer of the gene are, for example, DOTMA, DOPE, and DC-Chol; *see, e.g.*, Tonkinson *et al.*, Cancer Investigation, 14(1): 54-65 (1996)). The most preferred vectors for use in gene therapy are viruses, most preferably adenoviruses, AAV, lentiviruses, or retroviruses. A viral vector such as a retroviral vector includes at least one transcriptional promoter/enhancer or locus-defining element(s), or other

elements that control gene expression by other means such as alternate splicing, nuclear RNA export, or post-translational modification of messenger. In addition, a viral vector such as a retroviral vector includes a nucleic acid molecule that, when transcribed in the presence of a gene encoding the PRO polypeptide, is operably linked thereto and acts as a translation initiation sequence. Such vector constructs also include a packaging signal, long terminal repeats (LTRs) or portions thereof, and positive and negative strand primer binding sites appropriate to the virus used (if these are not already present in the viral vector). In addition, such vector typically includes a signal sequence for secretion of the PRO polypeptide from a host cell in which it is placed. Preferably the signal sequence for this purpose is a mammalian signal sequence, most preferably the native signal sequence for the PRO polypeptide. Optionally, the vector construct may also include a signal that directs polyadenylation, as well as one or more restriction sites and a translation termination sequence. By way of example, such vectors will typically include a 5' LTR, a tRNA binding site, a packaging signal, an origin of second-strand DNA synthesis, and a 3' LTR or a portion thereof. Other vectors can be used that are non-viral, such as cationic lipids, polylysine, and dendrimers.

In some situations, it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell-surface membrane protein or the target cell, a ligand for a receptor on the target cell, etc. Where liposomes are employed, proteins that bind to a cell-surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake, *e.g.*, capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins that undergo internalization in cycling, and proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu *et al.*, J. Biol. Chem., **262**: 4429-4432 (1987); and Wagner *et al.*, Proc. Natl. Acad. Sci. USA, **87**: 3410-3414 (1990). For a review of the currently known gene marking and gene therapy protocols, see, Anderson *et al.*, Science, **256**: 808-813 (1992). See also WO 93/25673 and the references cited therein.

Suitable gene therapy and methods for making retroviral particles and structural proteins can be found in, *e.g.*, U.S. Pat. No. 5,681,746.

5.2.4.6. Use of Gene as a Diagnostic

This invention is also related to the use of the gene encoding the PRO polypeptide as a diagnostic. Detection of a mutated form of the PRO polypeptide will allow a diagnosis of a cardiovascular, endothelial, and angiogenic disease or a susceptibility to a cardiovascular, endothelial, and angiogenic disease, such as a tumor, since mutations in the PRO polypeptide may cause tumors.

Individuals carrying mutations in the genes encoding a human PRO polypeptide may be detected at the DNA level by a variety of techniques. Nucleic acids for diagnosis may be obtained from a patient's cells, such as from blood, urine, saliva, tissue biopsy, and autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR (Saiki *et al.*, Nature, **324**: 163-166 (1986)) prior to analysis. RNA or cDNA may also be used for the same purpose. As an example, PCR primers complementary to the nucleic acid encoding the PRO polypeptide can be used to identify and analyze the PRO polypeptide mutations. For example, deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal

genotype. Point mutations can be identified by hybridizing amplified DNA to radiolabeled RNA encoding the PRO polypeptide, or alternatively, radiolabeled antisense DNA sequences encoding the PRO polypeptide. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase A digestion or by differences in melting temperatures.

5 Genetic testing based on DNA sequence differences may be achieved by detection of alteration in electrophoretic mobility of DNA fragments in gels with or without denaturing agents. Small sequence deletions and insertions can be visualized by high resolution gel electrophoresis. DNA fragments of different sequences may be distinguished on denaturing formamide gradient gels in which the mobilities of different DNA fragments are retarded in the gel at different positions according to their specific melting or partial melting temperatures. See, 10 *e.g.*, Myers *et al.*, Science, 230: 1242 (1985).

Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method, for example, Cotton *et al.*, Proc. Natl. Acad. Sci. USA, 85: 4397-4401 (1985).

15 Thus, the detection of a specific DNA sequence may be achieved by methods such as hybridization, RNase protection, chemical cleavage, direct DNA sequencing, or the use of restriction enzymes, *e.g.*, restriction fragment length polymorphisms (RFLP), and Southern blotting of genomic DNA.

5.2.4.7. Use to Detect PRO Polypeptide Levels

In addition to more conventional gel-electrophoresis and DNA sequencing, mutations can also be detected by *in situ* analysis.

20 Expression of nucleic acid encoding the PRO polypeptide may be linked to vascular disease or neovascularization associated with tumor formation. If the PRO polypeptide has a signal sequence and the mRNA is highly expressed in endothelial cells and to a lesser extent in smooth muscle cells, this indicates that the PRO polypeptide is present in serum. Accordingly, an anti-PRO polypeptide antibody could be used to diagnose vascular disease or neovascularization associated with tumor formation, since an altered level of this PRO polypeptide may 25 be indicative of such disorders.

A competition assay may be employed wherein antibodies specific to the PRO polypeptide are attached to a solid support and the labeled PRO polypeptide and a sample derived from the host are passed over the solid support and the amount of label detected attached to the solid support can be correlated to a quantity of the PRO polypeptide in the sample.

30 5.2.4.8. Chromosome Mapping

The sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. Moreover, there is a current need for identifying particular sites on the chromosome. Few chromosome marking reagents based on actual sequence data (repeat polymorphisms) are presently available for marking chromosomal 35 location. The mapping of DNAs to chromosomes according to the present invention is an important first step in

correlating those sequences with genes associated with disease.

Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the cDNA. Computer analysis for the 3'- untranslated region is used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the primer will yield an amplified fragment.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular DNA to a particular chromosome. Using the present invention with the same oligonucleotide primers, sublocalization can be achieved with panels of fragments from specific chromosomes or pools of large genomic clones in an analogous manner. Other mapping strategies that can similarly be used to map to its chromosome include *in situ* hybridization, prescreening with labeled flow-sorted chromosomes, and preselection by hybridization to construct chromosome-specific cDNA libraries.

Fluorescence *in situ* hybridization (FISH) of a cDNA clone to a metaphase chromosomal spread can be used to provide a precise chromosomal location in one step. This technique can be used with cDNA as short as 500 or 600 bases; however, clones larger than 2,000 bp have a higher likelihood of binding to a unique chromosomal location with sufficient signal intensity for simple detection. FISH requires use of the clones from which the gene encoding the PRO polypeptide was derived, and the longer the better. For example, 2,000 bp is good, 4,000 bp is better, and more than 4,000 is probably not necessary to get good results a reasonable percentage of the time. For a review of this technique, see, Verma *et al.*, Human Chromosomes: a Manual of Basic Techniques (Pergamon Press, New York, 1988).

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available online through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region is then identified through linkage analysis (coinheritance of physically adjacent genes).

Next, it is necessary to determine the differences in the cDNA or genomic sequence between affected and unaffected individuals. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

With current resolution of physical mapping and genetic mapping techniques, a cDNA precisely localized to a chromosomal region associated with the disease could be one of between 50 and 500 potential causative genes. (This assumes 1 megabase mapping resolution and one gene per 20 kb).

5.2.4.9. Screening Assays for Drug Candidates

This invention encompasses methods of screening compounds to identify those that mimic the PRO polypeptide (agonists) or prevent the effect of the PRO polypeptide (antagonists). Screening assays for antagonist drug candidates are designed to identify compounds that bind or complex with the PRO polypeptide encoded by the genes identified herein, or otherwise interfere with the interaction of the encoded polypeptides with other

cellular proteins. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates.

The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays, and cell-based assays, which are well characterized in the art.

5 All assays for antagonists are common in that they call for contacting the drug candidate with a PRO polypeptide encoded by a nucleic acid identified herein under conditions and for a time sufficient to allow these two components to interact.

10 In binding assays, the interaction is binding and the complex formed can be isolated or detected in the reaction mixture. In a particular embodiment, the PRO polypeptide encoded by the gene identified herein or the drug candidate is immobilized on a solid phase, *e.g.*, on a microtiter plate, by covalent or non-covalent attachments. Non-covalent attachment generally is accomplished by coating the solid surface with a solution of the PRO polypeptide and drying. Alternatively, an immobilized antibody, *e.g.*, a monoclonal antibody, specific for the PRO polypeptide to be immobilized can be used to anchor it to a solid surface. The assay is performed by adding the non-immobilized component, which may be labeled by a detectable label, to the immobilized component, *e.g.*, the coated surface containing the anchored component. When the reaction is complete, the non-reacted components are removed, *e.g.*, by washing, and complexes anchored on the solid surface are detected. When the originally non-immobilized component carries a detectable label, the detection of label immobilized on the surface indicates that complexing occurred. Where the originally non-immobilized component does not carry a label, complexing can be detected, for example, by using a labeled antibody specifically binding the immobilized complex.

20 If the candidate compound interacts with but does not bind to a particular PRO polypeptide encoded by a gene identified herein, its interaction with that polypeptide can be assayed by methods well known for detecting protein-protein interactions. Such assays include traditional approaches, such as, *e.g.*, cross-linking, co-immunoprecipitation, and co-purification through gradients or chromatographic columns. In addition, protein-protein interactions can be monitored by using a yeast-based genetic system described by Fields and co-workers (Fields and Song, Nature (London), 340: 245-246 (1989); Chien *et al.*, Proc. Natl. Acad. Sci. USA, 88: 9578-9582 (1991)) as disclosed by Chevray and Nathans, Proc. Natl. Acad. Sci. USA, 89: 5789-5793 (1991). Many transcriptional activators, such as yeast GAL4, consist of two physically discrete modular domains, one acting as the DNA-binding domain, the other one functioning as the transcription-activation domain. The yeast expression system described in the foregoing publications (generally referred to as the "two-hybrid system") takes advantage of this property, and employs two hybrid proteins, one in which the target protein is fused to the DNA-binding domain of GAL4, and another, in which candidate activating proteins are fused to the activation domain. The expression of a GAL1-*lacZ* reporter gene under control of a GAL4-activated promoter depends on reconstitution of GAL4 activity via protein-protein interaction. Colonies containing interacting polypeptides are detected with a chromogenic substrate for β -galactosidase. A complete kit (MATCHMAKER™) for identifying protein-protein interactions between two specific proteins using the two-hybrid technique is commercially available from Clontech. This system can also be extended to map protein domains involved in specific protein interactions as well as to pinpoint amino acid residues that are crucial for these interactions.

Compounds that interfere with the interaction of a gene encoding a PRO polypeptide identified herein and other intra- or extracellular components can be tested as follows: usually a reaction mixture is prepared containing the product of the gene and the intra- or extracellular component under conditions and for a time allowing for the interaction and binding of the two products. To test the ability of a candidate compound to inhibit binding, the reaction is run in the absence and in the presence of the test compound. In addition, a placebo may be added to a third reaction mixture, to serve as positive control. The binding (complex formation) between the test compound and the intra- or extracellular component present in the mixture is monitored as described hereinabove. The formation of a complex in the control reaction(s) but not in the reaction mixture containing the test compound indicates that the test compound interferes with the interaction of the test compound and its reaction partner.

If the PRO polypeptide has the ability to stimulate the proliferation of endothelial cells in the presence of the co-mitogen ConA, then one example of a screening method takes advantage of this ability. Specifically, in the proliferation assay, human umbilical vein endothelial cells are obtained and cultured in 96-well flat-bottomed culture plates (Costar, Cambridge, MA) and supplemented with a reaction mixture appropriate for facilitating proliferation of the cells, the mixture containing Con-A (Calbiochem, La Jolla, CA). Con-A and the compound to be screened are added and after incubation at 37°C, cultures are pulsed with ³H-thymidine and harvested onto glass fiber filters (pH; Cambridge Technology, Watertown, MA). Mean ³H-thymidine incorporation (cpm) of triplicate cultures is determined using a liquid scintillation counter (Beckman Instruments, Irvine, CA). Significant ³(H)-thymidine incorporation indicates stimulation of endothelial cell proliferation.

To assay for antagonists, the assay described above is performed; however, in this assay the PRO polypeptide is added along with the compound to be screened and the ability of the compound to inhibit ³(H)thymidine incorporation in the presence of the PRO polypeptide indicates that the compound is an antagonist to the PRO polypeptide. Alternatively, antagonists may be detected by combining the PRO polypeptide and a potential antagonist with membrane-bound PRO polypeptide receptors or recombinant receptors under appropriate conditions for a competitive inhibition assay. The PRO polypeptide can be labeled, such as by radioactivity, such that the number of PRO polypeptide molecules bound to the receptor can be used to determine the effectiveness of the potential antagonist. The gene encoding the receptor can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting. Coligan *et al.*, Current Protocols in Immun., 1(2): Chapter 5 (1991). Preferably, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the PRO polypeptide and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the PRO polypeptide. Transfected cells that are grown on glass slides are exposed to the labeled PRO polypeptide. The PRO polypeptide can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase. Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an interactive sub-pooling and re-screening process, eventually yielding a single clone that encodes the putative receptor.

As an alternative approach for receptor identification, the labeled PRO polypeptide can be photoaffinity-linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is

resolved by PAGE and exposed to X-ray film. The labeled complex containing the receptor can be excised, resolved into peptide fragments, and subjected to protein micro-sequencing. The amino acid sequence obtained from micro-sequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the gene encoding the putative receptor.

5 In another assay for antagonists, mammalian cells or a membrane preparation expressing the receptor would be incubated with the labeled PRO polypeptide in the presence of the candidate compound. The ability of the compound to enhance or block this interaction could then be measured.

10 The compositions useful in the treatment of cardiovascular, endothelial, and angiogenic disorders include, without limitation, antibodies, small organic and inorganic molecules, peptides, phosphopeptides, antisense and ribozyme molecules, triple-helix molecules, etc., that inhibit the expression and/or activity of the target gene product.

15 More specific examples of potential antagonists include an oligonucleotide that binds to the fusions of immunoglobulin with a PRO polypeptide, and, in particular, antibodies including, without limitation, poly- and monoclonal antibodies and antibody fragments, single-chain antibodies, anti-idiotypic antibodies, and chimeric or humanized versions of such antibodies or fragments, as well as human antibodies and antibody fragments. Alternatively, a potential antagonist may be a closely related protein, for example, a mutated form of the PRO polypeptide that recognizes the receptor but imparts no effect, thereby competitively inhibiting the action of the PRO polypeptide.

20 Another potential PRO polypeptide antagonist is an antisense RNA or DNA construct prepared using antisense technology, where, *e.g.*, an antisense RNA or DNA molecule acts to block directly the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes the mature PRO polypeptides herein, is used to design an antisense RNA oligonucleotide
25 of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix - *see*, Lee *et al.*, Nucl. Acids Res., **6**:3073 (1979); Cooney *et al.*, Science, **241**: 456 (1988); Dervan *et al.*, Science, **251**:1360 (1991)), thereby preventing transcription and the production of the PRO polypeptide. A sequence "complementary" to a portion of an RNA, as referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex;
30 in the case of double-stranded antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex helix formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the longer the hybridizing nucleic acid, the more base mismatches with an RNA it may contain and still form a stable duplex (or triplex, as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point
35 of the hybridized complex. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into the PRO polypeptide (antisense - Okano, Neurochem., **56**:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression (CRC Press: Boca Raton, FL, 1988).

The antisense oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger, *et al.*, 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556; Lemaitre, *et al.*, 1987, *Proc. Natl. Acad. Sci. U.S.A.* 84:648-652; PCT Publication No. WO88/09810, published December 15, 1988) or the blood-brain barrier (see, *e.g.*, PCT Publication No. WO89/10134, published April 25, 1988), hybridization-triggered cleavage agents (see, *e.g.*, Krol *et al.*, 1988, *BioTechniques* 6:958-976) or intercalating agents (see, *e.g.*, Zon, 1988, *Pharm. Res.* 5:539-549). To this end, the oligonucleotide may be conjugated to another molecule, *e.g.*, a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group consisting of a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gautier, *et al.*, 1987, *Nucl. Acids Res.* 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue, *et al.*, 1987, *Nucl. Acids Res.* 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue, *et al.*, 1987, *FEBS Lett.* 215:327-330).

Oligonucleotides of the invention may be synthesized by standard methods known in the art, *e.g.*, by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein, *et al.* (1988, *Nucl. Acids Res.* 16:3209), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin, *et al.*, 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85:7448-7451), etc.

The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed *in vivo* to inhibit production of the PRO polypeptide. When antisense DNA is used, oligodeoxyribonucleotides derived from the translation-initiation site, *e.g.*, between about -10 and +10 positions of the target gene nucleotide sequence, are preferred.

5 Antisense RNA or DNA molecules are generally at least about 5 bases in length, about 10 bases in length, about 15 bases in length, about 20 bases in length, about 25 bases in length, about 30 bases in length, about 35 bases in length, about 40 bases in length, about 45 bases in length, about 50 bases in length, about 55 bases in length, about 60 bases in length, about 65 bases in length, about 70 bases in length, about 75 bases in length, about 80 bases in length, about 85 bases in length, about 90 bases in length, about 95 bases in length, about 100 bases in length,
10 or more.

Potential antagonists further include small molecules that bind to the active site, the receptor binding site, or growth factor or other relevant binding site of the PRO polypeptide, thereby blocking the normal biological activity of the PRO polypeptide. Examples of small molecules include, but are not limited to, small peptides or peptide-like molecules, preferably soluble peptides, and synthetic non-peptidyl organic or inorganic compounds.

15 Additional potential antagonists are ribozymes, which are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. Ribozymes act by sequence-specific hybridization to the complementary target RNA, followed by endonucleolytic cleavage. Specific ribozyme cleavage sites within a potential RNA target can be identified by known techniques. For further details *see, e.g.*, Rossi, Current Biology, 4: 469-471 (1994), and PCT publication No. WO 97/33551 (published September 18, 1997).

20 While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy target gene mRNAs, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions which form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Myers, 1995, *Molecular Biology and Biotechnology: A Comprehensive Desk Reference*, VCH Publishers, New York, (see especially Figure 4, page
25 833) and in Haseloff and Gerlach, 1988, *Nature*, 334:585-591, which is incorporated herein by reference in its entirety.

30 Preferably the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the target gene mRNA, *i.e.*, to increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

The ribozymes of the present invention also include RNA endoribonucleases (hereinafter "Cech-type ribozymes") such as the one which occurs naturally in *Tetrahymena thermophila* (known as the IVS, or L-19 IVS RNA) and which has been extensively described by Thomas Cech and collaborators (Zaug, *et al.*, 1984, *Science*, 224:574-578; Zaug and Cech, 1986, *Science*, 231:470-475; Zaug, *et al.*, 1986, *Nature*, 324:429-433; published
35 International patent application No. WO 88/04300 by University Patents Inc.; Been and Cech, 1986, *Cell*, 47:207-216). The Cech-type ribozymes have an eight base pair active site that hybridizes to a target RNA sequence

whereafter cleavage of the target RNA takes place. The invention encompasses those Cech-type ribozymes that target eight base-pair active site sequences that are present in the target gene.

As in the antisense approach, the ribozymes can be composed of modified oligonucleotides (*e.g.*, for improved stability, targeting, *etc.*) and should be delivered to cells that express the target gene *in vivo*. A preferred method of delivery involves using a DNA construct "encoding" the ribozyme under the control of a strong constitutive pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous target gene messages and inhibit translation. Because ribozymes, unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.

Nucleic acid molecules in triple-helix formation used to inhibit transcription should be single-stranded and composed of deoxynucleotides. The base composition of these oligonucleotides is designed such that it promotes triple-helix formation via Hoogsteen base-pairing rules, which generally require sizeable stretches of purines or pyrimidines on one strand of a duplex. For further details *see, e.g.*, PCT publication No. WO 97/33551, *supra*.

These small molecules can be identified by any one or more of the screening assays discussed hereinabove and/or by any other screening techniques well known for those skilled in the art.

5.2.4.10. Types of Cardiovascular, Endothelial, and Angiogenic Disorders to be Treated

The PRO polypeptides, or agonists or antagonists thereto, that have activity in the cardiovascular, angiogenic, and endothelial assays described herein, and/or whose gene product has been found to be localized to the cardiovascular system, are likely to have therapeutic uses in a variety of cardiovascular, endothelial, and angiogenic disorders, including systemic disorders that affect vessels, such as diabetes mellitus. Their therapeutic utility could include diseases of the arteries, capillaries, veins, and/or lymphatics. Examples of treatments hereunder include treating muscle wasting disease, treating osteoporosis, aiding in implant fixation to stimulate the growth of cells around the implant and therefore facilitate its attachment to its intended site, increasing IGF stability in tissues or in serum, if applicable, and increasing binding to the IGF receptor (since IGF has been shown *in vitro* to enhance human marrow erythroid and granulocytic progenitor cell growth).

The PRO polypeptides or agonists or antagonists thereto may also be employed to stimulate erythropoiesis or granulopoiesis, to stimulate wound healing or tissue regeneration and associated therapies concerned with re-growth of tissue, such as connective tissue, skin, bone, cartilage, muscle, lung, or kidney, to promote angiogenesis, to stimulate or inhibit migration of endothelial cells, and to proliferate the growth of vascular smooth muscle and endothelial cell production. The increase in angiogenesis mediated by the PRO polypeptide or agonist would be beneficial to ischemic tissues and to collateral coronary development in the heart subsequent to coronary stenosis. Antagonists are used to inhibit the action of such polypeptides, for example, to limit the production of excess connective tissue during wound healing or pulmonary fibrosis if the PRO polypeptide promotes such production. This would include treatment of acute myocardial infarction and heart failure.

Moreover, the present invention provides the treatment of cardiac hypertrophy, regardless of the underlying cause, by administering a therapeutically effective dose of the PRO polypeptide, or agonist or antagonist thereto. If the objective is the treatment of human patients, the PRO polypeptide preferably is recombinant human PRO

polypeptide (rhPRO polypeptide). The treatment for cardiac hypertrophy can be performed at any of its various stages, which may result from a variety of diverse pathologic conditions, including myocardial infarction, hypertension, hypertrophic cardiomyopathy, and valvular regurgitation. The treatment extends to all stages of the progression of cardiac hypertrophy, with or without structural damage of the heart muscle, regardless of the underlying cardiac disorder.

The decision of whether to use the molecule itself or an agonist thereof for any particular indication, as opposed to an antagonist to the molecule, would depend mainly on whether the molecule herein promotes cardiovascularization, genesis of endothelial cells, or angiogenesis or inhibits these conditions. For example, if the molecule promotes angiogenesis, an antagonist thereof would be useful for treatment of disorders where it is desired to limit or prevent angiogenesis. Examples of such disorders include vascular tumors such as haemangioma, tumor angiogenesis, neovascularization in the retina, choroid, or cornea, associated with diabetic retinopathy or premature infant retinopathy or macular degeneration and proliferative vitreoretinopathy, rheumatoid arthritis, Crohn's disease, atherosclerosis, ovarian hyperstimulation, psoriasis, endometriosis associated with neovascularization, restenosis subsequent to balloon angioplasty, scar tissue overproduction, for example, that seen in a keloid that forms after surgery, fibrosis after myocardial infarction, or fibrotic lesions associated with pulmonary fibrosis.

If, however, the molecule inhibits angiogenesis, it would be expected to be used directly for treatment of the above conditions.

On the other hand, if the molecule stimulates angiogenesis it would be used itself (or an agonist thereof) for indications where angiogenesis is desired such as peripheral vascular disease, hypertension, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, arterial restenosis, thrombophlebitis, lymphangitis, lymphedema, wound healing and tissue repair, ischemia reperfusion injury, angina, myocardial infarctions such as acute myocardial infarctions, chronic heart conditions, heart failure such as congestive heart failure, and osteoporosis.

If, however, the molecule inhibits angiogenesis, an antagonist thereof would be used for treatment of those conditions where angiogenesis is desired.

Specific types of diseases are described below, where the PRO polypeptide herein or agonists or antagonists thereof may serve as useful for vascular-related drug targeting or as therapeutic targets for the treatment or prevention of the disorders. Atherosclerosis is a disease characterized by accumulation of plaques of intimal thickening in arteries, due to accumulation of lipids, proliferation of smooth muscle cells, and formation of fibrous tissue within the arterial wall. The disease can affect large, medium, and small arteries in any organ. Changes in endothelial and vascular smooth muscle cell function are known to play an important role in modulating the accumulation and regression of these plaques.

Hypertension is characterized by raised vascular pressure in the systemic arterial, pulmonary arterial, or portal venous systems. Elevated pressure may result from or result in impaired endothelial function and/or vascular disease.

Inflammatory vasculitides include giant cell arteritis, Takayasu's arteritis, polyarteritis nodosa (including the microangiopathic form), Kawasaki's disease, microscopic polyangiitis, Wegener's granulomatosis, and a variety

of infectious-related vascular disorders (including Henoch-Schonlein purpura). Altered endothelial cell function has been shown to be important in these diseases.

Reynaud's disease and Reynaud's phenomenon are characterized by intermittent abnormal impairment of the circulation through the extremities on exposure to cold. Altered endothelial cell function has been shown to be important in this disease.

Aneurysms are saccular or fusiform dilatations of the arterial or venous tree that are associated with altered endothelial cell and/or vascular smooth muscle cells.

Arterial restenosis (restenosis of the arterial wall) may occur following angioplasty as a result of alteration in the function and proliferation of endothelial and vascular smooth muscle cells.

Thrombophlebitis and lymphangitis are inflammatory disorders of veins and lymphatics, respectively, that may result from, and/or in, altered endothelial cell function. Similarly, lymphedema is a condition involving impaired lymphatic vessels resulting from endothelial cell function.

The family of benign and malignant vascular tumors are characterized by abnormal proliferation and growth of cellular elements of the vascular system. For example, lymphangiomas are benign tumors of the lymphatic system that are congenital, often cystic, malformations of the lymphatics that usually occur in newborns. Cystic tumors tend to grow into the adjacent tissue. Cystic tumors usually occur in the cervical and axillary region. They can also occur in the soft tissue of the extremities. The main symptoms are dilated, sometimes reticular, structured lymphatics and lymphocysts surrounded by connective tissue. Lymphangiomas are assumed to be caused by improperly connected embryonic lymphatics or their deficiency. The result is impaired local lymph drainage. Griener *et al.*, *Lymphology*, 4: 140-144 (1971).

Another use for the PRO polypeptides herein or agonists or antagonists thereto is in the prevention of tumor angiogenesis, which involves vascularization of a tumor to enable it to grow and/or metastasize. This process is dependent on the growth of new blood vessels. Examples of neoplasms and related conditions that involve tumor angiogenesis include breast carcinomas, lung carcinomas, gastric carcinomas, esophageal carcinomas, colorectal carcinomas, liver carcinomas, ovarian carcinomas, thecomas, arrhenoblastomas, cervical carcinomas, endometrial carcinoma, endometrial hyperplasia, endometriosis, fibrosarcomas, choriocarcinoma, head and neck cancer, nasopharyngeal carcinoma, laryngeal carcinomas, hepatoblastoma, Kaposi's sarcoma, melanoma, skin carcinomas, hemangioma, cavernous hemangioma, hemangioblastoma, pancreas carcinomas, retinoblastoma, astrocytoma, glioblastoma, Schwannoma, oligodendroglioma, medulloblastoma, neuroblastomas, rhabdomyosarcoma, osteogenic sarcoma, leiomyosarcomas, urinary tract carcinomas, thyroid carcinomas, Wilm's tumor, renal cell carcinoma, prostate carcinoma, abnormal vascular proliferation associated with phakomatoses, edema (such as that associated with brain tumors), and Meigs' syndrome.

Age-related macular degeneration (AMD) is a leading cause of severe visual loss in the elderly population. The exudative form of AMD is characterized by choroidal neovascularization and retinal pigment epithelial cell detachment. Because choroidal neovascularization is associated with a dramatic worsening in prognosis, the PRO polypeptide or agonist or antagonist thereto is expected to be useful in reducing the severity of AMD.

Healing of trauma such as wound healing and tissue repair is also a targeted use for the PRO polypeptides herein or their agonists or antagonists. Formation and regression of new blood vessels is essential for tissue healing and repair. This category includes bone, cartilage, tendon, ligament, and/or nerve tissue growth or regeneration, as well as wound healing and tissue repair and replacement, and in the treatment of burns, incisions, and ulcers.

5 A PRO polypeptide or agonist or antagonist thereof that induces cartilage and/or bone growth in circumstances where bone is not normally formed has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a PRO polypeptide or agonist or antagonist thereof may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. *De novo* bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma-

10 induced, or oncologic, resection-induced craniofacial defects, and also is useful in cosmetic plastic surgery.

PRO polypeptides or agonists or antagonists thereto may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

15 It is expected that a PRO polypeptide or agonist or antagonist thereto may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, or endothelium), muscle (smooth, skeletal, or cardiac), and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring to allow normal tissue to regenerate.

20 A PRO polypeptide herein or agonist or antagonist thereto may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage. Also, the PRO polypeptide or agonist or antagonist thereto may be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells, or for inhibiting the growth of tissues described above.

25 A PRO polypeptide or agonist or antagonist thereto may also be used in the treatment of periodontal diseases and in other tooth-repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells, or induce differentiation of progenitors of bone-forming cells. A PRO polypeptide herein or an agonist or an antagonist thereto may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes, since blood

30 vessels play an important role in the regulation of bone turnover and growth.

Another category of tissue regeneration activity that may be attributable to the PRO polypeptide herein or agonist or antagonist thereto is tendon/ligament formation. A protein that induces tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed has application in the healing of tendon or ligament tears, deformities, and other tendon or ligament defects in humans and other animals. Such

35 a preparation may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. *De novo* tendon/ligament-like tissue formation induced by a composition of the PRO polypeptide herein or

agonist or antagonist thereto contributes to the repair of congenital, trauma-induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions herein may provide an environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions herein may also be useful in the treatment of tendinitis, carpal tunnel syndrome, and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The PRO polypeptide or its agonist or antagonist may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, *i.e.*, for the treatment of central and peripheral nervous system disease and neuropathies, as well as mechanical and traumatic disorders, that involve degeneration, death, or trauma to neural cells or nerve tissue. More specifically, a PRO polypeptide or its agonist or antagonist may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions that may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head trauma, and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a PRO polypeptide herein or agonist or antagonist thereto.

Ischemia-reperfusion injury is another indication. Endothelial cell dysfunction may be important in both the initiation of, and in regulation of the sequelae of events that occur following ischemia-reperfusion injury.

Rheumatoid arthritis is a further indication. Blood vessel growth and targeting of inflammatory cells through the vasculature is an important component in the pathogenesis of rheumatoid and sero-negative forms of arthritis.

A PRO polypeptide or its agonist or antagonist may also be administered prophylactically to patients with cardiac hypertrophy, to prevent the progression of the condition, and avoid sudden death, including death of asymptomatic patients. Such preventative therapy is particularly warranted in the case of patients diagnosed with massive left ventricular cardiac hypertrophy (a maximal wall thickness of 35 mm or more in adults, or a comparable value in children), or in instances when the hemodynamic burden on the heart is particularly strong.

A PRO polypeptide or its agonist or antagonist may also be useful in the management of atrial fibrillation, which develops in a substantial portion of patients diagnosed with hypertrophic cardiomyopathy.

Further indications include angina, myocardial infarctions such as acute myocardial infarctions, and heart failure such as congestive heart failure. Additional non-neoplastic conditions include psoriasis, diabetic and other proliferative retinopathies including retinopathy of prematurity, retrolental fibroplasia, neovascular glaucoma, thyroid hyperplasias (including Grave's disease), corneal and other tissue transplantation, chronic inflammation, lung inflammation, nephrotic syndrome, preeclampsia, ascites, pericardial effusion (such as that associated with pericarditis), and pleural effusion.

In view of the above, the PRO polypeptides or agonists or antagonists thereof described herein, which are shown to alter or impact endothelial cell function, proliferation, and/or form, are likely to play an important role in the etiology and pathogenesis of many or all of the disorders noted above, and as such can serve as therapeutic targets to augment or inhibit these processes or for vascular-related drug targeting in these disorders.

5.2.4.11. Administration Protocols, Schedules, Doses, and Formulations

The molecules herein and agonists and antagonists thereto are pharmaceutically useful as a prophylactic and therapeutic agent for various disorders and diseases as set forth above.

Therapeutic compositions of the PRO polypeptides or agonists or antagonists are prepared for storage by mixing the desired molecule having the appropriate degree of purity with optional pharmaceutically acceptable carriers, excipients, or stabilizers (Remington's Pharmaceutical Sciences, 16th edition, Osol, A. ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrins; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (*e.g.*, Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG).

Additional examples of such carriers include ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts, or electrolytes such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, and polyethylene glycol. Carriers for topical or gel-based forms of agonist or antagonist include polysaccharides such as sodium carboxymethylcellulose or methylcellulose, polyvinylpyrrolidone, polyacrylates, polyoxyethylene-polyoxypropylene-block polymers, polyethylene glycol, and wood wax alcohols. For all administrations, conventional depot forms are suitably used. Such forms include, for example, microcapsules, nano-capsules, liposomes, plasters, inhalation forms, nose sprays, sublingual tablets, and sustained-release preparations. The PRO polypeptides or agonists or antagonists will typically be formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml.

Another formulation comprises incorporating a PRO polypeptide or agonist or antagonist thereof into formed articles. Such articles can be used in modulating endothelial cell growth and angiogenesis. In addition, tumor invasion and metastasis may be modulated with these articles.

PRO polypeptides or agonists or antagonists to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution. PRO polypeptides ordinarily will be stored in lyophilized form or in solution if administered systemically. If in lyophilized form, the PRO polypeptide or agonist or antagonist thereto is typically formulated in combination with other ingredients for reconstitution with an appropriate diluent at the time for use. An example of a liquid formulation of a PRO polypeptide or agonist or antagonist is a sterile, clear, colorless unpreserved solution filled in a single-dose vial for subcutaneous injection. Preserved pharmaceutical compositions suitable for repeated use may contain, for example, depending mainly on the indication and type of polypeptide:

- a) PRO polypeptide or agonist or antagonist thereto;
- b) a buffer capable of maintaining the pH in a range of maximum stability of the polypeptide or other molecule in solution, preferably about 4-8;
- c) a detergent/surfactant primarily to stabilize the polypeptide or molecule against agitation-induced aggregation;
- d) an isotonicifier;
- e) a preservative selected from the group of phenol, benzyl alcohol and a benzethonium halide, *e.g.*, chloride; and
- f) water.

If the detergent employed is non-ionic, it may, for example, be polysorbates (*e.g.*, POLYSORBATE™ (TWEEN™) 20, 80, etc.) or poloxamers (*e.g.*, POLOXAMER™ 188). The use of non-ionic surfactants permits the formulation to be exposed to shear surface stresses without causing denaturation of the polypeptide. Further, such surfactant-containing formulations may be employed in aerosol devices such as those used in a pulmonary dosing, and needleless jet injector guns (*see, e.g.*, EP 257,956).

An isotonicifier may be present to ensure isotonicity of a liquid composition of the PRO polypeptide or agonist or antagonist thereto, and includes polyhydric sugar alcohols, preferably trihydric or higher sugar alcohols, such as glycerin, erythritol, arabitol, xylitol, sorbitol, and mannitol. These sugar alcohols can be used alone or in combination. Alternatively, sodium chloride or other appropriate inorganic salts may be used to render the solutions isotonic.

The buffer may, for example, be an acetate, citrate, succinate, or phosphate buffer depending on the pH desired. The pH of one type of liquid formulation of this invention is buffered in the range of about 4 to 8, preferably about physiological pH.

The preservatives phenol, benzyl alcohol and benzethonium halides, *e.g.*, chloride, are known antimicrobial agents that may be employed.

Therapeutic PRO polypeptide compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle. The formulations are preferably administered as repeated intravenous (*i.v.*), subcutaneous (*s.c.*), or intramuscular (*i.m.*) injections, or as aerosol formulations suitable for intranasal or intrapulmonary delivery (for intrapulmonary delivery *see, e.g.*, EP 257,956).

PRO polypeptides can also be administered in the form of sustained-released preparations. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the protein, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (*e.g.*, poly(2-hydroxyethyl-methacrylate) as described by Langer *et al.*, J. Biomed. Mater. Res., **15**: 167-277 (1981) and Langer, Chem. Tech., **12**: 98-105 (1982) or poly(vinylalcohol)), polylactides (U.S. Patent No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman *et al.*, Biopolymers, **22**: 547-556 (1983)), non-degradable ethylene-vinyl acetate (Langer *et al.*, *supra*), degradable lactic acid-glycolic acid copolymers such as the Lupron Depot™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid (EP 133,988).

While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated proteins remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for protein stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

Sustained-release PRO polypeptide compositions also include liposomally entrapped PRO polypeptides. Liposomes containing the PRO polypeptide are prepared by methods known *per se*: DE 3,218,121; Epstein *et al.*, Proc. Natl. Acad. Sci. USA, **82**: 3688-3692 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. USA, **77**: 4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese patent application 83-118008; U.S. Patent Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. % cholesterol, the selected proportion being adjusted for the optimal therapy.

The therapeutically effective dose of a PRO polypeptide or agonist or antagonist thereto will, of course, vary depending on such factors as the pathological condition to be treated (including prevention), the method of administration, the type of compound being used for treatment, any co-therapy involved, the patient's age, weight, general medical condition, medical history, etc., and its determination is well within the skill of a practicing physician. Accordingly, it will be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the maximal therapeutic effect. If the PRO polypeptide has a narrow host range, for the treatment of human patients formulations comprising human PRO polypeptide, more preferably native-sequence human PRO polypeptide, are preferred. The clinician will administer the PRO polypeptide until a dosage is reached that achieves the desired effect for treatment of the condition in question. For example, if the objective is the treatment of CHF, the amount would be one that inhibits the progressive cardiac hypertrophy associated with this condition. The progress of this therapy is easily monitored by echo cardiography. Similarly, in patients with hypertrophic cardiomyopathy, the PRO polypeptide can be administered on an empirical basis.

With the above guidelines, the effective dose generally is within the range of from about 0.001 to about 1.0 mg/kg, more preferably about 0.01-1.0 mg/kg, most preferably about 0.01-0.1 mg/kg.

For non-oral use in treating human adult hypertension, it is advantageous to administer the PRO polypeptide in the form of an injection at about 0.01 to 50 mg, preferably about 0.05 to 20 mg, most preferably 1 to 20 mg, per kg body weight, 1 to 3 times daily by intravenous injection. For oral administration, a molecule based on the PRO polypeptide is preferably administered at about 5 mg to 1 g, preferably about 10 to 100 mg, per kg body weight, 1 to 3 times daily. It should be appreciated that endotoxin contamination should be kept minimally at a safe level, for example, less than 0.5 ng/mg protein. Moreover, for human administration, the formulations preferably meet sterility, pyrogenicity, general safety, and purity as required by FDA Office and Biologics standards.

The dosage regimen of a pharmaceutical composition containing the PRO polypeptide to be used in tissue regeneration will be determined by the attending physician considering various factors that modify the action of the polypeptides, *e.g.*, amount of tissue weight desired to be formed, the site of damage, the condition of the damaged tissue, the size of a wound, type of damaged tissue (*e.g.*, bone), the patient's age, sex, and diet, the severity of any infection, time of administration, and other clinical factors. The dosage may vary with the type of matrix used in the reconstitution and with inclusion of other proteins in the pharmaceutical composition. For example, the addition of other known growth factors, such as IGF-I, to the final composition may also affect the dosage. Progress can be monitored by periodic assessment of tissue/bone growth and/or repair, for example, X-rays, histomorphometric determinations, and tetracycline labeling.

The route of PRO polypeptide or antagonist or agonist administration is in accord with known methods, *e.g.*, by injection or infusion by intravenous, intramuscular, intracerebral, intraperitoneal, intracerebrospinal, subcutaneous, intraocular, intraarticular, intrasynovial, intrathecal, oral, topical, or inhalation routes, or by sustained-release systems as noted below. The PRO polypeptide or agonist or antagonists thereof also are suitably administered by intratumoral, peritumoral, intralesional, or perilesional routes, to exert local as well as systemic therapeutic effects. The intraperitoneal route is expected to be particularly useful, for example, in the treatment of ovarian tumors.

If a peptide or small molecule is employed as an antagonist or agonist, it is preferably administered orally or non-orally in the form of a liquid or solid to mammals.

Examples of pharmacologically acceptable salts of molecules that form salts and are useful hereunder include alkali metal salts (*e.g.*, sodium salt, potassium salt), alkaline earth metal salts (*e.g.*, calcium salt, magnesium salt), ammonium salts, organic base salts (*e.g.*, pyridine salt, triethylamine salt), inorganic acid salts (*e.g.*, hydrochloride, sulfate, nitrate), and salts of organic acid (*e.g.*, acetate, oxalate, p-toluenesulfonate).

For compositions herein that are useful for bone, cartilage, tendon, or ligament regeneration, the therapeutic method includes administering the composition topically, systemically, or locally as an implant or device. When administered, the therapeutic composition for use is in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage, or tissue damage. Topical administration may be suitable for wound healing and tissue repair. Preferably, for bone and/or cartilage formation, the composition would include a matrix capable of delivering the

protein-containing composition to the site of bone and/or cartilage damage, providing a structure for the developing bone and cartilage and preferably capable of being resorbed into the body. Such matrices may be formed of materials presently in use for other implanted medical applications.

The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance, and interface properties. The particular application of the compositions will define the appropriate formulation. Potential matrices for the compositions may be biodegradable and chemically defined calcium sulfate, tricalcium phosphate, hydroxyapatite, polylactic acid, polyglycolic acid, and polyanhydrides. Other potential materials are biodegradable and biologically well-defined, such as bone or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are nonbiodegradable and chemically defined, such as sintered hydroxyapatite, bioglass, aluminates, or other ceramics. Matrices may be comprised of combinations of any of the above-mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalcium phosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size, particle shape, and biodegradability.

One specific embodiment is a 50:50 (mole weight) copolymer of lactic acid and glycolic acid in the form of porous particles having diameters ranging from 150 to 800 microns. In some applications, it will be useful to utilize a sequestering agent, such as carboxymethyl cellulose or autologous blood clot, to prevent the polypeptide compositions from disassociating from the matrix.

One suitable family of sequestering agents is cellulosic materials such as alkylcelluloses (including hydroxyalkylcelluloses), including methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, and carboxymethylcellulose, one preferred being cationic salts of carboxymethylcellulose (CMC). Other preferred sequestering agents include hyaluronic acid, sodium alginate, poly(ethylene glycol), polyoxyethylene oxide, carboxyvinyl polymer, and poly(vinyl alcohol). The amount of sequestering agent useful herein is 0.5-20 wt%, preferably 1-10 wt%, based on total formulation weight, which represents the amount necessary to prevent desorption of the polypeptide (or its antagonist) from the polymer matrix and to provide appropriate handling of the composition, yet not so much that the progenitor cells are prevented from infiltrating the matrix, thereby providing the polypeptide (or its antagonist) the opportunity to assist the osteogenic activity of the progenitor cells.

5.2.4.12. Combination Therapies

The effectiveness of the PRO polypeptide or an agonist or antagonist thereof in preventing or treating the disorder in question may be improved by administering the active agent serially or in combination with another agent that is effective for those purposes, either in the same composition or as separate compositions.

For example, for treatment of cardiac hypertrophy, PRO polypeptide therapy can be combined with the administration of inhibitors of known cardiac myocyte hypertrophy factors, e.g., inhibitors of α -adrenergic agonists such as phenylephrine; endothelin-1 inhibitors such as BOSENTANTM and MOXONODINTM; inhibitors to CT-1

(U.S. Pat. No. 5,679,545); inhibitors to LIF; ACE inhibitors; des-aspartate-angiotensin I inhibitors (U.S. Pat. No. 5,773,415), and angiotensin II inhibitors.

For treatment of cardiac hypertrophy associated with hypertension, the PRO polypeptide can be administered in combination with β -adrenergic receptor blocking agents, *e.g.*, propranolol, timolol, tertalolol, carteolol, nadolol, betaxolol, penbutolol, acetobutolol, atenolol, metoprolol, or carvedilol; ACE inhibitors, *e.g.*, quinapril, captopril, enalapril, ramipril, benazepril, fosinopril, or lisinopril; diuretics, *e.g.*, chlorothiazide, hydrochlorothiazide, hydroflumethazide, methylchlorothiazide, benzthiazide, dichlorphenamide, acetazolamide, or indapamide; and/or calcium channel blockers, *e.g.*, diltiazem, nifedipine, verapamil, or nicardipine. Pharmaceutical compositions comprising the therapeutic agents identified herein by their generic names are commercially available, and are to be administered following the manufacturers' instructions for dosage, administration, adverse effects, contraindications, etc. *See, e.g., Physicians' Desk Reference* (Medical Economics Data Production Co.: Montvale, N.J., 1997), 51th Edition.

Preferred candidates for combination therapy in the treatment of hypertrophic cardiomyopathy are β -adrenergic-blocking drugs (*e.g.*, propranolol, timolol, tertalolol, carteolol, nadolol, betaxolol, penbutolol, acetobutolol, atenolol, metoprolol, or carvedilol), verapamil, diltiazem, or diltiazem. Treatment of hypertrophy associated with high blood pressure may require the use of antihypertensive drug therapy, using calcium channel blockers, *e.g.*, diltiazem, nifedipine, verapamil, or nicardipine; β -adrenergic blocking agents; diuretics, *e.g.*, chlorothiazide, hydrochlorothiazide, hydroflumethazide, methylchlorothiazide, benzthiazide, dichlorphenamide, acetazolamide, or indapamide; and/or ACE-inhibitors, *e.g.*, quinapril, captopril, enalapril, ramipril, benazepril, fosinopril, or lisinopril.

For other indications, PRO polypeptides or their agonists or antagonists may be combined with other agents beneficial to the treatment of the bone and/or cartilage defect, wound, or tissue in question. These agents include various growth factors such as EGF, PDGF, TGF- α or TGF- β , IGF, FGF, and CTGF.

In addition, PRO polypeptides or their agonists or antagonists used to treat cancer may be combined with cytotoxic, chemotherapeutic, or growth-inhibitory agents as identified above. Also, for cancer treatment, the PRO polypeptide or agonist or antagonist thereof is suitably administered serially or in combination with radiological treatments, whether involving irradiation or administration of radioactive substances.

The effective amounts of the therapeutic agents administered in combination with the PRO polypeptide or agonist or antagonist thereof will be at the physician's or veterinarian's discretion. Dosage administration and adjustment is done to achieve maximal management of the conditions to be treated. For example, for treating hypertension, these amounts ideally take into account use of diuretics or digitalis, and conditions such as hyper- or hypotension, renal impairment, etc. The dose will additionally depend on such factors as the type of the therapeutic agent to be used and the specific patient being treated. Typically, the amount employed will be the same dose as that used, if the given therapeutic agent is administered without the PRO polypeptide.

5.2.4.13. Articles of Manufacture

An article of manufacture such as a kit containing the PRO polypeptide or agonists or antagonists thereof useful for the diagnosis or treatment of the disorders described above comprises at least a container and a label. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition that is effective for diagnosing or treating the condition and may have a sterile access port (for example, the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The active agent in the composition is the PRO polypeptide or an agonist or antagonist thereto. The label on, or associated with, the container indicates that the composition is used for diagnosing or treating the condition of choice. The article of manufacture may further comprise a second container comprising a pharmaceutically-acceptable buffer, such as phosphate-buffered saline, Ringer's solution, and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for use. The article of manufacture may also comprise a second or third container with another active agent as described above.

5.2.5. Antibodies

Some of the most promising drug candidates according to the present invention are antibodies and antibody fragments that may inhibit the production or the gene product of the genes identified herein and/or reduce the activity of the gene products.

5.2.5.1. Polyclonal Antibodies

Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include the PRO polypeptide or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include, but are not limited to, keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants that may be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A or synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

5.2.5.2. Monoclonal Antibodies

The anti-PRO antibodies may, alternatively, be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, *Nature*, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal is typically immunized with an

immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized *in vitro*.

The immunizing agent will typically include the PRO polypeptide or a fusion protein thereof. Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell. Goding, Monoclonal Antibodies: Principles and Practice (New York: Academic Press, 1986), pp. 59-103. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine, and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high-level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies. Kozbor, J. Immunol., 133:3001 (1984); Brodeur *et al.*, Monoclonal Antibody Production Techniques and Applications (Marcel Dekker, Inc.: New York, 1987) pp. 51-63.

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against the PRO polypeptide. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980).

After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods. Goding, *supra*. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells may be grown *in vivo* as ascites in a mammal.

The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the

invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy- and light-chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; Morrison *et al.*, *supra*) or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

The antibodies may be monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy-chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

In vitro methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce fragments thereof, particularly Fab fragments, can be accomplished using routine techniques known in the art.

5.2.5.3. Human and Humanized Antibodies

The anti-PRO antibodies may further comprise humanized antibodies or human antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains, or fragments thereof (such as Fv, Fab, Fab', F(ab')₂, or other antigen-binding subsequences of antibodies) that contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a CDR of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat, or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues that are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin, and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody preferably also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. Jones *et al.*, Nature, **321**: 522-525 (1986); Riechmann *et al.*, Nature, **332**: 323-329 (1988); Presta, Curr. Op. Struct. Biol., **2**: 593-596 (1992).

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source that is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al.*, Nature, **321**: 522-525 (1986); Riechmann *et al.*, Nature, **332**: 323-327 (1988); Verhoeven *et al.*, Science, **239**: 1534-

1536 (1988)), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

Human antibodies can also be produced using various techniques known in the art, including phage display libraries. Hoogenboom and Winter, *J. Mol. Biol.*, 227: 381 (1991); Marks *et al.*, *J. Mol. Biol.*, 222: 581 (1991). The techniques of Cole *et al.* and Boerner *et al.* are also available for the preparation of human monoclonal antibodies. Cole *et al.*, *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985) and Boerner *et al.*, *J. Immunol.*, 147(1): 86-95 (1991). Similarly, human antibodies can be made by introducing human immunoglobulin loci into transgenic animals, *e.g.*, mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed that closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; and 5,661,016, and in the following scientific publications: Marks *et al.*, *Bio/Technology*, 10: 779-783 (1992); Lonberg *et al.*, *Nature*, 368: 856-859 (1994); Morrison, *Nature*, 368: 812-813 (1994); Fishwild *et al.*, *Nature Biotechnology*, 14: 845-851 (1996); Neuberger, *Nature Biotechnology*, 14: 826 (1996); Lonberg and Huszar, *Intern. Rev. Immunol.*, 13: 65-93 (1995).

5.2.5.4. Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for the PRO polypeptide, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities. Milstein and Cuello, *Nature*, 305: 537-539 (1983). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker *et al.*, *EMBO J.*, 10: 3655-3659 (1991).

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant-domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further

details of generating bispecific antibodies, *see*, for example, Suresh *et al.*, Methods in Enzymology, **121**: 210 (1986).

5.2.5.5. Heteroconjugate Antibodies

Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune-system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection. WO 91/00360; WO 92/200373; EP 03089. It is contemplated that the antibodies may be prepared *in vitro* using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide-exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

5.2.5.6. Effector Function Engineering

It may be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, *e.g.*, the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) may be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). *See*, Caron *et al.*, J. Exp. Med., **176**: 1191-1195 (1992) and Shopes, J. Immunol., **148**: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using heterobifunctional cross-linkers as described in Wolff *et al.*, Cancer Research, **53**: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. *See*, Stevenson *et al.*, Anti-Cancer Drug Design, **3**: 219-230 (1989).

5.2.5.7. Immunoconjugates

The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (*e.g.*, an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (*i.e.*, a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolacca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y , and ^{186}Re .

Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives

of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta *et al.*, Science, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See, WO94/11026.

In another embodiment, the antibody may be conjugated to a "receptor" (such as streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) that is conjugated to a cytotoxic agent (e.g., a radionucleotide).

5.2.5.8. Immunoliposomes

The antibodies disclosed herein may also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein *et al.*, Proc. Natl. Acad. Sci. USA, 82: 3688 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. USA, 77: 4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin *et al.*, J. Biol. Chem., 257: 286-288 (1982) via a disulfide-interchange reaction. A chemotherapeutic agent (such as Doxorubicin) is optionally contained within the liposome. See, Gabizon *et al.*, J. National Cancer Inst., 81(19): 1484 (1989).

5.2.5.9. Pharmaceutical Compositions of Antibodies

Antibodies specifically binding a PRO polypeptide identified herein, as well as other molecules identified by the screening assays disclosed hereinbefore, can be administered for the treatment of various disorders as noted above and below in the form of pharmaceutical compositions.

If the PRO polypeptide is intracellular and whole antibodies are used as inhibitors, internalizing antibodies are preferred. However, lipofections or liposomes can also be used to deliver the antibody, or an antibody fragment, into cells. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. See, e.g., Marasco *et al.*, Proc. Natl. Acad. Sci. USA, 90: 7889-7893 (1993).

The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition may comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, *supra*.

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOTTM (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

5.2.5.10. Methods of Treatment using the Antibody

It is contemplated that the antibodies to a PRO polypeptide may be used to treat various cardiovascular, endothelial, and angiogenic conditions as noted above.

The antibodies are administered to a mammal, preferably a human, in accord with known methods, such as intravenous administration as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or inhalation routes. Intravenous administration of the antibody is preferred.

Other therapeutic regimens may be combined with the administration of the antibodies of the instant invention as noted above. For example, if the antibodies are to treat cancer, the patient to be treated with such

antibodies may also receive radiation therapy. Alternatively, or in addition, a chemotherapeutic agent may be administered to the patient. Preparation and dosing schedules for such chemotherapeutic agents may be used according to manufacturers' instructions or as determined empirically by the skilled practitioner. Preparation and dosing schedules for such chemotherapy are also described in Chemotherapy Service, Ed., M.C. Perry (Williams & Wilkins: Baltimore, MD, 1992). The chemotherapeutic agent may precede, or follow administration of the antibody, or may be given simultaneously therewith. The antibody may be combined with an anti-estrogen compound such as tamoxifen or EVISTA™ or an anti-progesterone such as onapristone (*see*, EP 616812) in dosages known for such molecules.

If the antibodies are used for treating cancer, it may be desirable also to administer antibodies against other tumor-associated antigens, such as antibodies that bind to one or more of the ErbB2, EGFR, ErbB3, ErbB4, or VEGF receptor(s). These also include the agents set forth above. Also, the antibody is suitably administered serially or in combination with radiological treatments, whether involving irradiation or administration of radioactive substances. Alternatively, or in addition, two or more antibodies binding the same or two or more different antigens disclosed herein may be co-administered to the patient. Sometimes, it may be beneficial also to administer one or more cytokines to the patient. In a preferred embodiment, the antibodies herein are co-administered with a growth-inhibitory agent. For example, the growth-inhibitory agent may be administered first, followed by an antibody of the present invention. However, simultaneous administration or administration of the antibody of the present invention first is also contemplated. Suitable dosages for the growth-inhibitory agent are those presently used and may be lowered due to the combined action (synergy) of the growth-inhibitory agent and the antibody herein.

In one embodiment, vascularization of tumors is attacked in combination therapy. The anti-PRO polypeptide antibody and another antibody (*e.g.*, anti-VEGF) are administered to tumor-bearing patients at therapeutically effective doses as determined, for example, by observing necrosis of the tumor or its metastatic foci, if any. This therapy is continued until such time as no further beneficial effect is observed or clinical examination shows no trace of the tumor or any metastatic foci. Then TNF is administered, alone or in combination with an auxiliary agent such as alpha-, beta-, or gamma-interferon, anti-HER2 antibody, heregulin, anti-heregulin antibody, D-factor, interleukin-1 (IL-1), interleukin-2 (IL-2), granulocyte-macrophage colony stimulating factor (GM-CSF), or agents that promote microvascular coagulation in tumors, such as anti-protein C antibody, anti-protein S antibody, or C4b binding protein (*see*, WO 91/01753, published 21 February 1991), or heat or radiation.

Since the auxiliary agents will vary in their effectiveness, it is desirable to compare their impact on the tumor by matrix screening in conventional fashion. The administration of anti-PRO polypeptide antibody and TNF is repeated until the desired clinical effect is achieved. Alternatively, the anti-PRO polypeptide antibody is administered together with TNF and, optionally, auxiliary agent(s). In instances where solid tumors are found in the limbs or in other locations susceptible to isolation from the general circulation, the therapeutic agents described herein are administered to the isolated tumor or organ. In other embodiments, a FGF or PDGF antagonist, such as an anti-FGF or an anti-PDGF neutralizing antibody, is administered to the patient in conjunction with the anti-PRO

polypeptide antibody. Treatment with anti-PRO polypeptide antibodies preferably may be suspended during periods of wound healing or desirable neovascularization.

For the prevention or treatment of cardiovascular, endothelial, and angiogenic disorder, the appropriate dosage of an antibody herein will depend on the type of disorder to be treated, as defined above, the severity and course of the disease, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antibody, and the discretion of the attending physician. The antibody is suitably administered to the patient at one time or over a series of treatments.

For example, depending on the type and severity of the disorder, about 1 $\mu\text{g/kg}$ to 50 mg/kg (e.g., 0.1-20 mg/kg) of antibody is an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. A typical daily or weekly dosage might range from about 1 $\mu\text{g/kg}$ to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is repeated or sustained until a desired suppression of disorder symptoms occurs. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays, including, for example, radiographic tumor imaging.

5.2.5.11. Articles of Manufacture with Antibodies

An article of manufacture containing a container with the antibody and a label is also provided. Such articles are described above, wherein the active agent is an anti-PRO antibody.

5.2.5.12. Diagnosis and Prognosis of Tumors using Antibodies

If the indication for which the antibodies are used is cancer, while cell-surface proteins, such as growth receptors over expressed in certain tumors, are excellent targets for drug candidates or tumor (e.g., cancer) treatment, the same proteins along with PRO polypeptides find additional use in the diagnosis and prognosis of tumors. For example, antibodies directed against the PRO polypeptides may be used as tumor diagnostics or prognostics.

For example, antibodies, including antibody fragments, can be used qualitatively or quantitatively to detect the expression of genes including the gene encoding the PRO polypeptide. The antibody preferably is equipped with a detectable, e.g., fluorescent label, and binding can be monitored by light microscopy, flow cytometry, fluorimetry, or other techniques known in the art. Such binding assays are performed essentially as described above.

In situ detection of antibody binding to the marker gene products can be performed, for example, by immunofluorescence or immunoelectron microscopy. For this purpose, a histological specimen is removed from the patient, and a labeled antibody is applied to it, preferably by overlaying the antibody on a biological sample. This procedure also allows for determining the distribution of the marker gene product in the tissue examined. It will be apparent to those skilled in the art that a wide variety of histological methods are readily available for *in situ* detection.

The following Examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

The disclosures of all patent and literature references cited in the present specification are hereby incorporated by reference in their entirety.

6. EXAMPLES

Commercially available reagents referred to in the Examples were used according to manufacturer's instructions unless otherwise indicated. The source of those cells identified in the following Examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Manassas, VA. Unless otherwise noted, the present invention uses standard procedures of recombinant DNA technology, such as those described hereinabove and in the following textbooks: Sambrook *et al.*, supra; Ausubel *et al.*, Current Protocols in Molecular Biology (Green Publishing Associates and Wiley Interscience, N.Y., 1989); Innis *et al.*, PCR Protocols: A Guide to Methods and Applications (Academic Press, Inc.: N.Y., 1990); Harlow *et al.*, Antibodies: A Laboratory Manual (Cold Spring Harbor Press: Cold Spring Harbor, 1988); Gait, Oligonucleotide Synthesis (IRL Press: Oxford, 1984); Freshney, Animal Cell Culture, 1987; Coligan *et al.*, Current Protocols in Immunology, 1991.

6.1. EXAMPLE 1: Extracellular Domain Homology Screening to Identify Novel Polypeptides and cDNA Encoding Therefor

The extracellular domain (ECD) sequences (including the secretion signal sequence, if any) from about 950 known secreted proteins from the Swiss-Prot public database were used to search EST databases. The EST databases included public databases (*e.g.*, Dayhoff, GenBank), and proprietary databases (*e.g.* LIFESEQ®, Incyte Pharmaceuticals, Palo Alto, CA). The search was performed using the computer program BLAST or BLAST-2 (Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)) as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequences. Those comparisons with a BLAST score of 70 (or in some cases, 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, WA).

Using this extracellular domain homology screen, consensus DNA sequences were assembled relative to the other identified EST sequences using phrap. In addition, the consensus DNA sequences obtained were often (but not always) extended using repeated cycles of BLAST or BLAST-2 and phrap to extend the consensus sequence as far as possible using the sources of EST sequences discussed above.

Based upon the consensus sequences obtained as described above, oligonucleotides were then synthesized and used to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for a PRO polypeptide. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In some cases, additional oligonucleotides are synthesized when the consensus sequence is greater than about 1-1.5 kbp. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel *et al.*, Current Protocols in Molecular

Biology, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs.

5 The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; *see*, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

6.2. EXAMPLE 2: Isolation of cDNA Clones by Amylase Screening

10 6.2.1. Preparation of oligo dT primed cDNA library

mRNA was isolated from a human tissue of interest using reagents and protocols from Invitrogen, San Diego, CA (Fast Track 2). This RNA was used to generate an oligo dT primed cDNA library in the vector pRK5D using reagents and protocols from Life Technologies, Gaithersburg, MD (Super Script Plasmid System). In this procedure, the double stranded cDNA was sized to greater than 1000 bp and the SalI/NotI linked cDNA was 15 cloned into XhoI/NotI cleaved vector. pRK5D is a cloning vector that has an sp6 transcription initiation site followed by an SfiI restriction enzyme site preceding the XhoI/NotI cDNA cloning sites.

6.2.2. Preparation of random primed cDNA library

A secondary cDNA library was generated in order to preferentially represent the 5' ends of the primary cDNA clones. Sp6 RNA was generated from the primary library (described above), and this RNA was used 20 to generate a random primed cDNA library in the vector pSST-AMY.0 using reagents and protocols from Life Technologies (Super Script Plasmid System, referenced above). In this procedure the double stranded cDNA was sized to 500-1000 bp, linked with blunt to NotI adaptors, cleaved with SfiI, and cloned into SfiI/NotI cleaved vector. pSST-AMY.0 is a cloning vector that has a yeast alcohol dehydrogenase promoter preceding the cDNA cloning sites and the mouse amylase sequence (the mature sequence without the secretion signal) followed by the 25 yeast alcohol dehydrogenase terminator, after the cloning sites. Thus, cDNAs cloned into this vector that are fused in frame with amylase sequence will lead to the secretion of amylase from appropriately transfected yeast colonies.

6.2.3. Transformation and Detection

DNA from the library described in paragraph 2 above was chilled on ice to which was added electrocompetent DH10B bacteria (Life Technologies, 20 ml). The bacteria and vector mixture was then 30 electroporated as recommended by the manufacturer. Subsequently, SOC media (Life Technologies, 1 ml) was added and the mixture was incubated at 37°C for 30 minutes. The transformants were then plated onto 20 standard 150 mm LB plates containing ampicillin and incubated for 16 hours (37°C). Positive colonies were scraped off the plates and the DNA was isolated from the bacterial pellet using standard protocols, *e.g.*, CsCl-gradient. The purified DNA was then carried on to the yeast protocols below.

The yeast methods were divided into three categories: (1) Transformation of yeast with the plasmid/cDNA combined vector; (2) Detection and isolation of yeast clones secreting amylase; and (3) PCR amplification of the insert directly from the yeast colony and purification of the DNA for sequencing and further analysis.

5 The yeast strain used was HD56-5A (ATCC-90785). This strain has the following genotype: MAT alpha, ura3-52, leu2-3, leu2-112, his3-11, his3-15, MAL⁺, SUC⁺, GAL⁺. Preferably, yeast mutants can be employed that have deficient post-translational pathways. Such mutants may have translocation deficient alleles in *sec71*, *sec72*, *sec62*, with truncated *sec71* being most preferred. Alternatively, antagonists (including antisense nucleotides and/or ligands) which interfere with the normal operation of these genes, other proteins implicated in this post translation pathway (e.g., SEC61p, SEC72p, SEC62p, SEC63p, TDJ1p or SSA1p-4p) or the complex formation
10 of these proteins may also be preferably employed in combination with the amylase-expressing yeast.

Transformation was performed based on the protocol outlined by Gietz *et al.*, Nucl. Acid. Res., 20:1425 (1992). Transformed cells were then inoculated from agar into YEPD complex media broth (100 ml) and grown overnight at 30°C. The YEPD broth was prepared as described in Kaiser *et al.*, Methods in Yeast Genetics, Cold Spring Harbor Press, Cold Spring Harbor, NY, p. 207 (1994). The overnight culture was then diluted to about 2
15 x 10⁶ cells/ml (approx. OD₆₀₀=0.1) into fresh YEPD broth (500 ml) and regrown to 1 x 10⁷ cells/ml (approx. OD₆₀₀=0.4-0.5).

The cells were then harvested and prepared for transformation by transfer into GS3 rotor bottles in a Sorval GS3 rotor at 5,000 rpm for 5 minutes, the supernatant discarded, and then resuspended into sterile water, and centrifuged again in 50 ml falcon tubes at 3,500 rpm in a Beckman GS-6KR centrifuge. The supernatant was
20 discarded and the cells were subsequently washed with LiAc/TE (10 ml, 10 mM Tris-HCl, 1 mM EDTA pH 7.5, 100 mM Li₂OOCCH₃), and resuspended into LiAc/TE (2.5 ml).

Transformation took place by mixing the prepared cells (100 µl) with freshly denatured single stranded salmon testes DNA (Lofstrand Labs, Gaithersburg, MD) and transforming DNA (1 µg, vol. < 10 µl) in microfuge tubes. The mixture was mixed briefly by vortexing, then 40% PEG/TE (600 µl, 40% polyethylene glycol-4000,
25 10 mM Tris-HCl, 1 mM EDTA, 100 mM Li₂OOCCH₃, pH 7.5) was added. This mixture was gently mixed and incubated at 30°C while agitating for 30 minutes. The cells were then heat shocked at 42°C for 15 minutes, and the reaction vessel centrifuged in a microfuge at 12,000 rpm for 5-10 seconds, decanted and resuspended into TE (500 µl, 10 mM Tris-HCl, 1 mM EDTA pH 7.5) followed by recentrifugation. The cells were then diluted into TE (1 ml) and aliquots (200 µl) were spread onto the selective media previously prepared in 150 mm growth plates
30 (VWR).

Alternatively, instead of multiple small reactions, the transformation was performed using a single, large scale reaction, wherein reagent amounts were scaled up accordingly.

The selective media used was a synthetic complete dextrose agar lacking uracil (SCD-Ura) prepared as described in Kaiser *et al.*, Methods in Yeast Genetics, Cold Spring Harbor Press, Cold Spring Harbor, NY, p. 208-
35 210 (1994). Transformants were grown at 30°C for 2-3 days.

The detection of colonies secreting amylase was performed by including red starch in the selective growth media. Starch was coupled to the red dye (Reactive Red-120, Sigma) as per the procedure described by Biely *et*

al., Anal. Biochem., 172:176-179 (1988). The coupled starch was incorporated into the SCD-Ura agar plates at a final concentration of 0.15% (w/v), and was buffered with potassium phosphate to a pH of 7.0 (50-100 mM final concentration).

The positive colonies were picked and streaked across fresh selective media (onto 150 mm plates) in order to obtain well isolated and identifiable single colonies. Well isolated single colonies positive for amylase secretion were detected by direct incorporation of red starch into buffered SCD-Ura agar. Positive colonies were determined by their ability to break down starch resulting in a clear halo around the positive colony visualized directly.

6.2.4. Isolation of DNA by PCR Amplification

When a positive colony was isolated, a portion of it was picked by a toothpick and diluted into sterile water (30 µl) in a 96 well plate. At this time, the positive colonies were either frozen and stored for subsequent analysis or immediately amplified. An aliquot of cells (5 µl) was used as a template for the PCR reaction in a 25 µl volume containing: 0.5 µl KlenTaq (Clontech, Palo Alto, CA); 4.0 µl 10 mM dNTP's (Perkin Elmer-Cetus); 2.5 µl Kentaq buffer (Clontech); 0.25 µl forward oligo 1; 0.25 µl reverse oligo 2; 12.5 µl distilled water. The sequence of the forward oligonucleotide 1 was:

5'-TGTAACGACGGCCAGTTAAATAGACCTGCAATTATTAATCT-3' (SEQ ID NO:382)

The sequence of reverse oligonucleotide 2 was:

5'-CAGGAAACAGCTATGACCACCTGCACACCTGCAAATCCATT-3' (SEQ ID NO:383)

PCR was then performed as follows:

a.	Denature	92°C, 5 minutes
b.	3 cycles of:	
	Denature	92°C, 30 seconds
	Anneal	59°C, 30 seconds
	Extend	72°C, 60 seconds
c.	3 cycles of:	
	Denature	92°C, 30 seconds
	Anneal	57°C, 30 seconds
	Extend	72°C, 60 seconds
d.	25 cycles of:	
	Denature	92°C, 30 seconds
	Anneal	55°C, 30 seconds
	Extend	72°C, 60 seconds
e.	Hold	4°C

The underlined regions of the oligonucleotides annealed to the ADH promoter region and the amylase region, respectively, and amplified a 307 bp region from vector pSST-AMY.0 when no insert was present. Typically, the first 18 nucleotides of the 5' end of these oligonucleotides contained annealing sites for the sequencing primers. Thus, the total product of the PCR reaction from an empty vector was 343 bp. However, signal sequence-fused cDNA resulted in considerably longer nucleotide sequences.

Following the PCR, an aliquot of the reaction (5 μ l) was examined by agarose gel electrophoresis in a 1% agarose gel using a Tris-Borate-EDTA (TBE) buffering system as described by Sambrook *et al.*, *supra*. Clones resulting in a single strong PCR product larger than 400 bp were further analyzed by DNA sequencing after purification with a 96 Qiaquick PCR clean-up column (Qiagen Inc., Chatsworth, CA).

6.3. EXAMPLE 3: Isolation of cDNA Clones Using Signal Algorithm Analysis

Various polypeptide-encoding nucleic acid sequences were identified by applying a proprietary signal sequence finding algorithm developed by Genentech, Inc., (South San Francisco, CA) upon ESTs as well as clustered and assembled EST fragments from public (e.g., GenBank) and/or private (LIFESEQ®, Incyte Pharmaceuticals, Inc., Palo Alto, CA) databases. The signal sequence algorithm computes a secretion signal score based on the character of the DNA nucleotides surrounding the first and optionally the second methionine codon(s) (ATG) at the 5'-end of the sequence or sequence fragment under consideration. The nucleotides following the first ATG must code for at least 35 unambiguous amino acids without any stop codons. If the first ATG has the required amino acids, the second is not examined. If neither meets the requirement, the candidate sequence is not scored. In order to determine whether the EST sequence contains an authentic signal sequence, the DNA and corresponding amino acid sequences surrounding the ATG codon are scored using a set of seven sensors (evaluation parameters) known to be associated with secretion signals. Use of this algorithm resulted in the identification of numerous polypeptide-encoding nucleic acid sequences.

6.4. EXAMPLE 4: Isolation of cDNA clones Encoding Human PRO Polypeptides

Using the techniques described in Examples 1 to 3 above, numerous full-length cDNA clones were identified as encoding PRO polypeptides as disclosed herein. These cDNAs were then deposited under the terms of the Budapest Treaty with the American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209, USA (ATCC) as shown in Table 7 below.

Table 7

	<u>Material</u>	<u>ATCC Dep. No.</u>	<u>Deposit Date</u>
25	23330-1390	209775	4/14/1998
	23339-1130	209282	9/18/1997
	26846-1397	203406	10/27/1998
	26847-1395	209772	4/14/1998
	27865-1091	209296	9/23/1997
30	30868-1156	1437-PTA	3/2/2000
	30871-1157	209380	10/16/1997
	32286-1191	209385	10/16/1997
	33089-1132	209262	9/16/1997
	33092-1202	209420	10/28/1997

	33100-1159	209377	10/16/1997
	33223-1136	209264	9/16/1997
	34392-1170	209526	12/10/1997
	34431-1177	209399	10/17/1997
5	34433-1308	209719	3/31/1998
	34434-1139	209252	9/16/1997
	35600-1162	209370	10/16/1997
	35673-1201	209418	10/28/1997
	35880-1160	209379	10/16/1997
10	35918-1174	209402	10/17/1997
	36350-1158	209378	10/16/1997
	36638-1056	209456	11/12/1997
	38268-1188	209421	10/28/1997
	40370-1217	209485	11/21/1997
15	40628-1216	209432	11/7/1997
	43316-1237	209487	11/21/1997
	44196-1353	209847	5/6/1998
	45409-2511	203579	1/12/1999
	45419-1252	209616	2/5/1998
20	46777-1253	209619	2/5/1998
	48336-1309	209669	3/11/1998
	48606-1479	203040	7/1/1998
	49435-1219	209480	11/21/1997
	49631-1328	209806	4/28/1998
25	50919-1361	209848	5/6/1998
	50920-1325	209700	3/26/1998
	50921-1458	209859	5/12/1998
	52758-1399	209773	4/14/1998
	53517-1366-1	209802	4/23/1998
30	53915-1258	209593	1/21/1998
	53974-1401	209774	4/14/1998
	53987-1438	209858	5/12/1998
	56047-1456	209948	6/9/1998
	56050-1455	203011	6/23/1998
35	56110-1437	203113	8/11/1998
	56405-1357	209849	5/6/1998
	56433-1406	209857	5/12/1998

	56439-1376	209864	5/14/1998
	56529-1647	203293	9/29/1998
	56865-1491	203022	6/23/1998
	56965-1356	209842	5/6/1998
5	57033-1403-1	209905	5/27/1998
	57037-1444	209903	5/27/1998
	57039-1402	209777	4/14/1998
	57689-1385	209869	5/14/1998
	57690-1374	209950	6/9/1998
10	57694-1341	203017	6/23/1998
	57695-1340	203006	6/23/1998
	57699-1412	203020	6/23/1998
	57700-1408	203583	1/12/1999
	57708-1411	203021	6/23/1998
15	57838-1337	203014	6/23/1998
	58847-1383	209879	5/20/1998
	58852-1637	203271	9/22/1998
	58853-1423	203016	6/23/1998
	59212-1627	203245	9/9/1998
20	59220-1514	209962	6/9/1998
	59493-1420	203050	7/1/1998
	59497-1496	209941	6/4/1998
	59586-1520	203288	9/29/1998
	59588-1571	203106	8/11/1998
25	59620-1463	209989	6/16/1998
	59622-1334	209984	6/16/1998
	59777-1480	203111	8/11/1998
	59848-1512	203088	8/4/1998
	59849-1504	209986	6/16/1998
30	60621-1516	203091	8/4/1998
	60622-1525	203090	8/4/1998
	60764-1533	203452	11/10/1998
	60783-1611	203130	8/18/1998
	61755-1554	203112	8/11/1998
35	62306-1570	203254	9/9/1998
	62312-2558	203836	3/9/1999
	62814-1521	203093	8/4/1998

	62872-1509	203100	8/4/1998
	64883-1526	203253	9/9/1998
	64886-1601	203241	9/9/1998
	64889-1541	203250	9/9/1998
5	64896-1539	203238	9/9/1998
	64897-1628	203216	9/15/1998
	64903-1553	203223	9/15/1998
	64908-1163-1	203243	9/9/1998
	64950-1590	203224	9/15/1998
10	65402-1540	203252	9/9/1998
	65404-1551	203244	9/9/1998
	65405-1547	203476	11/17/1998
	65410-1569	203231	9/15/1998
	65412-1523	203094	8/4/1998
15	66307-2661	431-PTA	7/27/1999
	66526-1616	203246	9/9/1998
	66659-1593	203269	9/22/1998
	66660-1585	203279	9/22/1998
	66667-1596	203267	9/22/1998
20	66672-1586	203265	9/22/1998
	66675-1587	203282	9/22/1998
	67300-1605	203163	8/25/1998
	68818-2536	203657	2/9/1999
	68862-2546	203652	2/9/1999
25	68872-1620	203160	8/25/1998
	71290-1630	203275	9/22/1998
	73736-1657	203466	11/17/1998
	73739-1645	203270	9/22/1998
	73742-1662	203316	10/6/1998
30	76385-1692	203664	2/9/1999
	76393-1664	203323	10/6/1998
	76399-1700	203472	11/17/1998
	76400-2528	203573	1/12/1999
	76510-2504	203477	11/17/1998
35	76529-1666	203315	10/6/1998
	76532-1702	203473	11/17/1998
	76541-1675	203409	10/27/1998

	77503-1686	203362	10/20/1998
	77624-2515	203553	12/22/1998
	79230-2525	203549	12/22/1998
	79862-2522	203550	12/22/1998
5	80145-2594	204-PTA	6/8/1999
	80899-2501	203539	12/15/1998
	81754-2532	203542	12/15/1998
	81757-2512	203543	12/15/1998
	81761-2583	203862	3/23/1999
10	82358-2738	510-PTA	8/10/1999
	82364-2538	203603	1/20/1999
	82403-2959	2317-PTA	8/1/2000
	83500-2506	203391	10/29/1998
	83560-2569	203816	3/2/1999
15	84210-2576	203818	3/2/1999
	84920-2614	203966	4/27/1999
	86576-2595	203868	3/23/1999
	92218-2554	203834	3/9/1999
	92233-2599	134-PTA	5/25/1999
20	92256-2596	203891	3/30/1999
	92265-2669	256-PTA	6/22/1999
	92274-2617	203971	4/27/1999
	92929-2534-1	203586	1/12/1999
	93011-2637	20-PTA	5/4/1999
25	94854-2586	203864	3/23/1999
	96787-2534-1	203589	1/12/1999
	96867-2620	203972	4/27/1999
	96872-2674	550-PTA	8/17/1999
	96878-2626	23-PTA	5/4/1999
30	96889-2641	119-PTA	5/25/1999
	100312-2645	44-PTA	5/11/1999
	105782-2693	387-PTA	7/20/1999
	105849-2704	473-PTA	8/3/1999
	108725-2766	863-PTA	10/19/1999
35	108769-2765	861-PTA	10/19/1999
	119498-2965	2298-PTA	7/25/2000
	119535-2756	613-PTA	8/31/1999

	125185-2806	1031-PTA	12/7/1999
	131639-2874	1784-PTA	4/25/2000
	139623-2893	1670-PTA	4/11/2000
	143076-2787	1028-PTA	12/7/1999
5	143276-2975	2387-PTA	8/8/2000
	164625-2890	1535-PTA	3/21/2000
	167678-2963	2302-PTA	7/25/2000
	170021-2923	1906-PTA	5/23/2000
	170212-3000	2583-PTA	10/10/2000
10	177313-2982	2251-PTA	7/19/2000

These deposits were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable culture of the deposit for 30 years from the date of deposit. The deposits will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the culture of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC § 122 and the Commissioner's rules pursuant thereto (including 37 CFR § 1.14 with particular reference to 886 OG 638).

The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same. Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

6.5 EXAMPLE 5: Isolation of cDNA clones Encoding Human PRO1873, PRO7223, PRO7248, PRO730, PRO532, PRO7261, PRO734, PRO771, PRO2010, PRO5723, PRO3444, PRO9940, PRO3562, PRO10008, PRO5730, PRO6008, PRO4527, PRO4538 and PRO4553

DNA molecules encoding the PRO1873, PRO7223, PRO7248, PRO730, PRO532, PRO7261, PRO734, PRO771, PRO2010, PRO5723, PRO3444, PRO9940, PRO3562, PRO10008, PRO5730, PRO6008, PRO4527, PRO4538 and PRO4553 polypeptides shown in the accompanying figures were obtained through GenBank.

6.6. EXAMPLE 6: Use of PRO as a Hybridization Probe

The following method describes use of a nucleotide sequence encoding PRO as a hybridization probe.

DNA comprising the coding sequence of full-length or mature PRO (as shown in accompanying figures) or a fragment thereof is employed as a probe to screen for homologous DNAs (such as those encoding naturally-occurring variants of PRO) in human tissue cDNA libraries or human tissue genomic libraries.

5 Hybridization and washing of filters containing either library DNAs is performed under the following high-stringency conditions. Hybridization of radiolabeled probe derived from the gene encoding PRO polypeptide to the filters is performed in a solution of 50% formamide, 5x SSC, 0.1% SDS, 0.1% sodium pyrophosphate, 50 mM sodium phosphate, pH 6.8, 2x Denhardt's solution, and 10% dextran sulfate at 42°C for 20 hours. Washing of the filters is performed in an aqueous solution of 0.1x SSC and 0.1% SDS at 42°C.

10 DNAs having a desired sequence identity with the DNA encoding full-length native sequence can then be identified using standard techniques known in the art.

6.7. EXAMPLE 7: Expression of PRO in *E. coli*

This example illustrates preparation of an unglycosylated form of PRO by recombinant expression in *E. coli*.

15 The DNA sequence encoding PRO is initially amplified using selected PCR primers. The primers should contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector. A variety of expression vectors may be employed. An example of a suitable vector is pBR322 (derived from *E. coli*; see, Bolivar *et al.*, Gene, 2:95 (1977)) which contains genes for ampicillin and tetracycline resistance. The vector is digested with restriction enzyme and dephosphorylated. The PCR amplified sequences are then ligated into the vector. The vector will preferably include sequences which encode for an antibiotic resistance gene, a trp promoter, a poly-His leader (including the first six STII codons, poly-His sequence, and enterokinase cleavage site),
20 the PRO coding region, lambda transcriptional terminator, and an argU gene.

The ligation mixture is then used to transform a selected *E. coli* strain using the methods described in Sambrook *et al.*, *supra*. Transformants are identified by their ability to grow on LB plates and antibiotic resistant colonies are then selected. Plasmid DNA can be isolated and confirmed by restriction analysis and DNA
25 sequencing.

Selected clones can be grown overnight in liquid culture medium such as LB broth supplemented with antibiotics. The overnight culture may subsequently be used to inoculate a larger scale culture. The cells are then grown to a desired optical density, during which the expression promoter is turned on.

30 After culturing the cells for several more hours, the cells can be harvested by centrifugation. The cell pellet obtained by the centrifugation can be solubilized using various agents known in the art, and the solubilized PRO protein can then be purified using a metal chelating column under conditions that allow tight binding of the protein.

PRO may be expressed in *E. coli* in a poly-His tagged form, using the following procedure. The DNA encoding PRO is initially amplified using selected PCR primers. The primers will contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector, and other useful sequences
35 providing for efficient and reliable translation initiation, rapid purification on a metal chelation column, and proteolytic removal with enterokinase. The PCR-amplified, poly-His tagged sequences are then ligated into an

expression vector, which is used to transform an *E. coli* host based on strain 52 (W3110 fuhA(tonA) lon galE rpoHts(htpRts) clpP(lacIq). Transformants are first grown in LB containing 50 mg/ml carbenicillin at 30°C with shaking until an OD₆₀₀ of 3-5 is reached. Cultures are then diluted 50-100 fold into CRAP media (prepared by mixing 3.57 g (NH₄)₂SO₄, 0.71 g sodium citrate·2H₂O, 1.07 g KCl, 5.36 g Difco yeast extract, 5.36 g Sheffield hycase SF in 500 ml water, as well as 110 mM MPOS, pH 7.3, 0.55% (w/v) glucose and 7 mM MgSO₄) and grown for approximately 20-30 hours at 30°C with shaking. Samples are removed to verify expression by SDS-PAGE analysis, and the bulk culture is centrifuged to pellet the cells. Cell pellets are frozen until purification and refolding.

E. coli paste from 0.5 to 1 L fermentations (6-10 g pellets) is resuspended in 10 volumes (w/v) in 7 M guanidine, 20 mM Tris, pH 8 buffer. Solid sodium sulfite and sodium tetrathionate is added to make final concentrations of 0.1M and 0.02 M, respectively, and the solution is stirred overnight at 4°C. This step results in a denatured protein with all cysteine residues blocked by sulfitolization. The solution is centrifuged at 40,000 rpm in a Beckman Ultracentrifuge for 30 min. The supernatant is diluted with 3-5 volumes of metal chelate column buffer (6 M guanidine, 20 mM Tris, pH 7.4) and filtered through 0.22 micron filters to clarify. The clarified extract is loaded onto a 5 ml Qiagen Ni²⁺-NTA metal chelate column equilibrated in the metal chelate column buffer. The column is washed with additional buffer containing 50 mM imidazole (Calbiochem, Utrol grade), pH 7.4. The protein is eluted with buffer containing 250 mM imidazole. Fractions containing the desired protein are pooled and stored at 4°C. Protein concentration is estimated by its absorbance at 280 nm using the calculated extinction coefficient based on its amino acid sequence.

The proteins are refolded by diluting the sample slowly into freshly prepared refolding buffer consisting of: 20 mM Tris, pH 8.6, 0.3 M NaCl, 2.5 M urea, 5 mM cysteine, 20 mM glycine and 1 mM EDTA. Refolding volumes are chosen so that the final protein concentration is between 50 to 100 micrograms/ml. The refolding solution is stirred gently at 4°C for 12-36 hours. The refolding reaction is quenched by the addition of TFA to a final concentration of 0.4% (pH of approximately 3). Before further purification of the protein, the solution is filtered through a 0.22 micron filter and acetonitrile is added to 2-10% final concentration. The refolded protein is chromatographed on a Poros R1/H reversed phase column using a mobile buffer of 0.1% TFA with elution with a gradient of acetonitrile from 10 to 80%. Aliquots of fractions with A₂₈₀ absorbance are analyzed on SDS polyacrylamide gels and fractions containing homogeneous refolded protein are pooled. Generally, the properly refolded species of most proteins are eluted at the lowest concentrations of acetonitrile since those species are the most compact with their hydrophobic interiors shielded from interaction with the reversed phase resin. Aggregated species are usually eluted at higher acetonitrile concentrations. In addition to resolving misfolded forms of proteins from the desired form, the reversed phase step also removes endotoxin from the samples.

Fractions containing the desired folded PRO polypeptide are pooled and the acetonitrile removed using a gentle stream of nitrogen directed at the solution. Proteins are formulated into 20 mM Hepes, pH 6.8 with 0.14 M sodium chloride and 4% mannitol by dialysis or by gel filtration using G25 Superfine (Pharmacia) resins equilibrated in the formulation buffer and sterile filtered.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

6.8. EXAMPLE 8: Expression of PRO in mammalian cells

This example illustrates preparation of a potentially glycosylated form of PRO by recombinant expression in mammalian cells.

The vector, pRK5 (see EP 307,247, published March 15, 1989), is employed as the expression vector. Optionally, the PRO DNA is ligated into pRK5 with selected restriction enzymes to allow insertion of the PRO DNA using ligation methods such as described in Sambrook *et al.*, *supra*. The resulting vector is called pRK5-PRO.

In one embodiment, the selected host cells may be 293 cells. Human 293 cells (ATCC CCL 1573) are grown to confluence in tissue culture plates in medium such as DMEM supplemented with fetal calf serum and optionally, nutrient components and/or antibiotics. About 10 µg pRK5-PRO DNA is mixed with about 1 µg DNA encoding the VA RNA gene [Thimmappaya *et al.*, *Cell*, 31:543 (1982)] and dissolved in 500 µl of 1 mM Tris-HCl, 0.1 mM EDTA, 0.227 M CaCl₂. To this mixture is added, dropwise, 500 µl of 50 mM HEPES (pH 7.35), 280 mM NaCl, 1.5 mM NaPO₄, and a precipitate is allowed to form for 10 minutes at 25°C. The precipitate is suspended and added to the 293 cells and allowed to settle for about four hours at 37°C. The culture medium is aspirated off and 2 ml of 20% glycerol in PBS is added for 30 seconds. The 293 cells are then washed with serum free medium, fresh medium is added and the cells are incubated for about 5 days.

Approximately 24 hours after the transfections, the culture medium is removed and replaced with culture medium (alone) or culture medium containing 200 µCi/ml ³⁵S-cysteine and 200 µCi/ml ³⁵S-methionine. After a 12 hour incubation, the conditioned medium is collected, concentrated on a spin filter, and loaded onto a 15% SDS gel. The processed gel may be dried and exposed to film for a selected period of time to reveal the presence of the PRO polypeptide. The cultures containing transfected cells may undergo further incubation (in serum free medium) and the medium is tested in selected bioassays.

In an alternative technique, PRO may be introduced into 293 cells transiently using the dextran sulfate method described by Sompariyac *et al.*, *Proc. Natl. Acad. Sci.*, 12:7575 (1981). 293 cells are grown to maximal density in a spinner flask and 700 µg pRK5-PRO DNA is added. The cells are first concentrated from the spinner flask by centrifugation and washed with PBS. The DNA-dextran precipitate is incubated on the cell pellet for four hours. The cells are treated with 20% glycerol for 90 seconds, washed with tissue culture medium, and re-introduced into the spinner flask containing tissue culture medium, 5 µg/ml bovine insulin and 0.1 µg/ml bovine transferrin. After about four days, the conditioned media is centrifuged and filtered to remove cells and debris. The sample containing expressed PRO can then be concentrated and purified by any selected method, such as dialysis and/or column chromatography.

In another embodiment, PRO can be expressed in CHO cells. The pRK5-PRO can be transfected into CHO cells using known reagents such as CaPO₄ or DEAE-dextran. As described above, the cell cultures can be incubated, and the medium replaced with culture medium (alone) or medium containing a radiolabel such as ³⁵S-methionine. After determining the presence of a PRO polypeptide, the culture medium may be replaced with serum free medium. Preferably, the cultures are incubated for about 6 days, and then the conditioned medium is harvested.

The medium containing the expressed PRO polypeptide can then be concentrated and purified by any selected method.

Epitope-tagged PRO may also be expressed in host CHO cells. The PRO may be subcloned out of the pRK5 vector. The subclone insert can undergo PCR to fuse in frame with a selected epitope tag such as a poly-His tag into a Baculovirus expression vector. The poly-His tagged PRO insert can then be subcloned into a SV40 driven vector containing a selection marker such as DHFR for selection of stable clones. Finally, the CHO cells can be transfected (as described above) with the SV40 driven vector. Labeling may be performed, as described above, to verify expression. The culture medium containing the expressed poly-His tagged PRO can then be concentrated and purified by any selected method, such as by Ni²⁺-chelate affinity chromatography.

PRO may also be expressed in CHO and/or COS cells by a transient expression procedure or in CHO cells by another stable expression procedure.

Stable expression in CHO cells is performed using the following procedure. The proteins are expressed as an IgG construct (immunoadhesin), in which the coding sequences for the soluble forms (*e.g.*, extracellular domains) of the respective proteins are fused to an IgG1 constant region sequence containing the hinge, CH2 and CH2 domains and/or as a poly-His tagged form.

Following PCR amplification, the respective DNAs are subcloned in a CHO expression vector using standard techniques as described in Ausubel *et al.*, Current Protocols of Molecular Biology, Unit 3.16, John Wiley and Sons (1997). CHO expression vectors are constructed to have compatible restriction sites 5' and 3' of the DNA of interest to allow the convenient shuttling of cDNA's. The vector used in expression in CHO cells is as described in Lucas *et al.*, Nucl. Acids Res., 24:9 (1774-1779 (1996), and uses the SV40 early promoter/enhancer to drive expression of the cDNA of interest and dihydrofolate reductase (DHFR). DHFR expression permits selection for stable maintenance of the plasmid following transfection.

Twelve micrograms of the desired plasmid DNA is introduced into approximately 10 million CHO cells using commercially available transfection reagents Superfect® (Qiagen), Dosper® or Fugene® (Boehringer Mannheim). The cells are grown as described in Lucas *et al.*, *supra*. Approximately 3 x 10⁷ cells are frozen in an ampule for further growth and production as described below.

The ampules containing the plasmid DNA are thawed by placement into a water bath and mixed by vortexing. The contents are pipetted into a centrifuge tube containing 10 ml of media and centrifuged at 1000 rpm for 5 minutes. The supernatant is aspirated and the cells are resuspended in 10 ml of selective media (0.2 µm filtered PS20 with 5% 0.2 µm diafiltered fetal bovine serum). The cells are then aliquoted into a 100 ml spinner containing 90 ml of selective media. After 1-2 days, the cells are transferred into a 250 ml spinner filled with 150 ml selective growth medium and incubated at 37°C. After another 2-3 days, 250 ml, 500 ml and 2000 ml spinners are seeded with 3 x 10⁵ cells/ml. The cell media is exchanged with fresh media by centrifugation and resuspension in production medium. Although any suitable CHO media may be employed, a production medium described in U.S. Patent No. 5,122,469, issued June 16, 1992 may actually be used. A 3L production spinner is seeded at 1.2 x 10⁶ cells/ml. On day 0, the cell number and pH is determined. On day 1, the spinner is sampled and sparging with filtered air is commenced. On day 2, the spinner is sampled, the temperature shifted to 33°C, and 30 ml of 500 g/L

glucose and 0.6 ml of 10% antifoam (e.g., 35% polydimethylsiloxane emulsion, Dow Corning 365 Medical Grade Emulsion) taken. Throughout the production, the pH is adjusted as necessary to keep it at around 7.2. After 10 days, or until the viability drops below 70%, the cell culture is harvested by centrifugation and filtering through a 0.22 μ m filter. The filtrate is either stored at 4°C or immediately loaded onto columns for purification.

5 For the poly-His tagged constructs, the proteins are purified using a Ni²⁺-NTA column (Qiagen). Before purification, imidazole is added to the conditioned media to a concentration of 5 mM. The conditioned media is pumped onto a 6 ml Ni²⁺-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 4°C. After loading, the column is washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly
10 purified protein is subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80°C.

Immunoadhesin (Fc-containing) constructs are purified from the conditioned media as follows. The conditioned medium is pumped onto a 5 ml Protein A column (Pharmacia) which has been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column is washed extensively with equilibration buffer before
15 elution with 100 mM citric acid, pH 3.5. The eluted protein is immediately neutralized by collecting 1 ml fractions into tubes containing 275 μ l of 1 M Tris buffer, pH 9. The highly purified protein is subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity is assessed by SDS polyacrylamide gels and by N-terminal amino acid sequencing by Edman degradation.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

20 6.9. EXAMPLE 9: Expression of PRO in Yeast

The following method describes recombinant expression of PRO in yeast.

First, yeast expression vectors are constructed for intracellular production or secretion of PRO from the ADH2/GAPDH promoter. DNA encoding PRO and the promoter is inserted into suitable restriction enzyme sites in the selected plasmid to direct intracellular expression of PRO. For secretion, DNA encoding PRO can be cloned
25 into the selected plasmid, together with DNA encoding the ADH2/GAPDH promoter, a native PRO signal peptide or other mammalian signal peptide, or, for example, a yeast alpha-factor or invertase secretory signal/leader sequence, and linker sequences (if needed) for expression of PRO.

Yeast cells, such as yeast strain AB110, can then be transformed with the expression plasmids described above and cultured in selected fermentation media. The transformed yeast supernatants can be analyzed by
30 precipitation with 10% trichloroacetic acid and separation by SDS-PAGE, followed by staining of the gels with Coomassie Blue stain.

Recombinant PRO can subsequently be isolated and purified by removing the yeast cells from the fermentation medium by centrifugation and then concentrating the medium using selected cartridge filters. The concentrate containing PRO may further be purified using selected column chromatography resins.

35 Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

6.10. EXAMPLE 10: Expression of PRO in Baculovirus-Infected Insect Cells

The following method describes recombinant expression in Baculovirus-infected insect cells.

The sequence coding for PRO is fused upstream of an epitope tag contained within a baculovirus expression vector. Such epitope tags include poly-His tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pVL1393 (Novagen). Briefly, the sequence encoding PRO or the desired portion of the coding sequence of PRO (such as the sequence encoding the extracellular domain of a transmembrane protein or the sequence encoding the mature protein if the protein is extracellular) is amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector.

Recombinant baculovirus is generated by co-transfecting the above plasmid and BaculoGold™ virus DNA (Pharmingen) into *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711) using lipofectin (commercially available from GIBCO-BRL). After 4 - 5 days of incubation at 28°C, the released viruses are harvested and used for further amplifications. Viral infection and protein expression are performed as described by O'Reilley *et al.*, Baculovirus expression vectors: A Laboratory Manual, Oxford: Oxford University Press (1994).

Expressed poly-His tagged PRO can then be purified, for example, by Ni²⁺-chelate affinity chromatography as follows. Extracts are prepared from recombinant virus-infected Sf9 cells as described by Rupert *et al.*, Nature, 362:175-179 (1993). Briefly, Sf9 cells are washed, resuspended in sonication buffer (25 ml Hepes, pH 7.9; 12.5 mM MgCl₂; 0.1 mM EDTA; 10% glycerol; 0.1% NP-40; 0.4 M KCl), and sonicated twice for 20 seconds on ice. The sonicates are cleared by centrifugation, and the supernatant is diluted 50-fold in loading buffer (50 mM phosphate, 300 mM NaCl, 10% glycerol, pH 7.8) and filtered through a 0.45 µm filter. A Ni²⁺-NTA agarose column (commercially available from Qiagen) is prepared with a bed volume of 5 ml, washed with 25 ml of water and equilibrated with 25 ml of loading buffer. The filtered cell extract is loaded onto the column at 0.5 ml per minute. The column is washed to baseline A₂₈₀ with loading buffer, at which point fraction collection is started. Next, the column is washed with a secondary wash buffer (50 mM phosphate; 300 mM NaCl, 10% glycerol, pH 6.0), which elutes nonspecifically bound protein. After reaching A₂₈₀ baseline again, the column is developed with a 0 to 500 mM imidazole gradient in the secondary wash buffer. One ml fractions are collected and analyzed by SDS-PAGE and silver staining or Western blot with Ni²⁺-NTA-conjugated to alkaline phosphatase (Qiagen). Fractions containing the eluted His₁₀-tagged PRO are pooled and dialyzed against loading buffer.

Alternatively, purification of the IgG tagged (or Fc tagged) PRO can be performed using known chromatography techniques, including for instance, Protein A or protein G column chromatography.

Following PCR amplification, the respective coding sequences are subcloned into a baculovirus expression vector (pb.PH.IgG for IgG fusions and pb.PH.His.c for poly-His tagged proteins), and the vector and Baculogold® baculovirus DNA (Pharmingen) are co-transfected into 105 *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711), using Lipofectin (Gibco BRL). pb.PH.IgG and pb.PH.His are modifications of the commercially available baculovirus expression vector pVL1393 (Pharmingen), with modified polylinker regions to include the His or Fc tag sequences. The cells are grown in Hink's TNM-FH medium supplemented with 10% FBS (Hyclone). Cells are

incubated for 5 days at 28°C. The supernatant is harvested and subsequently used for the first viral amplification by infecting Sf9 cells in Hink's TNM-FH medium supplemented with 10% FBS at an approximate multiplicity of infection (MOI) of 10. Cells are incubated for 3 days at 28°C. The supernatant is harvested and the expression of the constructs in the baculovirus expression vector is determined by batch binding of 1 ml of supernatant to 25 ml of Ni²⁺-NTA beads (QIAGEN) for histidine tagged proteins or Protein-A Sepharose CL-4B beads (Pharmacia) for IgG tagged proteins followed by SDS-PAGE analysis comparing to a known concentration of protein standard by Coomassie blue staining.

The first viral amplification supernatant is used to infect a spinner culture (500 ml) of Sf9 cells grown in ESF-921 medium (Expression Systems LLC) at an approximate MOI of 0.1. Cells are incubated for 3 days at 28°C. The supernatant is harvested and filtered. Batch binding and SDS-PAGE analysis is repeated, as necessary, until expression of the spinner culture is confirmed.

The conditioned medium from the transfected cells (0.5 to 3 L) is harvested by centrifugation to remove the cells and filtered through 0.22 micron filters. For the poly-His tagged constructs, the protein construct is purified using a Ni²⁺-NTA column (Qiagen). Before purification, imidazole is added to the conditioned media to a concentration of 5 mM. The conditioned media is pumped onto a 6 ml Ni²⁺-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 4°C. After loading, the column is washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein is subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80°C.

Immunoadhesin (Fc containing) constructs of proteins are purified from the conditioned media as follows. The conditioned media is pumped onto a 5 ml Protein A column (Pharmacia) which has been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column is washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein is immediately neutralized by collecting 1 ml fractions into tubes containing 275 ml of 1 M Tris buffer, pH 9. The highly purified protein is subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity of the proteins is verified by SDS polyacrylamide gel (PEG) electrophoresis and N-terminal amino acid sequencing by Edman degradation.

Alternatively, a modified baculovirus procedure may be used incorporating high-5 cells. In this procedure, the DNA encoding the desired sequence is amplified with suitable systems, such as Pfu (Stratagene), or fused upstream (5'-of) of an epitope tag contained with a baculovirus expression vector. Such epitope tags include poly-His tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pIE1-1 (Novagen). The pIE1-1 and pIE1-2 vectors are designed for constitutive expression of recombinant proteins from the baculovirus ie1 promoter in stably-transformed insect cells (1). The plasmids differ only in the orientation of the multiple cloning sites and contain all promoter sequences known to be important for ie1-mediated gene expression in uninfected insect cells as well as the hr5 enhancer element. pIE1-1 and pIE1-2 include the translation initiation site and can be used to produce fusion proteins. Briefly, the desired sequence or the desired portion of the sequence (such as the sequence encoding the

extracellular domain of a transmembrane protein) is amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector. For example, derivatives of pIE1-1 can include the Fc region of human IgG (pb.PH.IgG) or an 8 histidine (pb.PH.His) tag downstream (3'-of) the desired sequence. Preferably, the vector construct is sequenced for confirmation.

High-5 cells are grown to a confluency of 50% under the conditions of, 27°C, no CO₂, NO pen/strep. For each 150 mm plate, 30 µg of pIE based vector containing the sequence is mixed with 1 ml Ex-Cell medium (Media: Ex-Cell 401 + 1/100 L-Glu JRH Biosciences #14401-78P (note: this media is light sensitive)), and in a separate tube, 100 µl of CellFectin (CellFECTIN (GibcoBRL #10362-010) (vortexed to mix)) is mixed with 1 ml of Ex-Cell medium. The two solutions are combined and allowed to incubate at room temperature for 15 minutes. 8 ml of Ex-Cell media is added to the 2 ml of DNA/CellFECTIN mix and this is layered on high-5 cells that have been washed once with Ex-Cell media. The plate is then incubated in darkness for 1 hour at room temperature. The DNA/CellFECTIN mix is then aspirated, and the cells are washed once with Ex-Cell to remove excess CellFECTIN, 30 ml of fresh Ex-Cell media is added and the cells are incubated for 3 days at 28°C. The supernatant is harvested and the expression of the sequence in the baculovirus expression vector is determined by batch binding of 1 ml of supernatant to 25 ml of Ni²⁺-NTA beads (QIAGEN) for histidine tagged proteins or Protein-A Sepharose CL-4B beads (Pharmacia) for IgG tagged proteins followed by SDS-PAGE analysis comparing to a known concentration of protein standard by Coomassie blue staining.

The conditioned media from the transfected cells (0.5 to 3 L) is harvested by centrifugation to remove the cells and filtered through 0.22 micron filters. For the poly-His tagged constructs, the protein comprising the sequence is purified using a Ni²⁺-NTA column (Qiagen). Before purification, imidazole is added to the conditioned media to a concentration of 5 mM. The conditioned media is pumped onto a 6 ml Ni²⁺-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 48°C. After loading, the column is washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein is then subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80°C.

Immunoadhesin (Fc containing) constructs of proteins are purified from the conditioned media as follows. The conditioned media is pumped onto a 5 ml Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column is washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein is immediately neutralized by collecting 1 ml fractions into tubes containing 275 ml of 1 M Tris buffer, pH 9. The highly purified protein is subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity of the sequence is assessed by SDS polyacrylamide gels and by N-terminal amino acid sequencing by Edman degradation and other analytical procedures as desired or necessary.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

6.11. EXAMPLE 11: Preparation of Antibodies that Bind PRO

This example illustrates preparation of monoclonal antibodies which can specifically bind the PRO polypeptide or an epitope on the PRO polypeptide without substantially binding to any other polypeptide or polypeptide epitope.

5 Techniques for producing the monoclonal antibodies are known in the art and are described, for instance, in Goding, *supra*. Immunogens that may be employed include purified PRO, fusion proteins containing PRO, and cells expressing recombinant PRO on the cell surface. Selection of the immunogen can be made by the skilled artisan without undue experimentation.

10 Mice, such as Balb/c, are immunized with the PRO immunogen emulsified in complete Freund's adjuvant and injected subcutaneously or intraperitoneally in an amount from 1-100 micrograms. Alternatively, the immunogen is emulsified in MPL-TDM adjuvant (Ribi Immunochemical Research, Hamilton, MT) and injected into the animal's hind foot pads. The immunized mice are then boosted 10 to 12 days later with additional immunogen emulsified in the selected adjuvant. Thereafter, for several weeks, the mice may also be boosted with additional immunization injections. Serum samples may be periodically obtained from the mice by retro-orbital
15 bleeding for testing in ELISA assays to detect anti-PRO antibodies.

After a suitable antibody titer has been detected, the animals "positive" for antibodies can be injected with a final intravenous injection of PRO. Three to four days later, the mice are sacrificed and the spleen cells are harvested. The spleen cells are then fused (using 35% polyethylene glycol) to a selected murine myeloma cell line such as P3X63AgU.1, available from ATCC, No. CRL 1597. The fusions generate hybridoma cells which can then
20 be plated in 96 well tissue culture plates containing HAT (hypoxanthine, aminopterin, and thymidine) medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

The hybridoma cells will be screened in an ELISA for reactivity against PRO. Determination of "positive" hybridoma cells secreting the desired monoclonal antibodies against PRO is within the skill in the art.

25 The positive hybridoma cells can be injected intraperitoneally into syngeneic Balb/c mice to produce ascites containing the anti-PRO monoclonal antibodies. Alternatively, the hybridoma cells can be grown in tissue culture flasks or roller bottles. Purification of the monoclonal antibodies produced in the ascites can be accomplished using ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein G can be employed.

6.12. EXAMPLE 12: Purification of PRO Polypeptides Using Specific Antibodies

30 Native or recombinant PRO polypeptides may be purified by a variety of standard techniques in the art of protein purification. For example, pro-PRO polypeptide, mature PRO polypeptide, or pre-PRO polypeptide is purified by immunoaffinity chromatography using antibodies specific for the PRO polypeptide of interest. In general, an immunoaffinity column is constructed by covalently coupling the anti-PRO polypeptide antibody to an activated chromatographic resin.

35 Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway, N.J.). Likewise,

monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated SEPHAROSE™ (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

Such an immunoaffinity column is utilized in the purification of PRO polypeptide by preparing a fraction from cells containing PRO polypeptide in a soluble form. This preparation is derived by solubilization of the whole cell or of a subcellular fraction obtained via differential centrifugation by the addition of detergent or by other methods well known in the art. Alternatively, soluble PRO polypeptide containing a signal sequence may be secreted in useful quantity into the medium in which the cells are grown.

A soluble PRO polypeptide-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PRO polypeptide (*e.g.*, high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/PRO polypeptide binding (*e.g.*, a low pH buffer such as approximately pH 2-3, or a high concentration of a chaotrope such as urea or thiocyanate ion), and PRO polypeptide is collected.

6.13. EXAMPLE 13: Drug Screening

This invention is particularly useful for screening compounds by using PRO polypeptides or binding fragment thereof in any of a variety of drug screening techniques. The PRO polypeptide or fragment employed in such a test may either be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the PRO polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may measure, for example, the formation of complexes between PRO polypeptide or a fragment and the agent being tested. Alternatively, one can examine the diminution in complex formation between the PRO polypeptide and its target cell or target receptors caused by the agent being tested.

Thus, the present invention provides methods of screening for drugs or any other agents which can affect a PRO polypeptide-associated disease or disorder. These methods comprise contacting such an agent with an PRO polypeptide or fragment thereof and assaying (i) for the presence of a complex between the agent and the PRO polypeptide or fragment, or (ii) for the presence of a complex between the PRO polypeptide or fragment and the cell, by methods well known in the art. In such competitive binding assays, the PRO polypeptide or fragment is typically labeled. After suitable incubation, free PRO polypeptide or fragment is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of the particular agent to bind to PRO polypeptide or to interfere with the PRO polypeptide/cell complex.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to a polypeptide and is described in detail in WO 84/03564, published on September 13, 1984. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such

as plastic pins or some other surface. As applied to a PRO polypeptide, the peptide test compounds are reacted with PRO polypeptide and washed. Bound PRO polypeptide is detected by methods well known in the art. Purified PRO polypeptide can also be coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies can be used to capture the peptide and immobilize it on the solid support.

This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding PRO polypeptide specifically compete with a test compound for binding to PRO polypeptide or fragments thereof. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PRO polypeptide.

6.14. EXAMPLE 14: Rational Drug Design

The goal of rational drug design is to produce structural analogs of biologically active polypeptide of interest (*i.e.*, a PRO polypeptide) or of small molecules with which they interact, *e.g.*, agonists, antagonists, or inhibitors. Any of these examples can be used to fashion drugs which are more active or stable forms of the PRO polypeptide or which enhance or interfere with the function of the PRO polypeptide *in vivo* (*c.f.*, Hodgson, Bio/Technology, 9: 19-21 (1991)).

In one approach, the three-dimensional structure of the PRO polypeptide, or of an PRO polypeptide-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the PRO polypeptide must be ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of the PRO polypeptide may be gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design analogous PRO polypeptide-like molecules or to identify efficient inhibitors. Useful examples of rational drug design may include molecules which have improved activity or stability as shown by Braxton and Wells, Biochemistry, 31:7796-7801 (1992) or which act as inhibitors, agonists, or antagonists of native peptides as shown by Athauda *et al.*, J. Biochem., 113:742-746 (1993).

It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon which subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids would be expected to be an analog of the original receptor. The anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides would then act as the pharmacore.

By virtue of the present invention, sufficient amounts of the PRO polypeptide may be made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the PRO polypeptide amino acid sequence provided herein will provide guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

6.15. EXAMPLE 15: Stimulation of Endothelial Cell Proliferation (Assay 8)

This assay is designed to determine whether PRO polypeptides of the present invention show the ability to stimulate adrenal cortical capillary endothelial cell (ACE) growth. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of conditions or disorders where angiogenesis would be beneficial including, for example, wound healing, and the like (as would agonists of these PRO polypeptides). Antagonists of the PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of cancerous tumors.

Bovine adrenal cortical capillary endothelial (ACE) cells (from primary culture, maximum of 12-14 passages) were plated in 96-well plates at 500 cells/well per 100 microliter. Assay media included low glucose DMEM, 10% calf serum, 2 mM glutamine, and 1X penicillin/streptomycin/fungizone. Control wells included the following: (1) no ACE cells added; (2) ACE cells alone; (3) ACE cells plus VEGF (5 ng/ml); and (4) ACE cells plus FGF (5ng/ml). The control or test sample, (in 100 microliter volumes), was then added to the wells (at dilutions of 1%, 0.1% and 0.01%, respectively). The cell cultures were incubated for 6-7 days at 37°C/5% CO₂. After the incubation, the media in the wells was aspirated, and the cells were washed 1X with PBS. An acid phosphatase reaction mixture (100 microliter; 0.1M sodium acetate, pH 5.5, 0.1% Triton X-100, 10 mM p-nitrophenyl phosphate) was then added to each well. After a 2 hour incubation at 37°C, the reaction was stopped by addition of 10 microliters 1N NaOH. Optical density (OD) was measured on a microplate reader at 405 nm.

The activity of a PRO polypeptide was calculated as the fold increase in proliferation (as determined by the acid phosphatase activity, OD 405 nm) relative to (1) cell only background, and (2) relative to maximum stimulation by VEGF. VEGF (at 3-10 ng/ml) and FGF (at 1-5 ng/ml) were employed as an activity reference for maximum stimulation. Results of the assay were considered "positive" if the observed stimulation was \geq 50% increase over background. VEGF (5 ng/ml) control at 1% dilution gave 1.24 fold stimulation; FGF (5 ng/ml) control at 1% dilution gave 1.46 fold stimulation.

PRO21 tested positive in this assay.

6.16. EXAMPLE 16: Inhibition of Vascular Endothelial Growth Factor (VEGF) Stimulated Proliferation of Endothelial Cell Growth (Assay 9)

The ability of various PRO polypeptides to inhibit VEGF stimulated proliferation of endothelial cells was tested. Polypeptides testing positive in this assay are useful for inhibiting endothelial cell growth in mammals where such an effect would be beneficial, e.g., for inhibiting tumor growth.

Specifically, bovine adrenal cortical capillary endothelial cells (ACE) (from primary culture, maximum of 12-14 passages) were plated in 96-well plates at 500 cells/well per 100 microliter. Assay media included low glucose DMEM, 10% calf serum, 2 mM glutamine, and 1X penicillin/streptomycin/fungizone. Control wells included the following: (1) no ACE cells added; (2) ACE cells alone; (3) ACE cells plus 5 ng/ml FGF; (4) ACE cells plus 3 ng/ml VEGF; (5) ACE cells plus 3 ng/ml VEGF plus 1 ng/ml TGF-beta; and (6) ACE cells plus 3 ng/ml VEGF plus 5 ng/ml LIF. The test samples, poly-his tagged PRO polypeptides (in 100 microliter volumes), were then added to the wells (at dilutions of 1%, 0.1% and 0.01%, respectively). The cell cultures were

incubated for 6-7 days at 37°C/5% CO₂. After the incubation, the media in the wells was aspirated, and the cells were washed 1X with PBS. An acid phosphatase reaction mixture (100 microliter; 0.1M sodium acetate, pH 5.5, 0.1% Triton X-100, 10 mM p-nitrophenyl phosphate) was then added to each well. After a 2 hour incubation at 37°C, the reaction was stopped by addition of 10 microliters 1N NaOH. Optical density (OD) was measured on a microplate reader at 405 nm.

The activity of PRO polypeptides was calculated as the percent inhibition of VEGF (3 ng/ml) stimulated proliferation (as determined by measuring acid phosphatase activity at OD 405 nm) relative to the cells without stimulation. TGF-beta was employed as an activity reference at 1 ng/ml, since TGF-beta blocks 70-90% of VEGF-stimulated ACE cell proliferation. The results are indicative of the utility of the PRO polypeptides in cancer therapy and specifically in inhibiting tumor angiogenesis. Numerical values (relative inhibition) are determined by calculating the percent inhibition of VEGF stimulated proliferation by the PRO polypeptides relative to cells without stimulation and then dividing that percentage into the percent inhibition obtained by TGF-β at 1 ng/ml which is known to block 70-90% of VEGF stimulated cell proliferation. The results are considered positive if the PRO polypeptide exhibits 30% or greater inhibition of VEGF stimulation of endothelial cell growth (relative inhibition 30% or greater).

PRO247, PRO720 and PRO4302 tested positive in this assay.

6.17. EXAMPLE 17: Enhancement of Heart Neonatal Hypertrophy Induced by LIF+ET-1 (Assay 75)

This assay is designed to determine whether PRO polypeptides of the present invention show the ability to enhance neonatal heart hypertrophy induced by LIF and endothelin-1 (ET-1). A test compound that provides a positive response in the present assay would be useful for the therapeutic treatment of cardiac insufficiency diseases or disorders characterized or associated with an undesired level of hypertrophy of the cardiac muscle.

Cardiac myocytes from 1-day old Harlan Sprague Dawley rats (180 μl at 7.5 x 10⁴/ml, serum <0.1, freshly isolated) are introduced on day 1 to 96-well plates previously coated with DMEM/F12 + 4%FCS. Test PRO polypeptide samples or growth medium alone (negative control) are then added directly to the wells on day 2 in 20 μl volume. LIF + ET-1 are then added to the wells on day 3. The cells are stained after an additional 2 days in culture and are then scored visually the next day. A positive in the assay occurs when the PRO polypeptide treated myocytes obtain a score greater than zero. A score of zero represents non-responsive cells whereas scores of 1 or 2 represent enhancement (*i.e.* they are visually larger on the average or more numerous than the untreated myocytes).

PRO21 polypeptides tested positive in this assay.

6.18. EXAMPLE 18: Detection of Endothelial Cell Apoptosis (FACS) (Assay 96)

The ability of PRO polypeptides of the present invention to induce apoptosis in endothelial cells was tested in human venous umbilical vein endothelial cells (HUVEC, Cell Systems) in gelatinized T175 flasks using HUVEC cells below passage 10. PRO polypeptides testing positive in this assay are expected to be useful for

therapeutically treating conditions where apoptosis of endothelial cells would be beneficial including, for example, the therapeutic treatment of tumors.

On day one, the cells were split [420,000 cells per gelatinized 6 cm dishes - (11 x 10³ cells/cm² Falcon, Primaria)] and grown in media containing serum (CS-C, Cell System) overnight or for 16 hours to 24 hours.

On day 2, the cells were washed 1x with 5 ml PBS ; 3 ml of 0% serum medium was added with VEGF (100 ng/ml); and 30 μ l of the PRO test compound (final dilution 1%) or 0% serum medium (negative control) was added. The mixtures were incubated for 48 hours before harvesting.

The cells were then harvested for FACS analysis. The medium was aspirated and the cells washed once with PBS. 5 ml of 1 x trypsin was added to the cells in a T-175 flask, and the cells were allowed to stand until they were released from the plate (about 5-10 minutes). Trypsinization was stopped by adding 5 ml of growth media. The cells were spun at 1000 rpm for 5 minutes at 4°C. The media was aspirated and the cells were resuspended in 10 ml of 10% serum complemented medium (Cell Systems), 5 μ l of Annexin-FITC (BioVison) added and chilled tubes were submitted for FACS. A positive result was determined to be enhanced apoptosis in the PRO polypeptide treated samples as compared to the negative control.

PRO4302 polypeptide tested positive in this assay.

6.19. EXAMPLE 19: Induction of c-fos in HUVEC Cells (Assay 123)

This assay is designed to determine whether PRO polypeptides show the ability to induce c-fos in HUVEC cells. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of conditions or disorders where angiogenesis would be beneficial including, for example, wound healing, and the like (as would agonists of these PRO polypeptides). Antagonists of the PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of cancerous tumors.

Human venous umbilical vein endothelial cells (HUVEC, Cell Systems) in growth media (50% Ham's F12 w/o GHT: low glucose, and 50% DMEM without glycine: with NaHCO₃, 1% glutamine, 10 mM HEPES, 10% FBS, 10 ng/ml bFGF) were plated on 96-well microtiter plates at a cell density of 5x10³ cells/well. The day after plating (day 2), the cells were starved for 24 hours by removing the growth media and replacing with serum free media. On day 3, the cells are treated with 100 μ l/well test samples and controls (positive control = growth media; negative control = Protein 32 buffer = 10 mM HEPES, 140 mM NaCl, 4% (w/v) mannitol, pH 6.8). One plate of cells was incubated for 30 minutes at 37°C, in 5% CO₂. Another plate of cells was incubated for 60 minutes at 37°C, in 5% CO₂. The samples were removed, and RNA was harvested using the RNeasy 96 kit (Qiagen). Next, the RNA was assayed for c-fos, egr-1 and GAPDH induction using Taqman.

The measure of activity of the fold increase over the negative control (Protein 32/HEPES buffer described above) value was by obtained by calculating the fold increase of the ratio of c-fos to GAPDH in test samples as compared to the negative control. The results are considered positive if the PRO polypeptide exhibits at least a two-fold value over the negative buffer control.

PRO1376 polypeptide tested positive in this assay.

6.20. EXAMPLE 20: Normal Human Iliac Artery Endothelial Cell Proliferation (Assay 138)

This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce proliferation of human iliac artery endothelial cells in culture and, therefore, function as useful growth factors.

On day 0, human iliac artery endothelial cells (from cell lines, maximum of 12-14 passages) were plated in 96-well plates at 1000 cells/well per 100 microliter and incubated overnight in complete media [epithelial cell growth media (EGM, Clonetics), plus supplements: human epithelial growth factor (hEGF), bovine brain extract (BBE), hydrocortisone, GA-1000, and fetal bovine serum (FBS, Clonetics)]. On day 1, complete media was replaced by basal media [EGM plus 1% FBS] and addition of PRO polypeptides at 1%, 0.1% and 0.01%. On day 7, an assessment of cell proliferation was performed by Alamar Blue assay followed by Crystal Violet. Results are expressed as % of the cell growth observed with control buffer.

The following PRO polypeptides tested positive in this assay: PRO214, PRO238, PRO256, PRO363, PRO365, PRO791, PRO836, PRO1025, PRO1029, PRO1186, PRO1192, PRO1272, PRO1274, PRO1279, PRO1306, PRO1325, PRO1329, PRO1376, PRO1411, PRO1419, PRO1508, PRO1787, PRO1868, PRO1890, PRO4324, PRO4333, PRO4408, PRO4499, PRO5725, PRO6006, PRO9821, PRO9873, PRO10008, PRO10096, PRO19670, PRO20040, PRO20044, PRO21384 and PRO28631.

6.21. EXAMPLE 21: Pooled Human Umbilical Vein Endothelial Cell Proliferation (Assay 139)

This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce proliferation of pooled human umbilical vein endothelial cells in culture and, therefore, function as useful growth factors.

On day 0, pooled human umbilical vein endothelial cells (from cell lines, maximum of 12-14 passages) were plated in 96-well plates at 1000 cells/well per 100 microliter and incubated overnight in complete media [epithelial cell growth media (EGM, Clonetics), plus supplements: human epithelial growth factor (hEGF), bovine brain extract (BBE), hydrocortisone, GA-1000, and fetal bovine serum (FBS, Clonetics)]. On day 1, complete media was replaced by basal media [EGM plus 1% FBS] and addition of PRO polypeptides at 1%, 0.1% and 0.01%. On day 7, an assessment of cell proliferation was performed by Alamar Blue assay followed by Crystal Violet. Results are expressed as % of the cell growth observed with control buffer.

The following PRO polypeptides tested positive in this assay: PRO181, PRO205, PRO221, PRO229, PRO231, PRO238, PRO241, PRO247, PRO256, PRO258, PRO263, PRO265, PRO295, PRO321, PRO322, PRO337, PRO363, PRO444, PRO533, PRO697, PRO725, PRO771, PRO788, PRO819, PRO827, PRO828, PRO846, PRO865, PRO1005, PRO1006, PRO1007, PRO1025, PRO1054, PRO1071, PRO1075, PRO1079, PRO1080, PRO1114, PRO1131, PRO1155, PRO1160, PRO1184, PRO1190, PRO1192, PRO1195, PRO1244, PRO1272, PRO1273, PRO1279, PRO1283, PRO1286, PRO1306, PRO1309, PRO1325, PRO1329, PRO1347, PRO1356, PRO1376, PRO1382, PRO1412, PRO1419, PRO1474, PRO1477, PRO1488, PRO1550, PRO1556, PRO1760, PRO1782, PRO1787, PRO1801, PRO1868, PRO1887, PRO3438, PRO3444, PRO4302, PRO4324, PRO4341, PRO4342, PRO4353, PRO4354, PRO4356, PRO4371, PRO4405, PRO4422, PRO4425, PRO5723,

PRO5725, PRO5737, PRO5776, PRO6029, PRO6071, PRO7436, PRO9771, PRO10008, PRO10096, PRO21055 and PRO21384.

6.22. EXAMPLE 22: Human Coronary Artery Smooth Muscle Cell Proliferation (Assay 140)

This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce proliferation of human coronary artery smooth muscle cells in culture and, therefore, function as useful growth factors.

On day 0, human coronary artery smooth muscle cells (from cell lines, maximum of 12-14 passages) were plated in 96-well plates at 1000 cells/well per 100 microliter and incubated overnight in complete media [smooth muscle growth media (SmGM, Clonetics), plus supplements: insulin, human epithelial growth factor (hEGF), human fibroblast growth factor (hFGF), GA-1000, and fetal bovine serum (FBS, Clonetics)]. On day 1, complete media was replaced by basal media [SmGM plus 1 % FBS] and addition of PRO polypeptides at 1 %, 0.1 % and 0.01 %. On day 7, an assessment of cell proliferation was performed by Alamar Blue assay followed by Crystal Violet. Results are expressed as % of the cell growth observed with control buffer.

The following PRO polypeptides tested positive in this assay: PRO162, PRO181, PRO182, PRO195, PRO204, PRO221, PRO230, PRO256, PRO258, PRO533, PRO697, PRO725, PRO738, PRO826, PRO836, PRO840, PRO846, PRO865, PRO982, PRO1025, PRO1029, PRO1071, PRO1080, PRO1083, PRO1134, PRO1160, PRO1182, PRO1184, PRO1186, PRO1192, PRO1265, PRO1274, PRO1279, PRO1283, PRO1306, PRO1308, PRO1309, PRO1325, PRO1337, PRO1338, PRO1343, PRO1376, PRO1387, PRO1411, PRO1412, PRO1415, PRO1434, PRO1474, PRO1488, PRO1550, PRO1556, PRO1567, PRO1600, PRO1754, PRO1758, PRO1760, PRO1787, PRO1865, PRO1868, PRO1917, PRO1928, PRO3438, PRO3562, PRO4302, PRO4333, PRO4345, PRO4353, PRO4354, PRO4405, PRO4408, PRO4430, PRO4503, PRO5725, PRO6714, PRO9771, PRO9820, PRO9940, PRO10096, PRO21055, PRO21184 and PRO21366.

6.23. EXAMPLE 23: Microarray Analysis to Detect Overexpression of PRO Polypeptides in HUVEC Cells Treated with Growth Factors

This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce angiogenesis by stimulating endothelial cell tube formation in HUVEC cells.

Nucleic acid microarrays, often containing thousands of gene sequences, are useful for identifying differentially expressed genes in tissues exposed to various stimuli (e.g., growth factors) as compared to their normal, unexposed counterparts. Using nucleic acid microarrays, test and control mRNA samples from test and control tissue samples are reverse transcribed and labeled to generate cDNA probes. The cDNA probes are then hybridized to an array of nucleic acids immobilized on a solid support. The array is configured such that the sequence and position of each member of the array is known. Hybridization of a labeled probe with a particular array member indicates that the sample from which the probe was derived expresses that gene. If the hybridization signal of a probe from a test (exposed tissue) sample is greater than hybridization signal of a probe from a control (normal, unexposed tissue) sample, the gene or genes overexpressed in the exposed tissue are identified. The

implication of this result is that an overexpressed protein in an exposed tissue may be involved in the functional changes within the tissue following exposure to the stimuli (*e.g.*, tube formation).

The methodology of hybridization of nucleic acids and microarray technology is well known in the art. In the present example, the specific preparation of nucleic acids for hybridization and probes, slides, and hybridization conditions are all detailed in U.S. Provisional Patent Application Serial No. 60/193,767, filed on March 31, 2000 and which is herein incorporated by reference.

In the present example, HUVEC cells grown in either collagen gels or fibrin gels were induced to form tubes by the addition of various growth factors. Specifically, collagen gels were prepared as described previously in Yang *et al.*, *American J. Pathology*, 1999, 155(3):887-895 and Xin *et al.*, *American J. Pathology*, 2001, 158(3):1111-1120. Following gelation of the HUVEC cells, 1X basal medium containing M199 supplemented with 1%FBS, 1X ITS, 2 mM L-glutamine, 50 µg/ml ascorbic acid, 26.5 mM NaHCO₃, 100U/ml penicillin and 100 U/ml streptomycin was added. Tube formation was elicited by the inclusion in the culture media of either a mixture of phorbol myrsitate acetate (50 nM), vascular endothelial cell growth factor (40 ng/ml) and basic fibroblast growth factor (40 ng/ml) ("PMA growth factor mix") or hepatocyte growth factor (40 ng/ml) and vascular endothelial cell growth factor (40 ng/ml) (HGF/VEGF mix) for the indicated period of time. Fibrin Gels were prepared by suspending Huvec (4 x 10⁵ cells/ml) in M199 containing 1% fetal bovine serum (Hyclone) and human fibrinogen (2.5mg/ml). Thrombin (50U/ml) was then added to the fibrinogen suspension at a ratio of 1 part thrombin solution:30 parts fibrinogen suspension. The solution was then layered onto 10 cm tissue culture plates (total volume: 15 ml/plate) and allowed to solidify at 37°C for 20 min. Tissue culture media (10 ml of BM containing PMA (50 nM), bFGF (40ng/ml) and VEGF (40 ng/ml)) was then added and the cells incubated at 37°C in 5%CO₂ in air for the indicated period of time.

Total RNA was extracted from the HUVEC cells incubated for 0, 4, 8, 24, 40 and 50 hours in the different matrix and media combinations using a TRIzol extraction followed by a second purification using RNAeasy Mini Kit (Qiagen). The total RNA was used to prepare cRNA which was then hybridized to the microarrays.

In the present experiments, nucleic acid probes derived from the herein described PRO polypeptide-encoding nucleic acid sequences were used in the creation of the microarray and RNA from the HUVEC cells described above were used for the hybridization thereto. Pairwise comparisons were made using time 0 chips as a baseline. Three replicate samples were analyzed for each experimental condition and time. Hence there were 3 time 0 samples for each treatment and 3 replicates of each successive time point. Therefore, a 3 by 3 comparison was performed for each time point compared against each time 0 point. This resulted in 9 comparisons per time point. Only those genes that had increased expression in all three non-time-0 replicates in each of the different matrix and media combinations as compared to any of the three time zero replicates were considered positive. Although this stringent method of data analysis does allow for false negatives, it minimizes false positives.

PRO178, PRO195, PRO228, PRO301, PRO302, PRO532, PRO724, PRO730, PRO734, PRO793, PRO871, PRO938, PRO1012, PRO1120, PRO1139, PRO1198, PRO1287, PRO1361, PRO1864, PRO1873,

PRO2010, PRO3579, PRO4313, PRO4527, PRO4538, PRO4553, PRO4995, PRO5730, PRO6008, PRO7223, PRO7248 and PRO7261 tested positive in this assay.

6.24. EXAMPLE 24: *In situ* Hybridization

In situ hybridization is a powerful and versatile technique for the detection and localization of nucleic acid sequences within cell or tissue preparations. It may be useful, for example, to identify sites of gene expression, analyze the tissue distribution of transcription, identify and localize viral infection, follow changes in specific mRNA synthesis, and aid in chromosome mapping.

In situ hybridization was performed following an optimized version of the protocol by Lu and Gillett, Cell Vision, 1: 169-176 (1994), using PCR-generated ³³P-labeled riboprobes. Briefly, formalin-fixed, paraffin-embedded human tissues were sectioned, deparaffinized, deproteinized in proteinase K (20 g/ml) for 15 minutes at 37°C, and further processed for *in situ* hybridization as described by Lu and Gillett, *supra*. A (³³-P)UTP-labeled antisense riboprobe was generated from a PCR product and hybridized at 55°C overnight. The slides were dipped in Kodak NTB2TM nuclear track emulsion and exposed for 4 weeks.

6.24.1. ³³P-Riboprobe synthesis

6.0 µl (125 mCi) of ³³P-UTP (Amersham BF 1002, SA <2000 Ci/mmol) were speed-vacuum dried. To each tube containing dried ³³P-UTP, the following ingredients were added:

2.0 µl 5x transcription buffer

1.0 µl DTT (100 mM)

2.0 µl NTP mix (2.5 mM: 10 µl each of 10 mM GTP, CTP & ATP + 10 µl H₂O)

1.0 µl UTP (50 µM)

1.0 µl RNAsin

1.0 µl DNA template (1 µg)

1.0 µl H₂O

1.0 µl RNA polymerase (for PCR products T3 = AS, T7 = S, usually)

The tubes were incubated at 37°C for one hour. A total of 1.0 µl RQ1 DNase was added, followed by incubation at 37°C for 15 minutes. A total of 90 µl TE (10 mM Tris pH 7.6/1 mM EDTA pH 8.0) was added, and the mixture was pipetted onto DE81 paper. The remaining solution was loaded in a MICROCON-50TM ultrafiltration unit, and spun using program 10 (6 minutes). The filtration unit was inverted over a second tube and spun using program 2 (3 minutes). After the final recovery spin, a total of 100 µl TE was added, then 1 µl of the final product was pipetted on DE81 paper and counted in 6 ml of BIOFLUOR IITM.

The probe was run on a TBE/urea gel. A total of 1-3 µl of the probe or 5 µl of RNA Mrk III was added to 3 µl of loading buffer. After heating on a 95°C heat block for three minutes, the gel was immediately placed on ice. The wells of gel were flushed, and the sample was loaded and run at 180-250 volts for 45 minutes. The gel was wrapped in plastic wrap (SARANTM brand) and exposed to XAR film with an intensifying screen in a -70°C freezer one hour to overnight.

6.24.2. ³³P-Hybridization6.24.2.1. *Pretreatment of frozen sections*

The slides were removed from the freezer, placed on aluminum trays, and thawed at room temperature for 5 minutes. The trays were placed in a 55°C incubator for five minutes to reduce condensation. The slides were fixed for 10 minutes in 4% paraformaldehyde on ice in the fume hood, and washed in 0.5 x SSC for 5 minutes, at room temperature (25 ml 20 x SSC + 975 ml SQ H₂O). After deproteinization in 0.5 µg/ml proteinase K for 10 minutes at 37°C (12.5 µl of 10 mg/ml stock in 250 ml prewarmed RNase-free RNase buffer), the sections were washed in 0.5 x SSC for 10 minutes at room temperature. The sections were dehydrated in 70%, 95%, and 100% ethanol, 2 minutes each.

6.24.2.2. *Pretreatment of paraffin-embedded sections*

The slides were deparaffinized, placed in SQ H₂O, and rinsed twice in 2 x SSC at room temperature, for 5 minutes each time. The sections were deproteinized in 20 µg/ml proteinase K (500 µl of 10 mg/ml in 250 ml RNase-free RNase buffer; 37°C, 15 minutes) for human embryo tissue, or 8 x proteinase K (100 µl in 250 ml RNase buffer, 37°C, 30 minutes) for formalin tissues. Subsequent rinsing in 0.5 x SSC and dehydration were performed as described above.

6.24.2.3. *Prehybridization*

The slides were laid out in a plastic box lined with Box buffer (4 x SSC, 50% formamide) - saturated filter paper. The tissue was covered with 50 µl of hybridization buffer (3.75 g dextran sulfate + 6 ml SQ H₂O), vortexed, and heated in the microwave for 2 minutes with the cap loosened. After cooling on ice, 18.75 ml formamide, 3.75 ml 20 x SSC, and 9 ml SQ H₂O were added, and the tissue was vortexed well and incubated at 42°C for 1-4 hours.

6.24.2.4. *Hybridization*

1.0 x 10⁶ cpm probe and 1.0 µl tRNA (50 mg/ml stock) per slide were heated at 95°C for 3 minutes. The slides were cooled on ice, and 48 µl hybridization buffer was added per slide. After vortexing, 50 µl ³³P mix was added to 50 µl prehybridization on the slide. The slides were incubated overnight at 55°C.

6.24.2.5. *Washes*

Washing was done for 2x10 minutes with 2xSSC, EDTA at room temperature (400 ml 20 x SSC + 16 ml 0.25 M EDTA, V_f=4L), followed by RNaseA treatment at 37°C for 30 minutes (500 µl of 10 mg/ml in 250 ml RNase buffer = 20 µg/ml). The slides were washed 2 x10 minutes with 2 x SSC, EDTA at room temperature. The stringency wash conditions were as follows: 2 hours at 55°C, 0.1 x SSC, EDTA (20 ml 20 x SSC + 16 ml EDTA, V_f=4L).

6.24.2.6. *Oligonucleotides*

In situ analysis was performed on three of the DNA sequences disclosed herein. The primers used to generate the probes and/or the probes employed for these analyses are as follows:

5 DNA33100-p1: 5'GGA TTC TAA TAC GAC TCA CTA TAG GGC CGG GTG GAG GTG GAA CAG AAA
3' (SEQ ID NO:375)

DNA33100-p2: 5' CTA TGA AAT TAA CCC TCA CTA AAG GGA CAC AGA CAG AGC CCC ATA CGC
3' (SEQ ID NO:376)

DNA34431-p1: 5'GGA TTC TAA TAC GAC TCA CTA TAG GGC CAG GGA AAT CCG GAT GTC TC
3' (SEQ ID NO:377)

10 DNA34431-p2: 5' CTA TGA AAT TAA CCC TCA CTA AAG GGA GTA AGG GGA TGC CAC CGA GTA
3' (SEQ ID NO:378)

DNA38268-p1: 5'GGA TTC TAA TAC GAC TCA CTA TAG GGC CAG CTA CCC GCA GGA GGA GG
3' (SEQ ID NO:379)

15 DNA38268-p2: 5'CTA TGA AAT TAA CCC TCA CTA AAG GGA TCC CAG GTG ATG AGG TCC AGA
3' (SEQ ID NO:380)

DNA64908 probe: 5'CCATCTCGGAGACCTTTGTGCAGCGTGTATACCAGCCTTACCTCACCA
CTTGCGACGGACACAGAGCCTGCAGCACCTACCGAACCATCTACCGGAC
TGCCTATCGCCGTAGCCCTGGGGTGACTCCCGCAAGCCTCGCTATGCTTG
CTGCCCTGGTTGGAAGAGGACCAGTGGGCTCCCTGGGGCTTGTGGAGCA
20 GCAATATGCCAGCCTCCATGTGGAATGGAGGGAGTTGCATCCGCCCAG
GACACTGCCGCTGCCCTGTGGGATGGCAGGGAGATACTTGCCAGACAGA
TGTTGATGAATGCAGTACAGGAGAGGCCAGTTGTCCCCAGCGCTGTGTC
AATACTGTGGGAAGTTACTGGTGCCAGGGATGGGAGGGACAAAGCCCAT
CTGCAGATGGGACGCGCTGCCTGTCTAAGGAGGGGCCCTCCCGGTGGCC
25 CCAACCCACAGCAGGAGTGGACAGCA3' (SEQ ID NO:381)

6.24.2.7. *Results*

In situ analysis was performed and the results from these analyses are as follows:

6.24.2.7.1. DNA33100-1159 (PRO229) (Scavenger-R/CD6
homologTNF motif)

A specific positive signal was observed in mononuclear phagocytes (macrophages) of fetal and adult spleen, liver, lymph node and thymus. All other tissues were negative.

5 6.24.2.7.2. DNA34431-1177 (PRO263) (CD44)

A specific positive signal was observed in human fetal tissues and placenta over mononuclear cells, with strong expression in epithelial cells of the adrenal cortex. All adult tissues were negative.

6.24.2.7.3. DNA38268-1188 (PRO295) (Integrin)

10 A specific positive signal was observed in human fetal ganglion cells, fetal neurons, adult adrenal medulla and adult neurons. All other tissues were negative.

6.24.2.7.4. DNA64908-1163-1 (PRO1449)

15 A specific positive signal was observed in the developing vasculature (from E7-E11), in endothelial cells and in progenitors of endothelial cells in wholemount in situ hybridizations of mouse embryos (Figure 375). Specific expression was also observed in a subset of blood vessels and epidermis from E12 onward. A mouse orthologue of PRO1449 which has about 78% amino acid identity with PRO1449 was used as the probe.

In normal adult tissues, expression was low to absent. When present, expression was confined to the vasculature (Figure 376). Figure 376 further shows that highest expression in adult tissues was observed regionally in vessels running within the white matter of the brain. Elevated expression was also observed in vasculature of many inflamed and diseased tissues, including, but not limited to, tumor vasculature.

20 Following electroporation of the mouse orthologue of PRO1449 into the choroid layer in the eyes of chicken embryos, new vessel formation was observed in the electroporated eye (top right), but not in the control side from the same embryo (top left), or an embryo that was electroporated with a control cDNA (bottom right) (Figure 377).

25 The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the construct(s) deposited, since the deposited embodiment(s) is/are intended as single illustration(s) of certain aspects of the invention and any constructs that are functionally equivalent are within the scope of this invention. The deposit of material(s) herein does not constitute an admission that the written description herein contained is inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific
30 illustrations that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule having at least 80% nucleic acid sequence identity to a nucleotide sequence that encodes an amino acid sequence selected from the group consisting of the amino acid sequence shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure

248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) and Figure 374 (SEQ ID NO:374).

2. An isolated nucleic acid molecule having at least 80% nucleic acid sequence identity to a nucleotide sequence selected from the group consisting of the nucleotide sequence shown in Figure 1 (SEQ ID NO:1), Figure 3 (SEQ ID NO:3), Figure 5 (SEQ ID NO:5), Figure 7 (SEQ ID NO:7), Figure 9 (SEQ ID NO:9), Figure 11 (SEQ ID NO:11), Figure 13 (SEQ ID NO:13), Figure 15 (SEQ ID NO:15), Figure 17 (SEQ ID NO:17), Figure 19 (SEQ ID NO:19), Figure 21 (SEQ ID NO:21), Figure 23 (SEQ ID NO:23), Figure 25 (SEQ ID NO:25), Figure 27 (SEQ ID NO:27), Figure 29 (SEQ ID NO:29), Figure 31 (SEQ ID NO:31), Figure 33 (SEQ ID NO:33), Figure 35 (SEQ ID NO:35), Figure 37 (SEQ ID NO:37), Figure 39 (SEQ ID NO:39), Figure 41 (SEQ ID NO:41), Figure 43 (SEQ ID NO:43), Figure 45 (SEQ ID NO:45), Figure 47 (SEQ ID NO:47), Figure 49 (SEQ ID NO:49), Figure 51 (SEQ ID NO:51), Figure 53 (SEQ ID NO:53), Figure 55 (SEQ ID NO:55), Figure 57 (SEQ ID NO:57), Figure 59 (SEQ ID NO:59), Figure 61 (SEQ ID NO:61), Figure 63 (SEQ ID NO:63), Figure 65 (SEQ ID NO:65), Figure 67 (SEQ ID NO:67), Figure 69 (SEQ ID NO:69), Figure 71 (SEQ ID NO:71), Figure 73 (SEQ ID NO:73), Figures 75A-75B (SEQ ID NO:75), Figure 77 (SEQ ID NO:77), Figure 79 (SEQ ID NO:79), Figure 81 (SEQ ID NO:81), Figure 83 (SEQ ID NO:83), Figure 85 (SEQ ID NO:85), Figure 87 (SEQ ID NO:87), Figure 89 (SEQ ID NO:89), Figure 91 (SEQ ID NO:91), Figure 93 (SEQ ID NO:93), Figure 95 (SEQ ID NO:95), Figure 97 (SEQ ID NO:97), Figure 99 (SEQ ID NO:99), Figure 101 (SEQ ID NO:101), Figure 103 (SEQ ID NO:103), Figure 105 (SEQ ID NO:105), Figure 107 (SEQ ID NO:107), Figure 109 (SEQ ID NO:109), Figure 111 (SEQ ID NO:111), Figure 113 (SEQ ID NO:113), Figure 115 (SEQ

ID NO:115), Figure 117 (SEQ ID NO:117), Figure 119 (SEQ ID NO:119), Figure 121 (SEQ ID NO:121), Figure 123 (SEQ ID NO:123), Figure 125 (SEQ ID NO:125), Figure 127 (SEQ ID NO:127), Figure 129 (SEQ ID NO:129), Figure 131 (SEQ ID NO:131), Figure 133 (SEQ ID NO:133), Figure 135 (SEQ ID NO:135), Figure 137 (SEQ ID NO:137), Figure 139 (SEQ ID NO:139), Figure 141 (SEQ ID NO:141), Figure 143 (SEQ ID NO:143), Figure 145 (SEQ ID NO:145), Figure 147 (SEQ ID NO:147), Figure 149 (SEQ ID NO:149), Figure 151 (SEQ ID NO:151), Figure 153 (SEQ ID NO:153), Figure 155 (SEQ ID NO:155), Figure 157 (SEQ ID NO:157), Figure 159 (SEQ ID NO:159), Figure 161 (SEQ ID NO:161), Figure 163 (SEQ ID NO:163), Figure 165 (SEQ ID NO:165), Figure 167 (SEQ ID NO:167), Figure 169 (SEQ ID NO:169), Figure 171 (SEQ ID NO:171), Figure 173 (SEQ ID NO:173), Figure 175 (SEQ ID NO:175), Figure 177 (SEQ ID NO:177), Figure 179 (SEQ ID NO:179), Figure 181 (SEQ ID NO:181), Figure 183 (SEQ ID NO:183), Figure 185 (SEQ ID NO:185), Figure 187 (SEQ ID NO:187), Figure 189 (SEQ ID NO:189), Figure 191 (SEQ ID NO:191), Figure 193 (SEQ ID NO:193), Figure 195 (SEQ ID NO:195), Figure 197 (SEQ ID NO:197), Figure 199 (SEQ ID NO:199), Figure 201 (SEQ ID NO:201), Figure 203 (SEQ ID NO:203), Figure 205 (SEQ ID NO:205), Figure 207 (SEQ ID NO:207), Figure 209 (SEQ ID NO:209), Figure 211 (SEQ ID NO:211), Figure 213 (SEQ ID NO:213), Figure 215 (SEQ ID NO:215), Figure 217 (SEQ ID NO:217), Figure 219 (SEQ ID NO:219), Figure 221 (SEQ ID NO:221), Figure 223 (SEQ ID NO:223), Figure 225 (SEQ ID NO:225), Figure 227 (SEQ ID NO:227), Figure 229 (SEQ ID NO:229), Figure 231 (SEQ ID NO:231), Figure 233 (SEQ ID NO:233), Figure 235 (SEQ ID NO:235), Figure 237 (SEQ ID NO:237), Figure 239 (SEQ ID NO:239), Figure 241 (SEQ ID NO:241), Figure 243 (SEQ ID NO:243), Figure 245 (SEQ ID NO:245), Figure 247 (SEQ ID NO:247), Figure 249 (SEQ ID NO:249), Figure 251 (SEQ ID NO:251), Figure 253 (SEQ ID NO:253), Figure 255 (SEQ ID NO:255), Figure 257 (SEQ ID NO:257), Figure 259 (SEQ ID NO:259), Figure 261 (SEQ ID NO:261), Figure 263 (SEQ ID NO:263), Figure 265 (SEQ ID NO:265), Figure 267 (SEQ ID NO:267), Figure 269 (SEQ ID NO:269), Figure 271 (SEQ ID NO:271), Figure 273 (SEQ ID NO:273), Figure 275 (SEQ ID NO:275), Figure 277 (SEQ ID NO:277), Figure 279 (SEQ ID NO:279), Figure 281 (SEQ ID NO:281), Figure 283 (SEQ ID NO:283), Figure 285 (SEQ ID NO:285), Figure 287 (SEQ ID NO:287), Figures 289A-289B (SEQ ID NO:289), Figure 291 (SEQ ID NO:291), Figure 293 (SEQ ID NO:293), Figure 295 (SEQ ID NO:295), Figure 297 (SEQ ID NO:297), Figure 299 (SEQ ID NO:299), Figure 301 (SEQ ID NO:301), Figure 303 (SEQ ID NO:303), Figure 305 (SEQ ID NO:305), Figure 307 (SEQ ID NO:307), Figure 309 (SEQ ID NO:309), Figures 311A-311B (SEQ ID NO:311), Figure 313 (SEQ ID NO:313), Figure 315 (SEQ ID NO:315), Figure 317 (SEQ ID NO:317), Figure 319 (SEQ ID NO:319), Figure 321 (SEQ ID NO:321), Figure 323 (SEQ ID NO:323), Figure 325 (SEQ ID NO:325), Figure 327 (SEQ ID NO:327), Figure 329 (SEQ ID NO:329), Figure 331 (SEQ ID NO:331), Figure 333 (SEQ ID NO:333), Figure 335 (SEQ ID NO:335), Figure 337 (SEQ ID NO:337), Figure 339 (SEQ ID NO:339), Figure 341 (SEQ ID NO:341), Figure 343 (SEQ ID NO:343), Figure 345 (SEQ ID NO:345), Figure 347 (SEQ ID NO:347), Figure 349 (SEQ ID NO:349), Figures 351A-351B (SEQ ID NO:351), Figure 353 (SEQ ID NO:353), Figure 355 (SEQ ID NO:355), Figure 357 (SEQ ID NO:357), Figure 359 (SEQ ID NO:359), Figure 361 (SEQ ID NO:361), Figure 363 (SEQ ID NO:363), Figure 365 (SEQ ID NO:365), Figure 367 (SEQ ID NO:367), Figure 369 (SEQ ID NO:369), Figure 371 (SEQ ID NO:371) and Figure 373 (SEQ ID NO:373).

3. An isolated nucleic acid molecule having at least 80% nucleic acid sequence identity to a nucleotide sequence selected from the group consisting of the full-length coding sequence of the nucleotide sequence shown in Figure 1 (SEQ ID NO:1), Figure 3 (SEQ ID NO:3), Figure 5 (SEQ ID NO:5), Figure 7 (SEQ ID NO:7), Figure 9 (SEQ ID NO:9), Figure 11 (SEQ ID NO:11), Figure 13 (SEQ ID NO:13), Figure 15 (SEQ ID NO:15), Figure 17 (SEQ ID NO:17), Figure 19 (SEQ ID NO:19), Figure 21 (SEQ ID NO:21), Figure 23 (SEQ ID NO:23), Figure 25 (SEQ ID NO:25), Figure 27 (SEQ ID NO:27), Figure 29 (SEQ ID NO:29), Figure 31 (SEQ ID NO:31), Figure 33 (SEQ ID NO:33), Figure 35 (SEQ ID NO:35), Figure 37 (SEQ ID NO:37), Figure 39 (SEQ ID NO:39), Figure 41 (SEQ ID NO:41), Figure 43 (SEQ ID NO:43), Figure 45 (SEQ ID NO:45), Figure 47 (SEQ ID NO:47), Figure 49 (SEQ ID NO:49), Figure 51 (SEQ ID NO:51), Figure 53 (SEQ ID NO:53), Figure 55 (SEQ ID NO:55), Figure 57 (SEQ ID NO:57), Figure 59 (SEQ ID NO:59), Figure 61 (SEQ ID NO:61), Figure 63 (SEQ ID NO:63), Figure 65 (SEQ ID NO:65), Figure 67 (SEQ ID NO:67), Figure 69 (SEQ ID NO:69), Figure 71 (SEQ ID NO:71), Figure 73 (SEQ ID NO:73), Figures 75A-75B (SEQ ID NO:75), Figure 77 (SEQ ID NO:77), Figure 79 (SEQ ID NO:79), Figure 81 (SEQ ID NO:81), Figure 83 (SEQ ID NO:83), Figure 85 (SEQ ID NO:85), Figure 87 (SEQ ID NO:87), Figure 89 (SEQ ID NO:89), Figure 91 (SEQ ID NO:91), Figure 93 (SEQ ID NO:93), Figure 95 (SEQ ID NO:95), Figure 97 (SEQ ID NO:97), Figure 99 (SEQ ID NO:99), Figure 101 (SEQ ID NO:101), Figure 103 (SEQ ID NO:103), Figure 105 (SEQ ID NO:105), Figure 107 (SEQ ID NO:107), Figure 109 (SEQ ID NO:109), Figure 111 (SEQ ID NO:111), Figure 113 (SEQ ID NO:113), Figure 115 (SEQ ID NO:115), Figure 117 (SEQ ID NO:117), Figure 119 (SEQ ID NO:119), Figure 121 (SEQ ID NO:121), Figure 123 (SEQ ID NO:123), Figure 125 (SEQ ID NO:125), Figure 127 (SEQ ID NO:127), Figure 129 (SEQ ID NO:129), Figure 131 (SEQ ID NO:131), Figure 133 (SEQ ID NO:133), Figure 135 (SEQ ID NO:135), Figure 137 (SEQ ID NO:137), Figure 139 (SEQ ID NO:139), Figure 141 (SEQ ID NO:141), Figure 143 (SEQ ID NO:143), Figure 145 (SEQ ID NO:145), Figure 147 (SEQ ID NO:147), Figure 149 (SEQ ID NO:149), Figure 151 (SEQ ID NO:151), Figure 153 (SEQ ID NO:153), Figure 155 (SEQ ID NO:155), Figure 157 (SEQ ID NO:157), Figure 159 (SEQ ID NO:159), Figure 161 (SEQ ID NO:161), Figure 163 (SEQ ID NO:163), Figure 165 (SEQ ID NO:165), Figure 167 (SEQ ID NO:167), Figure 169 (SEQ ID NO:169), Figure 171 (SEQ ID NO:171), Figure 173 (SEQ ID NO:173), Figure 175 (SEQ ID NO:175), Figure 177 (SEQ ID NO:177), Figure 179 (SEQ ID NO:179), Figure 181 (SEQ ID NO:181), Figure 183 (SEQ ID NO:183), Figure 185 (SEQ ID NO:185), Figure 187 (SEQ ID NO:187), Figure 189 (SEQ ID NO:189), Figure 191 (SEQ ID NO:191), Figure 193 (SEQ ID NO:193), Figure 195 (SEQ ID NO:195), Figure 197 (SEQ ID NO:197), Figure 199 (SEQ ID NO:199), Figure 201 (SEQ ID NO:201), Figure 203 (SEQ ID NO:203), Figure 205 (SEQ ID NO:205), Figure 207 (SEQ ID NO:207), Figure 209 (SEQ ID NO:209), Figure 211 (SEQ ID NO:211), Figure 213 (SEQ ID NO:213), Figure 215 (SEQ ID NO:215), Figure 217 (SEQ ID NO:217), Figure 219 (SEQ ID NO:219), Figure 221 (SEQ ID NO:221), Figure 223 (SEQ ID NO:223), Figure 225 (SEQ ID NO:225), Figure 227 (SEQ ID NO:227), Figure 229 (SEQ ID NO:229), Figure 231 (SEQ ID NO:231), Figure 233 (SEQ ID NO:233), Figure 235 (SEQ ID NO:235), Figure 237 (SEQ ID NO:237), Figure 239 (SEQ ID NO:239), Figure 241 (SEQ ID NO:241), Figure 243 (SEQ ID NO:243), Figure 245 (SEQ ID NO:245), Figure 247 (SEQ ID NO:247), Figure 249 (SEQ ID NO:249), Figure 251 (SEQ ID NO:251), Figure

253 (SEQ ID NO:253), Figure 255 (SEQ ID NO:255), Figure 257 (SEQ ID NO:257), Figure 259 (SEQ ID NO:259), Figure 261 (SEQ ID NO:261), Figure 263 (SEQ ID NO:263), Figure 265 (SEQ ID NO:265), Figure 267 (SEQ ID NO:267), Figure 269 (SEQ ID NO:269), Figure 271 (SEQ ID NO:271), Figure 273 (SEQ ID NO:273), Figure 275 (SEQ ID NO:275), Figure 277 (SEQ ID NO:277), Figure 279 (SEQ ID NO:279), Figure 281 (SEQ ID NO:281), Figure 283 (SEQ ID NO:283), Figure 285 (SEQ ID NO:285), Figure 287 (SEQ ID NO:287), Figures 289A-289B (SEQ ID NO:289), Figure 291 (SEQ ID NO:291), Figure 293 (SEQ ID NO:293), Figure 295 (SEQ ID NO:295), Figure 297 (SEQ ID NO:297), Figure 299 (SEQ ID NO:299), Figure 301 (SEQ ID NO:301), Figure 303 (SEQ ID NO:303), Figure 305 (SEQ ID NO:305), Figure 307 (SEQ ID NO:307), Figure 309 (SEQ ID NO:309), Figures 311A-311B (SEQ ID NO:311), Figure 313 (SEQ ID NO:313), Figure 315 (SEQ ID NO:315), Figure 317 (SEQ ID NO:317), Figure 319 (SEQ ID NO:319), Figure 321 (SEQ ID NO:321), Figure 323 (SEQ ID NO:323), Figure 325 (SEQ ID NO:325), Figure 327 (SEQ ID NO:327), Figure 329 (SEQ ID NO:329), Figure 331 (SEQ ID NO:331), Figure 333 (SEQ ID NO:333), Figure 335 (SEQ ID NO:335), Figure 337 (SEQ ID NO:337), Figure 339 (SEQ ID NO:339), Figure 341 (SEQ ID NO:341), Figure 343 (SEQ ID NO:343), Figure 345 (SEQ ID NO:345), Figure 347 (SEQ ID NO:347), Figure 349 (SEQ ID NO:349), Figures 351A-351B (SEQ ID NO:351), Figure 353 (SEQ ID NO:353), Figure 355 (SEQ ID NO:355), Figure 357 (SEQ ID NO:357), Figure 359 (SEQ ID NO:359), Figure 361 (SEQ ID NO:361), Figure 363 (SEQ ID NO:363), Figure 365 (SEQ ID NO:365), Figure 367 (SEQ ID NO:367), Figure 369 (SEQ ID NO:369), Figure 371 (SEQ ID NO:371) and Figure 373 (SEQ ID NO:373).

4. An isolated nucleic acid molecule having at least 80% nucleic acid sequence identity to the full-length coding sequence of the DNA deposited under any ATCC accession number shown in Table 7.

5. A vector comprising the nucleic acid of Claim 1.

6. A host cell comprising the vector of Claim 5.

7. The host cell of Claim 6, wherein said cell is a CHO cell.

8. The host cell of Claim 6, wherein said cell is an *E. coli*.

9. The host cell of Claim 6, wherein said cell is a yeast cell.

10. A process for producing a PRO polypeptide comprising culturing the host cell of Claim 6 under conditions suitable for expression of said PRO polypeptide and recovering said PRO polypeptide from the cell culture.

11. An isolated polypeptide having at least 80% amino acid sequence identity to an amino acid sequence selected from the group consisting of the amino acid sequence shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID

NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) and Figure 374 (SEQ ID NO:374).

12. An isolated polypeptide having at least 80% amino acid sequence identity to an amino acid sequence encoded by the full-length coding sequence of the DNA deposited under any ATCC accession number shown in Table 7.

13. A chimeric molecule comprising a polypeptide according to Claim 11 fused to a heterologous amino acid sequence.

14. The chimeric molecule of Claim 13, wherein said heterologous amino acid sequence is an epitope tag sequence.

15. The chimeric molecule of Claim 13, wherein said heterologous amino acid sequence is a Fc region of an immunoglobulin.

16. An antibody which specifically binds to a polypeptide according to Claim 11.

17. The antibody of Claim 16, wherein said antibody is a monoclonal antibody, a humanized antibody or a single-chain antibody.

18. An isolated nucleic acid molecule having at least 80% nucleic acid sequence identity to:

(a) a nucleotide sequence encoding the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26),
Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64),
Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102),
Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264)

NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), lacking its associated signal peptide;

(b) a nucleotide sequence encoding an extracellular domain of the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure

144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), with its associated signal peptide; or

(c) a nucleotide sequence encoding an extracellular domain of the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18

(SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure

284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), lacking its associated signal peptide.

19. An isolated polypeptide having at least 80% amino acid sequence identity to:

(a) an amino acid sequence of the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID

NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), lacking its associated signal peptide;

(b) an amino acid sequence of an extracellular domain of the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26

(SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure

292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), with its associated signal peptide; or

(c) an amino acid sequence of an extracellular domain of the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID

NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), lacking its associated signal peptide.

20. A method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal a therapeutically effective amount of a polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32),

Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure

298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), or agonist or antagonist thereof.

21. The method according to Claim 20, wherein the mammal is human.

22. The method of Claim 21, wherein the human has suffered myocardial infarction.

23. The method of Claim 21, wherein the human has cardiac hypertrophy, trauma, a cancer, or age-related macular degeneration.

24. The method of Claim 23, wherein the cardiac hypertrophy is characterized by the presence of an elevated level of $\text{PGF}_{2\alpha}$.

25. The method of Claim 20, wherein the polypeptide is administered together with a cardiovascular, endothelial or angiogenic agent.

26. The method of Claim 23, wherein the polypeptide is administered following primary angioplasty.

27. The method of Claim 20, wherein the cardiovascular, endothelial or angiogenic disorder is cancer.

28. The method of Claim 27, wherein the polypeptide is administered in combination with a chemotherapeutic agent, a growth inhibitory agent or a cytotoxic agent.

29. The method of Claim 20, wherein said agonist is an antibody to said polypeptide.

30. The method of Claim 20, wherein said antagonist is an antibody to said polypeptide.

31. A method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal a nucleic acid molecule that encodes a polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure

256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), or agonist or antagonist thereof.

32. The method of Claim 31, wherein said agonist is an antibody to said polypeptide.

33. The method of Claim 31, wherein said antagonist is an antibody to said polypeptide.

34. The method of Claim 31, wherein the mammal is human.

35. The method of Claim 31, wherein the nucleic acid molecule is administered via *ex vivo* gene therapy.

36. A method for inhibiting endothelial cell growth in a mammal comprising administering to the mammal a PRO247, PRO720 or PRO4302 polypeptide or agonist thereof, wherein endothelial cell growth in said mammal is inhibited.

37. A method for stimulating endothelial cell growth in a mammal comprising administering to the mammal a PRO21, PRO181, PRO205, PRO214, PRO221, PRO229, PRO231, PRO238, PRO241, PRO247, PRO256, PRO258, PRO263, PRO265, PRO295, PRO321, PRO322, PRO337, PRO363, PRO365, PRO444, PRO533, PRO697, PRO725, PRO771, PRO788, PRO791, PRO819, PRO827, PRO828, PRO836, PRO846, PRO865, PRO1005, PRO1006, PRO1007, PRO1025, PRO1029, PRO1054, PRO1071, PRO1075, PRO1079,

PRO1080, PRO1114, PRO1131, PRO1155, PRO1160, PRO1184, PRO1186, PRO1190, PRO1192, PRO1195, PRO1244, PRO1272, PRO1273, PRO1274, PRO1279, PRO1283, PRO1286, PRO1306, PRO1309, PRO1325, PRO1329, PRO1347, PRO1356, PRO1376, PRO1382, PRO1411, PRO1412, PRO1419, PRO1474, PRO1477, PRO1488, PRO1508, PRO1550, PRO1556, PRO1760, PRO1782, PRO1787, PRO1801, PRO1868, PRO1887, PRO1890, PRO3438, PRO3444, PRO4302, PRO4324, PRO4333, PRO4341, PRO4342, PRO4353, PRO4354, PRO4356, PRO4371, PRO4405, PRO4408, PRO4422, PRO4425, PRO4499, PRO5723, PRO5725, PRO5737, PRO5776, PRO6006, PRO6029, PRO6071, PRO7436, PRO9771, PRO9821, PRO9873, PRO10008, PRO10096, PRO19670, PRO20040, PRO20044, PRO21055, PRO21384 or PRO28631 polypeptide, or agonist thereof, wherein endothelial cell growth in said mammal is stimulated.

38. A method for inducing cardiac hypertrophy in a mammal comprising administering to the mammal a PRO21 polypeptide or agonist thereof, wherein cardiac hypertrophy in said mammal is induced.

39. A method for stimulating angiogenesis induced by a PRO1376 or PRO1449 polypeptide in a mammal comprising administering a therapeutically effective amount of said polypeptide to the mammal, wherein said angiogenesis is stimulated.

40. A method for inducing endothelial cell apoptosis comprising administering to the endothelial cell a PRO4302 polypeptide or agonist thereof, wherein apoptosis in said endothelial cell is induced.

41. A method for stimulating smooth muscle cell growth comprising administering to the smooth muscle cell a PRO162, PRO181, PRO182, PRO195, PRO204, PRO221, PRO230, PRO256, PRO258, PRO533, PRO697, PRO725, PRO738, PRO826, PRO836, PRO840, PRO846, PRO865, PRO982, PRO1025, PRO1029, PRO1071, PRO1080, PRO1083, PRO1134, PRO1160, PRO1182, PRO1184, PRO1186, PRO1192, PRO1265, PRO1274, PRO1279, PRO1283, PRO1306, PRO1308, PRO1309, PRO1325, PRO1337, PRO1338, PRO1343, PRO1376, PRO1387, PRO1411, PRO1412, PRO1415, PRO1434, PRO1474, PRO1488, PRO1550, PRO1556, PRO1567, PRO1600, PRO1754, PRO1758, PRO1760, PRO1787, PRO1865, PRO1868, PRO1917, PRO1928, PRO3438, PRO3562, PRO4302, PRO4333, PRO4345, PRO4353, PRO4354, PRO4405, PRO4408, PRO4430, PRO4503, PRO5725, PRO6714, PRO9771, PRO9820, PRO9940, PRO10096, PRO21055, PRO21184 or PRO21366 polypeptide, or agonist thereof, wherein smooth muscle cell growth in said smooth muscle cell is stimulated.

42. A method for inducing endothelial cell tube formation comprising administering to the endothelial cell a PRO178, PRO195, PRO228, PRO301, PRO302, PRO532, PRO724, PRO730, PRO734, PRO793, PRO871, PRO938, PRO1012, PRO1120, PRO1139, PRO1198, PRO1287, PRO1361, PRO1864, PRO1873, PRO2010, PRO3579, PRO4313, PRO4527, PRO4538, PRO4553, PRO4995, PRO5730, PRO6008, PRO7223, PRO7248 or PRO7261 polypeptide, or agonist thereof, wherein tube formation in said endothelial cell is induced.

1/392

FIGURE 1

GCCCCACGCGTCCGATGGCGTTACGTTTCGCGGCCTTCTGCTACATGCTGGCGCTGCTGCT
CACTGCCGCGCTCATCTTCTTCGCCATTTGGCACATTATAGCATTTGATGAGCTGAAGAC
TGATTACAAGAATCCTATAGACCAGTGTAATACCCTGAATCCCCTTGTAAGTCCCAGAGTA
CCTCATCCACGCTTTCCTTCTGTGTCATGTTTCTTTGTGCAGCAGAGTGGCTTACACTGGG
TCTCAATATGCCCCCTCTTGGCATATCATATTTGGAGGTATATGAGTAGACCAGTGATGAG
TGGCCAGGACTCTATGACCCTACAACCATCATGAATGCAGATATTCTAGCATATTGTCA
GAAGGAAGGATGGTGCAAATTAGCTTTTTTATCTTCTAGCATTTTTTTTACTACCTATATGG
CATGATCTATGTTTTGGTGAGCTCTTAGAACAACACACAGAAGAATTGGTCCAGTTAAGT
GCATGCAAAAAGCCACCAAATGAAGGGATTCTATCCAGCAAGATCCTGTCCAAGAGTAGC
CTGTGGAATCTGATCAGTTACTTTAAAAAATGACTCCTTATTTTTTAAATGTTTCCACAT
TTTTGCTTGTGGAAAGACTGTTTTCATATGTTATACTCAGATAAAGATTTTTAAATGGTAT
TACGTATAAATTAATATAAAATGATTACCTCTGGTGTTGACAGGTTTGAAGTTGCACTTC
TTAAGGAACAGCCATAATCCTCTGAATGATGCATTAATTACTGACTGTCCTAGTACATTG
GAAGCTTTTGTTTATAGGAAGTTGTAGGGCTCATTTTGGTTTCATTGAAACAGTATCTAA
TTATAAATTAGCTGTAGATATCAGGTGCTTCTGATGAAGTGAAAATGTATATCTGACTAG
TGGGAAAGTTTATGGGTTTCCTCATCTGTCATGTCGATGATTATATATGGATACATTTAC
AAAAATAAAAAGCGGGAATTTTCCCTTCGCTTGAATATTATCCCTGTATATTGCATGAAT
GAGAGATTTCCCATATTTCCATCAGAGTAATAAATATACTTGCTTTAATTCTTAAGCATA
AGTAAACATGATATAAAAAATATATGCTGAATTACTTGTGAAGAATGCATTTAAAGCTATT
TTAAATGTGTTTTTATTTGTAAGACATTACTTATTAAGAAATTGGTTATTATGCTTACTG
TTCTAATCTGGTGGTAAAGGTATTCTTAAGAATTTGCAGGTACTACAGATTTTCAAACT
GAATGAGAGAAAATTGTATAACCATCCTGCTGTTCTTTAGTGCAATACAATAAACTCT
GAAATTAAGACTC

2/392

FIGURE 2

MAFTFAAFCYMLALLLTAAALIFFAIWHIIAFDELKTDYKNPIDQCNTLNPLVLPEYLIHA
FFCVMFLCAAEWLTGLNMPLLAYHIWRYMSRPVMSGPLYDPTTIMNADILAYCQKEGW
CKLAFYLLAFFYYLYGMIYVLVSS

Important features:

Signal peptide:

amino acids 1-20

Type II transmembrane domain:

amino acids 11-31

Other transmembrane domain:

amino acids 57-77 and 123-143

3/392

FIGURE 3

GGCTCAGAGGCCCCACTGGACCCTCGGCTCTTCCTTGGACTTCTTGTGTGTTCTGTGAGC
TTCGCTGGATT CAGGGTCTTGGGCATCAGAGGTGAGAGGGTGGGAAGGTCCGCCGCGATG
GGGAAGCCCTGGCTGCGTGCGCTACAGCTGCTGCTCCTGCTGGGCGCGTCGTGGGCGCGG
GCGGGCGCCCCGCGCTGCACCTACACCTTCGTGCTGCCCCCGCAGAAGTTCACGGGCGCT
GTGTGCTGGAGCGGCCCCGCATCCACGCGGGCGACGCCCCGAGGCCGCAACGCCAGCGAG
CTGGCGGCGCTGCGCATGCGCGTCGGCCGCCACGAGGAGCTGTTACGCGAGCTGCAGAGG
CTGGCGGCGGCGGACGGCGCCGTGGCCGGCGAGGTGCGCGCGCTGCGCAAGGAGAGCCGC
GGCCTGAGCGCGCGCCTGGGCCAGTTGCGCGCGCAGCTGCAGCACGAGGCGGGGCCCGGG
GCGGGCCCCGGGGGCGGATCTGGGGGCGGAGCCTGCCGCGGCGCTGGCGCTGCTCGGGGAG
CGCGTGCTCAACGCGTCCGCCGAGGCTCAGCGCGCAGCCGCCCGGTTCCACCAGCTGGAC
GTCAAGTTCGCGAGCTGGCGCAGCTCGTCACCCAGCAGAGCAGTCTCATCGCCGCGCTG
GAGCGCCTGTGCCCCGGGAGGCGCGGGCGGGCAGCAGCAGGTCTGCCGCCACCCCCACTG
GTGCCTGTGGTTCCGGTCCGTCTTGTGGGTAGCACCCAGTGACACCAGTAGGATGCTGGAC
CCAGCCCCAGAGCCCCAGAGAGACCAGACCCAGAGACAGCAGGAGCCCATGGCTTCTCCC
ATGCCTGCAGGTACCCCTGCGGTCCCCACCAAGCCTGTGGGCCCGTGGCAGGATTGTGCA
GAGGCCCGCCAGGCAGGCCATGAACAGAGTGGAGTGTATGAACTGCGAGTGGGCCGTAC
GTAGTGT CAGTATGGTGTGAGCAGCAACTGGAGGGTGGAGGCTGGACTGTGATCCAGCGG
AGGCAAGATGGTT CAGTCAACTTCTTCACTACCTGGCAGCACTATAAGGCGGGCTTTGGG
CGGCCAGACGGAGAATACTGGCTGGGCCTTGAACCCGTGTATCAGCTGACCAGCCGTGGG
GACCATGAGCTGCTGGTTCTCCTGGAGGACTGGGGGGGCGGTGGAGCACGTGCCCACTAT
GATGGCTTCTCCCTGGAACCCGAGAGCGACCACTACCGCCTGCGGCTTGGCCAGTACCAT
GGTGATGCTGGAGACTCTCTTTCTGGCACAATGACAAGCCCTTCAGCACCGTGGATAGG
GACCGAGACTCCTATTCTGGTAACTGTGCCCTGTACCAGCGGGGAGGCTGGTGGTACCAT
GCCTGTGCCCACTCCAACCTCAACGGTGTGTGGCACCACGGCGGCCACTACCGAAGCCGC
TACCAGGATGGTGTCTACTGGGCTGAGTTTCGTGGTGGGGCATATTCTCTCAGGAAGGCC
GCCATGCTCATTGCGCCCCCTGAAGCTGTGACTCTGTGTTCTCTGTCCCCTAGGCCCTAG
AGGACATTGGTCAGCAGGAGCCCAAGTTGTTCTGGCCACACCTTCTTTGTGGCTCAGTGC
CAATGTGTCCACAGAACTTCCCACTGTGGATCTGTGACCCTGGGCGCTGAAAATGGGAC
CCAGGAATCCCCCGCTCAATATCTTGGCCTCAGATGGCTCCCCAAGGTCATTCATATCT
CGGTTTGAGCTCATATCTTATAATAACACAAAGTAGCCAC

4/392

FIGURE 4

MGKPWLRALQLLLLLLGASWARAGAPRCTYTFVLPPQKFTGAVCWSGPASTRATPEAANAS
ELAALRMVRGRHEELLRELQRLAAADGAVAGEVRALRKESRGLSARLGQLRAQLQHEAGP
GAGPGADLGAEPAAALALLGERVLNASAEAQRAAARFHQLDVKFRELAQLVTQQSSLIAR
LERLCPPGAGGQQQVLPPPPLVPVVPVRLVGSTSDTSRMLDPAPEPQRDQTORQQEPMAS
PMPAGHPAVPTKPVGPWQDCAEARQAGHEQSGVYELRVGRHVSVWCEQQLEGGGWTVIQ
RRQDGSVNFFTTWQHYKAGFGRPDGEYWLGLEPVYQLTSRGDHELLVLLEDWGGRGARAH
YDGFSLPEPESDHYRLRLGQYHGDAGDSLWHNDKPFSTVDRDRDSYSGNCALYQRGGWWY
HACAHSNLNGVWHHGGHYRSRYQDGVYWAEFRGGAYSLRKAAMLIRPLKL

Signal peptide:

Amino acids 1-20

N-glycosylation sites:

Amino acids 58-62;145-149

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 97-101

Tyrosine kinase phosphorylation site:

Amino acids 441-448

N-myristoylation sites:

Amino acids

16-22;23-29;87-93;108-114;121-127;125-131;129-135;187-193;29
3-299;353-359;378-384;445-451;453-459**Cell attachment sequence:**

Amino acids 340-343

Fibrinogen beta and gamma chains C-terminal domain signature:

Amino acids 418-431

5/392

FIGURE 5

CCCACGCGTCCGGCGCCGTGGCCTCGCGTCCATCTTTGCCGTTCTCTCGGACCTGTCACA
AAGGAGTCGCGCCGCGCCGCGCCGCCCCCTCCCTCCGGTGGGCCCCGGGAGGTAGAGAAAGT
CAGTGCCACAGCCCGACCGCGCTGCTCTGAGCCCTGGGCACGCGGAACGGGAGGGAGTCT
GAGGGTTGGGGACGTCTGTGAGGGAGGGGAACAGCCGCTCGAGCCTGGGGCGGGCGGACC
GGACTGGGGCCGGGGTAGGCTCTGGAAGGGCCCGGGAGAGAGGTGGCGTTGGTCAGAAC
CTGAGAAACAGCCGAGAGGTTTTCCACCGAGGCCCGCGCTTGAGGGATCTGAAGAGGTTT
CTAGAAGAGGGTGTTCCCTCTTTCCGGGGGTCTCACCAGAAGAGGTTCTTGGGGGTCTGCC
CTTCTGAGGAGGCTGCGGCTAACAGGGCCCAAGACTGCCATTGGATGTCCAGAATCCCCT
GTAGTTGATAATGTTGGGAATAAGCTCTGCAACTTTCTTTGGCATTTCAGTTGTTAAAAAC
AAATAGGATGCAAATTCCTCAACTCCAGGTTATGAAAACAGTACTTGAAAACTGAAAAC
TACCTAAATGATCGTCTTTGGTTGGGCCGTGTTCTTAGCGAGCAGAAGCCTTGGCCAGGG
TCTGTTGTTGACTCTCGAAGAGCACATAGCCCACTTCCTAGGGACTGGAGGTGCCGCTAC
TACCATGGGTAATTCCTGTATCTGCCGAGATGACAGTGGAACAGATGACAGTGTTGACAC
CCAACAGCAACAGGCCGAGAACAGTGACAGTACCCACTGCTGACACAAGGAGCCAACCACG
GGACCCTGTTCCGCCACCAAGGAGGGGCCGAGGACCTCATGAGCCAAGGAGAAAGAAACA
AAATGTGGATGGGCTAGTGTTGGACACACTGGCAGTAATACGGAATCTTGTAGATAAGTA
AGTATCTGACTCACGGTCACCTCCAGTGGAATGAAAAGTGTTCTGCCCCGAACCATGACT
TTAGGACTCCTTCAGTTCCTTTAGGACATACTCGCCAAGCCTTGTGCTCACAGGGCAAAG
GAGAATATTTTAATGCTCCGCTGATGGCAGAGTAAATGATAAGATTTGATGTTTTTGCTT
GCTGTCATCTACTTTGTCTGGAATGTCTAAATGTTTCTGTAGCAGAAAACACGATAAAG
CTATGATCTTTATTAGAG

6/392

FIGURE 6

MIVFGWAVFLASRSLGQGLLLTLEEHIAHFLGTGGAATTMGNSCICRDDSGTDDSVDTQQ
QQAENSAVPTADTRSQPRDPVRPPRRGRGPHEPRRKKQNV DGLVLDTLAVIRTLVDKO

Signal peptide:
amino acids 1-16

Casein kinase II phosphorylation site:
amino acids 22-26, 50-54, 113-117

N-myristoylation site:
amino acids 18-24, 32-38, 34-40, 35-41, 51-57

7/392

FIGURE 7

CGGACGCGTGGGGGAAACCCCTTCCGAGAAAACAGCAACAAGCTGAGCTGCTGTGACAGAG
GGGAACAAGATGGCGGCGCCGAAGGGGAGCCTCTGGGTGAGGACCCAACTGGGGCTCCCG
CCGCTGCTGCTGCTGACCATGGCCTTGCCCGGAGGTTTCGGGGACCGCTTCGGCTGAAGCA
TTTGACTCGGTCTTGCGGTGATACGGCGTCTTGCCACCGGGCCTGTCAGTTGACCTACCCC
TTGCACACCTACCCCTAAGGAAGAGGAGTTGTACGCATGTCAGAGAGGTTGCAGGCTGTTT
TCAATTTGTCAGTTTGTGGATGATGGAATTGACTTAAATCGAACTAAATTGGAATGTGAA
TCTGCATGTACAGAAGCATATTCCTCAATCTGATGAGCAATATGCTTGCCATCTTGTTGC
CAGAATCAGCTGCCATTCGCTGAACTGAGACAAGAACAACCTTATGTCCCTGATGCCAAAA
ATGCACCTACTCTTTCCTCTAACTCTGGTGAGGTCATTCTGGAGTGACATGATGGACTCC
GCACAGAGCTTCATAACCTCTTCATGGACTTTTTATCTTCAAGCCGATGACGGAAAAATA
GTTATATTCCAGTCTAAGCCAGAAATCCAGTACGCACCACATTTGGAGCAGGAGCCTACA
AATTTGAGAGAATCATCTCTAAGCAAAATGTCTTATCTGCAAATGAGAAATTCACAAGCG
CACAGGAATTTTCTTGAAGATGGAGAAAGTGATGGCTTTTTAAGATGCCTCTCTCTTAAC
TCTGGGTGGATTTTAACTACAACCTCTTGTCTCTCGGTGATGGTATTGCTTTGGATTTGT
TGTGCAACTGTTGCTACAGCTGTGGAGCAGTATGTTCCCTCTGAGAAGCTGAGTATCTAT
GGTGACTTGGAGTTTATGAATGAACAAAAGCTAAACAGATATCCAGCTTCTTCTCTTG
GTTGTTAGATCTAAACTGAAGATCATGAAGAAGCAGGGCCTCTACCTACAAAAGTGAAT
CTTGCTCATTCTGAAATTTAAGCATTCTTTTAAAGACAAGTGTAATAGACATCTAA
AATCCACTCCTCATAGAGCTTTTAAATGGTTTCATTGGATATAGGCCTTAAGAAATCA
CTATAAAATGCAAATAAAGTTACTCAAATCTGTG

8/392

FIGURE 8

MAAPKGS LWVRTQLGLPPLLLLTMALAGGSGTASAEAFDSVLGDTASCHRAQQLTYPLHT
YPKEEELYACQRCRLFSICQFVDDGIDLNRKLECESACTEAYSQSDEQYACHLGCONQ
LPFAELRQEQMLSLMPKMHLFPLTLVRSFWSMMDSAQSFITSSWTFYLQADDGKIVIF
QSKPEIQYAPHLEQEPTNLRSSLSKMSYLMRNSQAHRNFLEDGESDGFRLCLSLNSGW
ILTTTLVLSVMVLLWICCATVATAVEQYVPSEKLSIYGDLEFMNEQKLNRYPASSLVVVR
SKTEDHEEAGPLPTKVNLAHSEI

Important features:

Signal peptide:

amino acids 1-31

Transmembrane domain:

amino acids 241-260

N-glycosylation site:

amino acids 90-93

9/392

FIGURE 9

TATTTACCATATCAGATTCACATTCAGTCCTCAGCAAAATGAAGGGCTCCATTTTCACTC
TGTTTTTATTCTCTGTCCTATTTGCCATCTCAGAAGTGCGGAGCAAGGAGTCTGTGAGAC
TCTGTGGGCTAGAATACATACGGACAGTCATCTATATCTGTGCTAGCTCCAGGTGGAGAA
GGCATCTGGAGGGGATCCCTCAAGCTCAGCAAGCTGAGACAGGAACTCCTTCCAGCTCC
CACATAAACGTGAGTTTTCTGAGGAAAATCCAGCGCAAAACCTTCCGAAGGTGGATGCCT
CAGGGGAAGACCGTCTTTGGGGTGGACAGATGCCCACTGAAGAGCTTTGGAAGTCAAAGA
AGCATTCAAGTGATGTCAAGACAAGATTTACAACTTTGTGTTGCACTGATGGCTGTTCCA
TGACTGATTTGAGTGCTCTTTGCTAAGACAAGAGCAAATACCCAATGGGTGGCAGAGCTT
TATCACATGTTTAATTACAGTGTTTTACTGCCTGGTAGAACACTAATATTGTGTTATTAA
AATGATGGCTTTTGGGTAGGCAAACTTCTTTTCTAAAAGGTATAGCTGAGCGGTGAAA
CCACAGTGATCTCTATTTTCTCCCTTTGCCAAGGTTAATGAACTGTTCTTTTCAAATTCT
ACTAATGCTTTGAAATTTCAAATGCTGCGCAAAATTGCAATAAAAATGCTATAAA

10/392

FIGURE 10

MKGSIFTLFLFSVLFAISEVRSKESVRLCGLEYIRTVIYICASSRWRRLHLEGI PQAQQAE
TGNSFQLPHKREFSEENPAQNLPKVDASGEDRLWGGQMPTEELWKS KHSVMSRQDLQTL
CCTDGCSMTDLSALC

Important features:

Signal sequence:

amino acids 1-18

**cAMP- and cGMP-dependent protein kinase phosphorylation
site:**

amino acids 107-111

N-myristoylation sites:

amino acids 3-9, 52-58, 96-102, 125-131

Insulin family signature:

amino acids 121-136

Insulin family proteins:

amino acids 28-46

11/392

FIGURE 11

CCCACGCGTCCGGACAAACTGGAGGTGAAAGGAGCTGGTACTGTCCACTGTGCTGTCCGGT
GCTGAACCTGAGACGCGAGCGGACCAAGTTGCTCCAGCACCTGAAGGCAACGCCCTCTTGC
ACCCTCTGTGCCCTGTGGGACCCGCTTACCAACAGGACCCATATCAACTTGACAAAGGA
GTGTGGTATCGGACGTGGGAGAGAGTCTCTGTTTGCCACCTGGGCGCTCATTGAGGCGT
GACTTTGGAGATTTCTATAGTTTTAGACCAAACATATTTTTTTTTCCCCAGCTAAGACGAT
CTTTTGAGAGTTTTTTTTTTTTATTGTGATTTATATTTCCACAGCGTTTAGGAATCTTTCT
GGGGGACTTTTGTGACTGTAAATAAGGTGAAAAGCAATAAGGATGTTTAAAGTGCTGGT
CAGTTGTCTTGGTTCTCGGATTCATTTTTCTGGAGTCGGAAGGAAGGCCAACCAAAGAAG
GAGGATATGGCCTTAAATCCTATCAGCCTCTAATGAGATTGCGACATAAGCAGGAAAAAA
ATCAAGAAAGTTCAAGAGTCAAAGGATTTATGATTCAGGATGGCCCTTTTGGATCTTGTG
AAAATAAGTACTGTGGTTTGGGAAGACACTGTGTTACCAGCAGAGAGACAGGGCAAGCAG
AATGTGCCTGTATGGACCTTTGCAAACGTCCTACAAACCTGTGTGTGGATCTGACGGAG
AATTCTATGAAAACCACTGTGAAGTGCACAGAGCTGCTTGCCTGAAAAAACAAAAGATTA
CCATTGTTCACAATGAAGACTGCTTCTTTAAAGGAGATAAGTGCAAGACTACTGAATACA
GCAAGATGAAAAATATGCTATTAGATTTACAAATCAAAAATATATTATGCAAGAAAATG
AAAATCCTAATGGCGACGACATATCTCGGAAGAAGCTATTGGTGGATCAAATGTTTAAAT
ATTTTGATGCAGACAGTAATGGACTTGTAGATATTAATGAACTAACTCAGGTGATAAAAC
AGGAAGAACTTGGCAAGGATCTCTTTGATTGTACTTTGTATGTTCTATTGAAATATGATG
ATTTTAATGCTGACAAGCACCTGGCTCTTGAAGAATTTTATAGAGCATTCCAAGTGATCC
AGTTGAGTCTGCCAGAAGATCAGAACTAAGCATCACTGCAGCAACTGTGGGACAAAGTG
CTGTTCTGAGCTGTGCCATTCAAGGAACCCTGAGACCTCCCATTATCTGGAAAAGGAACA
ATATTATTCTAAATAATTTAGATTTGGAAGACATCAATGACTTTGGAGATGATGGGTCCT
TGTATATTACTAAGGTTACCACAACTCACGTTGGCAATTACACCTGCTATGCAGATGGCT
ATGAACAAGTCTATCAGACTCACATCTTCCAAGTGAATGTTCTCCAGTCATCC

12/392

FIGURE 12

MFKCWSVVLVLGFIFLESEGRPTKEGGYGLKSYQPLMRLRHKQEKQESSRVKGFMIQDG
PFGSCENKYCGLGRHCVTSRETGQAECACMDLCKRHYKPVCSDGEFYENHCEVHRAACL
KKQKITIVHNEDCFFKGDCKTTEYSKMKNMLLDLQNQKYIMQENENPNGDDISRKKLLV
DQMFKYFDADSNGLVDINELTQVIKQEELGKDLFDCTLYVLLKYDDFNADKHLALEEFYR
AFQVIQLSLPEDQKLSITAATVGQSAVLSCAIQGTLRPPIIWKRNNIILNNLDLEDINDF
GDDGSLYITKVTTHVGNVTCYADGYEQVYQTHIFQVNVPPVI

Signal sequence:

Amino acids 1-20

N-glycosylation site:

Amino acids 318-322

Tyrosine kinase phosphorylation sites:

Amino acids 21-29;211-220

N-myristoylation sites:

Amino acids 63-69;83-89;317-323

Prokaryotic membrane lipoprotein lipid attachment site:

Amino acids 260-271

13/392

FIGURE 13

TGCCGGGCTGCGGGGCGCCTTGACTCTCCCTCCACCCTGCCTCCTCGGGCTCCACTCGTC
TGCCCCTGGA CTCCCGTCTCCTCCTGTCTCCGGCTTCCCAGAGCTCCCTCCTTATGGCA
GCAGCTTCCCGCGTCTCCGGCGCAGCTTCTCAGCGGACGACCCTCTCGCTCCGGGGCTGA
GCCCAGTCCCTGGATGTTGCTGAAACTCTCGAGATCATGCGCGGGTTTGGCTGCTGCTTC
CCCGCCGGGTGCCACTGCCACCGCCGCCGCTCTGCTGCCGCCGTCCGCGGGATGCTCAG
TAGCCCGCTGCCCGGCCCCCGCGATCCTGTGTTCTCGGAAGCCGTTTGCTGCTGCAGAG
TTGCACGAACTAGTCATGTGCTGTGGGAGTCCCCGCGGCAGTGCAGCAGCTGGACACTT
TGCGAGGGCTTTTGCTGGCTGCTGCTGCTGCCCGTCATGCTACTCATCGTAGCCCGCCCG
GTGAAGCTCGCTGCTTTCCCTACCTCCTTAAGTGACTGCCAAACGCCCACCGGCTGGAAT
TGCTCTGTTTATGATGACAGAGAAAATGATCTCTTCCTCTGTGACACCAACACCTGTAAA
TTTGATGGGGAATGTTTAAGAATTGGAGACACTGTGACTTGCGTCTGTGAGTTCAAGTGC
AACAATGACTATGTGCCTGTGTGTGGCTCCAATGGGGAGAGCTACCAGAATGAGTGTTAC
CTGCGACAGGCTGCATGCAAACAGCAGAGTGAGATACTTGTGGTGTGAGAAGGATCATGT
GCCACAGATGCAGGATCAGGATCTGGAGATGGAGTCCATGAAGGCTCTGGAGAACTAGT
CAAAGGAGACATCCACCTGTGATATTTGCCAGTTTGGTGCAGAATGTGACGAAGATGCC
GAGGATGTCTGGTGTGTGTGTAATATTGACTGTTCTCAAACCAACTTCAATCCCCTCTGC
GCTTCTGATGGGAAATCTTATGATAATGCATGCCAAATCAAAGAAGCATCGTGTGAGAAA
CAGGAGAAAATTGAAGTCATGTCTTTGGGTCGATGTCAAGATAACACAACACTACAACACT
AAGTCTGAAGATGGGCATTATGCAAGAACAGATTATGCAGAGAATGCTAACAAATTAGAA
GAAAGTGCCAGAGAACACCACATACCTTGTCCGGAACATTACAATGGCTTCTGCATGCAT
GGGAAGTGTGAGCATTCTATCAATATGCAGGAGCCATCTTGCAGGTGTGATGCTGGTTAT
ACTGGACAACACTGTGAAAAAAGGACTACAGTGTTCTATACGTTGTTCCCGGTCCTGTA
CGATTTTCAGTATGTCTTAATCGCAGCTGTGATTGGAACAATTCAGATTGCTGTCATCTGT
GTGGTGGTCCTCTGCATCACAAGGAAATGCCCCAGAAGCAACAGAATTCACAGACAGAAG
CAAATAACAGGGCACTACAGTTCAGACAATAACAAGAGCGTCCACGAGGTTAATCTAA
AGGGAGCATGTTTCACAGTGGCTGGACTACCGAGAGCTTGGACTACACAATACAGTATTA
TAGACAAAAGAATAAGACAAGAGATCTACACATGTTGCCTTGCAATTTGTGGTAATCTACA
CCAATGAAAACATGTACTACAGCTATATTTGATTATGTATGGATATATTTGAAATAGTAT
ACATTGTCTTGATGTTTTTTCTGTAATGTAAATAAATACTATTTATATCACACAATATAGTT
TTTTCTTTCCCATGTATTTGTTATATATAATAAATACTCAGTGATGAG

14/392

FIGURE 14

MVLWESPRQCSSWTLCEGFCWLLLLPVMLLIVARPVKLAAFPTSLSDCQTPTGWNC SGY
DDRENDLFLCDTNTCKFDGECLRIGDTVTCVCQFKCNNDYVPVCGSNGESYQNECYLRQ
AACKQQSEILVVSEGSCATDAGSGSGDGVHEGSGETSQKETSTCDICQFGAECDEDAED
VWCVCNIDCSQTNFNPLCASDGKSYDNACQIKEASCQKQEKIEVMSLGRCQDNTTTT TK
SEDGHYARTDYAENANKLEESAREHHIPCPEHYNGFCMHGKCEHSINMQEPSCRC DAGY
TGQHCEKKDYSVLYVVPGPVRFQYVLIAAVIGTIQIAVICVVVLCITRK CPRSNRIHRQ
KQNTGHYSSDNTTRASTRLI

15/392

FIGURE 15

GGAGCTCAGCCCAGTGGGCAGTCTGAAGATGGCCAATTACACGCTGGCACCAGAGGATGA
ATATGATGTCCTCATAGAAGGTGAACTGGAGAGCGATGAGGCAGAGCAATGTGACAAGTA
TGACGCCCAGGCACCTCTCAGCCCAGCTGGTGCCATCACTCTGCTCTGCTGTGTTTGTGAT
CGGTGTCCTGGACAATCTCCTGGTTGTGCTTATCCTGGTAAATATAAAGGACTCAAACG
CGTGGAAAATATCTATCTTCTAAACTTTGGCAGTTTCTAACTTGTGTTTCTTGCTTACCCT
GCCCTTCTGGGCTCATGCTGGGGGCGATCCCATGTGTAAATCTCATTGGACTGTACTT
CGTGGGCCTGTACAGTGAGACATTTTTCAATTGCCTTCTGACTGTGCAAAGGTACCTAGT
GTTTTTGCACAAGGGCAACTTTTTCTCAGCCAGGAGGAGGGTGCCCTGTGGCATCATTAC
AAGTGTCTTGGCATGGGTAAACAGCCATTCTGGCCACTTTGCCTGAATACGTGGTTTATAA
ACCTCAGATGGAAGACCAGAAATACAAGTGTGCATTTAGCAGAACTCCCTTCTGCCAGC
TGATGAGACATTCTGGAAGCATTTTCTGACTTTAAAAATGAACATTTTCGGTTCTTGTCTT
CCCCCTATTTATTTTTACATTTCTCTATGTGCAAATGAGAAAAACACTAAGGTTTCAGGGA
GCAGAGGTATAGCCTTTTCAAGCTTGTTTTTGCCATAATGGTAGTCTTCCTTCTGATGTG
GGCGCCCTACAATATTGCATTTTTCTGTCCACTTTCAAAGAACACTTCTCCCTGAGTGA
CTGCAAGAGCAGCTACAATCTGGACAAAAGTGTTACATCACTAACTCATCGCCACCAC
CCTGCTGCATCAACCCTCTCCTGTATGCGTTTCTTGATGGGACATTTAGCAAATACCT
CTGCCGCTGTTTCCATCTGCGTAGTAACACCCCACTTCAACCCAGGGGGCAGTCTGCACA
AGGCACATCGAGGGAAGAACCTGACCATTCCACCGAAGTGTA~~AA~~ACTAGCATCCACCAAAT
GCAAGAAGAATAAACATGGATTTTCATCTTTCTGCATTATTTTCATGTAAATTTTCTACAC
ATTTGTATACAAAATCGGATACAGGAAGAAAAGGGAGAGGTGAGCTAACATTTGCTAAGC
ACTGAATTTGTCTCAGGCACCGTGCAAGGCTCTTTACAAACGTGAGCTCCTTCGCCTCCT
ACCACTTGTCCATAGTGTGGATAGGACTAGTCTCATTTCTCTGAGAAGAAAATAAGGCG
CGGAAATTTGTCTAAGATCACATAACTAGGAAGTGGCAGAACTGATTCTCCAGCCCTGGT
AGCATTTGCTCAGAGCCTACGCTTGGTCCAGAACATCAAACCTCAAACCCTGGGGACAAA
CGACATGAAATAAATGTATTTTAAACATCTAAAA

16/392

FIGURE 16

MANYTLAPEDEYDVLIEGELESDEAEQCDKYDAQALSAQLVPSLCSAVFVIGVLDNLLVV
LILVKYKGLKRVENIYLLNLAVSNLCFLLTLPFWAHAGGDPCKILIGLYFVGLYSETFF
NCLLTVQRYLVFLHKGNFSSARRRVPCGIITSVLAWVTAILATLPEYVVYKPMEDQKYK
CAFSRTPFLPADETFWKHFLTLKMNISVLVLPLFI FTFLYVQMRKTLRFREQRYSLFKLV
FAIMVVFLLMWAPYNI AFFLSTFKEHFSLSDCSSYNLDKSVHITKLIATTHCCINPLLY
AFLDGTFSKYLRCRCFHLRSNTPLQPRGQSAQGTSREEPDHSTEV

Signal sequence:

None

Transmembrane domain:

41-61, 76-96, 109-129, 147-167, 199-219, 237-257, 285-305

7 transmembrane receptor (rhodopsin family):

55-300

N-glycosylation site:

3-6, 205-208

Tyrosine kinase phosphorylation site:

70-76, 171-179, 228-234

N-myristoylation site:

52-57, 136-141, 148-153

G-protein coupled receptors:

55-85, 96-136, 209-220, 235-254, 292-308

17/392

FIGURE 17

CGGACGCGTG GGGCGGACGCGTG GGGCGGCCACGGCGCCCGCGGGCTGGGGCGGTGCTTC
TTCCTTCTCCGTGGCCTACGAGGGTCCCCAGCCTGGGTAAAGATGGCCCCATGGCCCCCG
AAGGGCCTAGTCCCAGCTGTGCTCTGGGGCCTCAGCCTCTTCCTCAACCTCCCAGGACCT
ATCTGGCTCCAGCCCTCTCCACCTCCCCAGTCTTCTCCCCCGCCTCAGCCCCATCCGTGT
CATACCTGCCGGGGACTGGTTGACAGCTTTAACAAGGGCCTGGAGAGAACCATCCGGGAC
AACTTTGGAGGTGGAAACACTGCCTGGGAGGAAGAGAATTTGTCCAAATACAAAGACAGT
GAGACCCGCCTGGTAGAGGTGCTGGAGGGTGTGTGCAGCAAGTCAGACTTCGAGTGCCAC
CGCCTGCTGGAGCTGAGTGAGGAGCTGGTGGAGAGCTGGTGGTTTCACAAGCAGCAGGAG
GCCCCGGACCTCTTCCAGTGGCTGTGCTCAGATTCCCTGAAGCTCTGCTGCCCCGCAGGC
ACCTTCGGGGCCCTCCTGCCTTCCCTGTCTGGGGGAACAGAGAGGCCCTGCGGTGGCTAC
GGGCAGTGTGAAGGAGAAGGGACACGAGGGGGCAGCGGGCACTGTGACTGCCAAGCCGGC
TACGGGGGTGAGGCCTGTGGCCAGTGTGGCCTTGGCTACTTTGAGGCAGAACGCAACGCC
AGCCATCTGGTATGTTCCGCTTGTTTTGGCCCCCTGTGCCCCGATGCTCAGGACCTGAGGAA
TCAAACCTGTTTGCAATGCAAGAAGGGCTGGGCCCTGCATCACCTCAAGTGTGTAGACATT
GATGAGTGTGGCACAGAGGGAGCCAACTGTGGAGCTGACCAATTCTGCGTGAACACTGAG
GGCTCCTATGAGTGCCGAGACTGTGCCAAGGCCTGCCTAGGCTGCATGGGGGCAGGGCCA
GGTCGCTGTAAGAAGTGTAGCCCTGGCTATCAGCAGGTGGGCTCCAAGTGTCTCGATGTG
GATGAGTGTGAGACAGAGGTGTGTCCGGGAGAGAAACAAGCAGTGTGAAAACACCGAGGGC
GGTTATCGCTGCATCTGTGCCGAGGGCTACAAGCAGATGGAAAGGCATCTGTGTGAAGGAG
CAGATCCCAGAGTCAGCAGGCTTCTTCTCAGAGATGACAGAAGACGAGTTGGTGGTGCTG
CAGCAGATGTTCTTTGGCATCATCATCTGTGCACTGGCCACGCTGGCTGCTAAGGGCGAC
TTGGTGTTACACGCCATCTTCATTGGGGCTGTGGCGGCCATGACTGGCTACTGGTTGTCA
GAGCGCAGTGACCGTGTGCTGGAGGGCTTCATCAAGGGCAGATAATCGCGGCCACCACCT
GTAGGACCTCCTCCCACCCACGCTGCCCCCAGAGCTTGGGCTGCCCTCCTGCTGGACACT
CAGGACAGCTTGGTTTATTTTTGAGAGTGGGGTAAGCACCCCTACCTGCCTTACAGAGCA
GCCCAGGTACCCAGGCCCGGGCAGACAAGGCCCTGGGGTAAAAAGTAGCCCTGAAGGTG
GATACCATGAGCTCTTCACCTGGCGGGGACTGGCAGGCTTCACAATGTGTGAATTTCAA
AGTTTTTCTTAATGGTGGCTGCTAGAGCTTTGGCCCCCTGCTTAGGATTAGGTGGTCCTC
ACAGGGGTGGGGCCATCACAGCTCCCTCCTGCCAGCTGCATGCTGCCAGTTCCTGTTCTG
TGTTACACCATCCCCACCCCCATTGCCACTTATTTATTCATCTCAGGAAATAAAGAAA
GGTCTTGGAAGTTAAAAAAAAAAAAAAAAAAAAAAAAA

18/392

FIGURE 18

MAPWPPKGLVPAVLWGLSLFLNLP GPIWLQPSPPPQSSPPPQPHPCHTCRGLVDSFNKGL
ERTIRDNFGGGNTAWEENLSKYKDSETRLVEVLEGVCSKSDFECHRLLELSEELVESWW
FHKQOEAPDLFQWLCSDSLKLCCPAGTFGPSCLPCPGGTERPCGGYGQCEGEGTRGSGH
CDCQAGYGGEACGQCGLGYFEAERNASHLVCSACFGPCARCSGP EESNCLQCKKGWALHH
LKCVDIDECGTEGANCGADQFCVNTEGSYECRDCAKACLGCMGAGPGRCKKCSPGYQQVG
SKCLDVDECETEVCPGENKQCENTEGGYRCICAEGYKQMEGICVKEQIPESAGFFSEMTE
DELVLQQMFFGIIICALATLAAKGLVFTAIFIGAVAAMTGYWLSERSDRVLEGFIKGR

19/392

FIGURE 19

GCCCCGGGACTGGCGCAAGGTGCCCAAGCAAGGAAAGAAATAATGAAGAGACACATGTGT
AGCTGCAGCCTTTTGAAACACGCAAGAAGGAAATCAATAGTGTGGACAGGGCTGGAACCT
TTACCACGCTTGTTGGAGTAGATGAGGAATGGGCTCGTGATTATGCTGACATTCCAGCAT
GAATCTGGTAGACCTGTGGTTAACCCGTTCCCTCTCCATGTGTCTCCTCCTACAAAGTTT
TGTTCTTATGATACTGTGCTTTTCATTCTGCCAGTATGTGTCCCAAGGGCTGTCTTTGTTC
TTCCTCTGGGGGTTTAAATGTCACCTGTAGCAATGCAAATCTCAAGGAAATACCTAGAGA
TCTTCCTCCTGAAACAGTCTTACTGTATCTGGACTCCAATCAGATCACATCTATTCCCAA
TGAAATTTTAAAGGACCTCCATCAACTGAGAGTTCTCAACCTGTCCAAAAATGGCATTGA
GTTTATCGATGAGCATGCCTTCAAAGGAGTAGCTGAAACCTTGCAGACTCTGGACTTGTC
CGACAATCGGATTCAAAGTGTGCACAAAAATGCCTTCAATAACCTGAAGGCCAGGGCCAG
AATTGCCAACAACCCCTGGCACTGCGACTGTACTCTACAGCAAGTTCTGAGGAGCATGGC
GTCCAATCATGAGACAGCCACAACGTGATCTGTAAACGTCCGTGTTGGATGAACATGC
TGGCAGACCATTCTCAATGCTGCCAACGACGCTGACCTTTGTAACCTCCCTAAAAAAC
TACCGATTATGCCATGCTGGTCACCATGTTTGGCTGGTTCATATGGTGATCTCATATGT
GGTATATTATGTGAGGCAAAATCAGGAGGATGCCCGGAGACACCTCGAATACTTGAAATC
CCTGCCAAGCAGGCAGAAGAAAGCAGATGAACCTGATGATATTAGCACTGTGGTATAGTG
TCCAAACTGACTGTCATTGAGAAAAGAAAGAAAGTAGTTTGCGATTGCAGTAGAAATAAGT
GGTTTACTTCTCCCATCCATTGTAAACATTTGAAACTTTGTATTTTCAGTTTTTTTTTGAAT
TATGCCACTGCTGAACTTTTAACAAACACTACAACATAAATAATTTGAGTTTAGGTGATC
CACCCCTTAATTGTACCCCGATGGTATATTTCTGAGTAAGCTACTATCTGAACATTAGT
TAGATCCATCTCACATTTTAATAATGAAATTTATTTTTTTTAATTTAAAAGCAAATAAAAG
CTTAACCTTTGAACCATGGGAAAAAAAAAAAAAAAAAAAAAAAAAACA

20/392

FIGURE 20

MNLVDLWLTRSLSMCLLLQSFVLMILCFHSASMCPCGCLCSSSGGLNVTCSNANLKEIPR
DLPPETVLLYLDNQITSIPNEIFKDLHQLRVNLNSKNGIEFIDEHAFKGVAETLQTLDL
SDNRIQSVHKNAFNNLKRARIANNPWHCDCTLQQVLRSMASNHETAHNVICKTSVLDEH
AGRPFLNAANDADLCNLPKTTDYAMLVTMFGWFTMVISYVVYYVRQNQEDARRHLEYLK
SLPSRQKKADEPDDISTVV

Signal sequence:

amino acids 1-33

Transmembrane domain:

amino acids 205-220

N-glycosylation site:

amino acids 47-51, 94-98

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 199-203

Casein kinase II phosphorylation site:

amino acids 162-166, 175-179

N-myristoylation site:

amino acids 37-43, 45-51, 110-116

21/392

FIGURE 21

CGCCACCACTGCGGCCACCGCCAATGAAACGCCTCCCGCTCCTAGTGGTTTTTTTCCACTT
TGTTGAATTGTTCTTATACTCAAAATTGCACCAAGACACCTTGTCTCCCAAATGCAAAAT
GTGAAATACGCAATGGAATTGAAGCCTGCTATTGCAACATGGGATTTTCAGGAAATGGTG
TCACAATTTGTGAAGATGATAATGAATGTGGAAATTTAACTCAGTCCTGTGGCGAAAATG
CTAATTGCACTAACACAGAAGGAAGTTATTATTGTATGTGTGTACCTGGCTTCAGATCCA
GCAGTAACCAAGACAGGTTTATCACTAATGATGGAACCGTCTGTATAGAAAATGTGAATG
CAAACCTGCCATTTAGATAATGTCTGTATAGCTGCAAATATTAATAAAACTTTAACAAAAA
TCAGATCCATAAAAGAACCCTGTGGCTTTGCTACAAGAAGTCTATAGAAATTCTGTGACAG
ATCTTTCACCAACAGATATAATTACATATATAGAAATATTAGCTGAATCATCTTCATTAC
TAGGTTACAAGAACAACACTATCTCAGCCAAGGACACCCTTTCTAACTCAACTCTTACTG
AATTTGTAAAAACCGTGAATAATTTTGTTCAAAGGGATACATTTGTAGTTTGGGACAAGT
TATCTGTGAATCATAGGAGAACACATCTTACAAAACCTCATGCACACTGTTGAACAAGCTA
CTTTAAGGATATCCCAGAGCTTCCAAAAGACCACAGAGTTTGATACAAATTCACCGGATA
TAGCTCTCAAAGTTTTCTTTTTTGATTATATAACATGAAACATATTCATCCTCATATGA
ATATGGATGGAGACTACATAAATATATTTCCAAAGAGAAAAGCTGCATATGATTCAAATG
GCAATGTTGCAGTTGCATTTTTATATTATAAGAGTATTGGTCCTTTGCTTTCATCATCTG
ACAACTTCTTATTGAAACCTCAAAATTATGATAATTCTGAAGAGGAGGAAAGAGTCATAT
CTTCAGTAATTTAGTCTCAATGAGCTCAAACCCACCCACATTATATGAACCTGAAAAAA
TAACATTTACATTAAGTCATCGAAAGGTCACAGATAGGTATAGGAGTCTATGTGCATTTT
GGAATTACTCACCTGATACCATGAATGGCAGCTGGTCTTCAGAGGGCTGTGAGCTGACAT
ACTCAAATGAGACCCACACCTCATGCCGCTGTAATCACCTGACACATTTTGCAATTTTGA
TGTCCTCTGGTCCTTCCATTGGTATTAAGATTATAATATTCTTACAAGGATCACTCAAC
TAGGAATAATTATTTCACTGATTGTCTTGCCATATGCATTTTTTACCTTCTGGTTCTTCA
GTGAAATTCAAAGCACCAGGACAACAATTCACAAAAATCTTTGCTGTAGCCTATTTCTTG
CTGAACTTGTTTTTCTTGTTGGGATCAATACAAATACTAATAAGCTCTTCTGTTCAATCA
TTGCCGGAAGTCTACACTACTTCTTTTTAGCTGCTTTTGCATGGATGTGCATTGAAGGCA
TACATCTCTATCTCATTGTTGTGGGTGTCATCTACAACAAGGGATTTTTTGCACAAGAATT
TTTATATCTTTGGCTATCTAAGCCCAGCCGTGGTAGTTGGATTTTCGGCAGCACTAGGAT
ACAGATATTATGGCACAACCAAAGTATGTTGGCTTAGCACCGAAAACAACCTTTATTTGGA
GTTTTATAGGACCAGCATGCCTAATCATTCCTTGTTAATCTCTTGGCTTTTGGAGTCATCA
TATACAAAGTTTTTTCGTCACACTGCAGGGTTGAAACCAGAAGTTAGTTGCTTTGAGAAAC
TAAGGTCTTGTCAGAGAGGAGCCCTCGCTCTTCTGTTCCCTTCTCGGCACCACCTGGATCT
TTGGGGTTCTCCATGTTGTGCACGCATCAGTGGTTACAGCTTACCTCTTCACAGTCAGCA
ATGCTTTCCAGGGGATGTTTCATTTTTTTTATTCTGTGTGTTTTATCTAGAAAGATTCAAG
AAGAATATTACAGATTGTTCAAAAATGTCCCCTGTTGTTTTGGATGTTTAAGGTAAACAT
AGAGAATGGTGGATAATTACAACCTGCACAAAAATAAAAATTCCAAGCTGTGGATGACCAA
TGTATAAAAATGACTCATCAAATTATCCAATTATTAACCTACTAGACAAAAAGTATTTTAA
ATCAGTTTTTCTGTTTATGCTATAGGAACGTAGATAATAAGGTAAAATTATGTATCATA
TAGATATACTATGTTTTTCTATGTGAAATAGTTCTGTCAAAAATAGTATTGCAGATATTT
GGAAAGTAATTGGTTTTCTCAGGAGTGATATCACTGCACCCAAGGAAAGATTTTCTTTCTA
ACACGAGAAGTATATGAATGTCCTGAAGGAAACCACTGGCTTGATATTTCTGTGACTCGT
GTTGCCTTTGAACTAGTCCCCTACCACCTCGGTAATGAGCTCCATTACAGAAAGTGGAA
CATAAGAGAATGAAGGGGCAGAATATCAAACAGTGAAAAGGGAATGATAAGATGTATTTT
GAATGAACTGTTTTTTCTGTAGACTAGCTGAGAAATTGTTGACATAAAATAAAGAATTGA
AGAAACACATTTTACCATTTTGTGAATTGTTCTGAACTTAAATGTCCACTAAAACAACCTT
AGACTTCTGTTTGCTAAATCTGTTTCTTTTCTAATATTCTAAAAAAGGTTT
ACCTCCACAAATTGAAA

22/392

FIGURE 22

MKRLPLLVFSTLLNCSYTQNCTKTPCLPNAKCEIRNGIEACYCNMGFSGNGVTICEDDN
ECGNLTQSCGENANCTNTEGSYYCMCVPGFRSSSNQDRFITNDGTVCIENVNANCHLDNV
CIAANINKTLTKIRSIKEPVALLQEVYRNSVTDLSPTDIITYIEILAESSLLGYKNNTI
SAKDTLSNSTLTEFVKTVNNFVQRDTFVVWDKLSVNHRRTHLTCLMHTVEQATLRISQSF
QKTTEFDTNSTDIALKVFFFDSDYNMKHIHPHMNDGDYINIFPKRKAAYDSNGNVAVAF
YYKSIGPLLSSSDNFLKLPQNYDNSEEEERVISSVISVSMSSNPPTLYELEKITFTLSHR
KVTDRYRSLCAFWNYSPTMNGSWSSEGCETYSNETHTSCRCNHLTHFAILMSSGPSIG
IKDYNILTRITQLGIIISLICLAICIFTFWFFSEIQSTRTTIHKNLCCSLFLAELVFLVG
INTNTNKLFCSSIAGLLHYFFLAFAWMCIEGIHLYLIVVGVIYNKGFLHKNFYIFGYLS
PAVVVGFSAAALGYRYYGTTKVCWLSTENNFIWSFIGPACLIILVNLLAFGVIIYKVRHT
AGLKPEVSCFENIRSCARGALALLFLLGTTWIFGVLHVHASVVTAYLFTVSNAFQGMFI
FLFLCVLSRKIQEYYRLFKNPCCFGCLR

23/392

FIGURE 23

CTCCTCTTAACATACTTGCAGCTAAAACTAAATATTGCTGCTTGGGGACCTCCTTCTAGC
CTTAAATTTTTCAGCTCATCACCTTCACCTGCCTTGGTTCATGGCTCTGCTATTCTCCTTGAT
CCTTGCCATTTTGCACCAGACCTGGATTCCCTAGCGTCTCCATCTGGAGTGCGGCTGGTGGG
GGGCTCCACCGCTGTGAAGGGCGGGTGGAGGTGGAACAGAAAGGCCAGTGGGGCACCGT
GTGTGATGACGGCTGGGACATTAAGGACGTGGCTGTGTTGTGCCGGGAGCTGGGCTGTGG
AGCTGCCAGCGGAACCCCTAGTGGTATTTTGTATGAGCCACCAGCAGAAAAAGAGCAAAA
GGTCCTCATCCAATCAGTCAGTTGCACAGGAACAGAAGATACATTGGCTCAGTGTGAGCA
AGAAGAAGTTTATGATTGTTTACATGATGAAGATGCTGGGGCATCGTGTGAGAACCAGCA
GAGCTCTTTCTCCCCAGTCCCAGAGGGTGTGAGGCTGGCTGACGGCCCTGGGCATTGCAA
GGGACGCGTGGAAGTGAAGCACCAGAACAGTGGTATACCGTGTGCCAGACAGGCTGGAG
CCTCCGGGGCCGCAAAGGTGGTGTGCCGGCAGCTGGGATGTGGGAGGGCTGTACTGACTCA
AAAACGCTGCAACAAGCATGCCTATGGCCGAAAACCCATCTGGCTGAGCCAGATGTGATG
CTCAGGACGAGAAGCAACCCTTCAGGATTGCCCTTCTGGGCCTTGGGGGAAGAACACCTG
CAACCATGATGAAGACACGTGGGTGCAATGTGAAGATCCCTTTGACTTGAGACTAGTAGG
AGGAGACAACCTCTGCTCTGGGCGACTGGAGGTGCTGCACAAGGGCGTATGGGGCTCTGT
CTGTGATGACAACTGGGGAGAAAAGGAGGACCAGGTGGTATGCAAGCAACTGGGCTGTGG
GAAGTCCCTCTCTCCCTCCTTCAGAGACCGGAAATGCTATGGCCCTGGGGTTGGCCGCAT
CTGGCTGGATAATGTTTCGTTGCTCAGGGGAGGAGCAGTCCCTGGAGCAGTGCCAGCACAG
ATTTTGGGGGTTTCACGACTGCACCCACCAGGAAGATGTGGCTGTCATCTGCTCAGTGTA
GGTGGGCATCATCTAATCTGTTGAGTGCCTGAATAGAAGAAAAACACAGAAGAAGGGAGC
ATTTACTGTCTACATGACTGCATGGGATGAACACTGATCTTCTTCTGCCCTTGGACTGGG
ACTTATACTTGGTGCCCTGATTCTCAGGCCTTCAGAGTTGGATCAGAAGTTACAACATC
AGGTCTAGTTCTCAGGCCATCAGACATAGTTTGGAACTACATCACCACCTTTTCTATGTC
TCCACATTGCACACAGCAGATTCCCAGCCTCCATAATTGTGTGTATCAACTACTTAAATA
CATTCTCATACACCATTTGTCC
TGTTTCTCTGAAGAACTCTGACAAAATACAGATTTTGGTACTGAAAGAGATTCTAGAGGA
ACGGAATTTTAAGGATAAATTTTCTGAATTGGTTATGGGGTTTCTGAAATTGGCTCTATA
ATCTAATTAGATATAAAATTCTGGTAACTTTATTTACAATAATAAAGATAGCACTATGTG
TTCAAA

24/392

FIGURE 24

MALLFSLILAICTRPGFLASPSGVRLVGGLHRCEGRVEVEQKGQWGTVCDDGWDIKDVAV
LCRELGCGAASGTPSGILYEPPAEKEQKVLIQSVSCTGTEDTLAQCEQEVEYDCSHDEDA
GASCENPESSFSPVPEGVRLADGPGHCKGRVEVKHQNQWYTVQCQTGWSLRAAKVVCRQLG
CGRAVLTQKRCNKHAYGRKPIWLSQMSCSGREATLQDCPSGPWGKNTCNHDEDTWVECED
PFDLRLVGGDNLCSGRLEVLHKGWGSVCDDNWGEKEDQVVCKQLGCGKSLSPSFRDRKC
YGPVGRIWLDNVRCSGEEQSLEQCQHRFWGFHDCTHQEDVAVICSV

Signal sequence:

amino acids 1-15

Casein kinase II phosphorylation site:amino acids 47-51, 97-101, 115-119, 209-213, 214-218, 234-238,
267-271, 294-298, 316-320, 336-340**N-myristoylation site:**amino acids 29-35, 43-49, 66-72, 68-74, 72-78, 98-104, 137-143,
180-186, 263-269, 286-292**Amidation site:**

amino acids 196-200

Speract receptor repeated domain signature:

amino acids 29-67, 249-287

25/392

FIGURE 25

CGGACGCGTGCGGCGTCCGGCGGTGCGCAGAGCCAGGAGGCGGAGGCGCGCGGGCCAGCCTG
GGCCCCAGCCCACACCTTCACCAGGGGCCAGGAGCCACCATGTGCGCATGTCCACTGGGG
CTACTGCTGTTGCTGCCGCTGGCTGGCCACTTGGCTCTGGGTGCCCAGCAGGGTCGTGGG
CGCCGGGAGCTAGCACCGGGTCTGCACCTGCCGGGCATCCGGGACGCGGGAGGCCGGTAC
TGCCAGGAGCAGGACCTGTGCTGCCGCGGGCGTCCGACGACTGTGCCCTGCCCTACCTG
GGCGCCATCTGTTACTGTGACCTCTTCTGCAACCGCACGGTCTCCGACTGCTGCCCTGAC
TTCTGGGACTTCTGCCTCGGCGTGCCACCCCTTTTCCCCCGATCCAAGGATGTATGCAT
GGAGGTCGTATCTATCCAGTCTTGGGAACGTACTGGGACAACGTAAACCGTTGCACCTGC
CAGGAGAACAGGCAGTGGCATGGTGGATCCAGACATGATCAAAGCCATCAACCAGGGCAA
CTATGGCTGGCAGGCTGGGAACCACAGCGCTTCTGGGGCATGACCCTGGATGAGGGCAT
TCGCTACCGCTGGGCACCATCCGCCATCTTCCTCGGTCATGAACATGCATGAAATTTA
TACAGTGCTGAACCCAGGGGAGGTGCTTCCCACAGCCTTCGAGGCCTCTGAGAAGTGGCC
CAACCTGATTCATGAGCCTCTTGACCAAGGCAACTGTGCAGGCTCCTGGGCCTTCTCCAC
AGCAGCTGTGGCATCCGATCGTGTCTCAATCCATTCTCTGGGACACATGACGCCTGTCT
GTCGCCCCAGAACCTGCTGTCTTGACACCCACCAGCAGCAGGGCTGCCGCGGTGGGCG
TCTCGATGGTGCTTGGTGGTTCCTGCGTCGCCGAGGGGTGGTGTCTGACCACTGCTACCC
CTTCTCGGGCCGTGAACGAGACGAGGCTGGCCCTGCGCCCCCTGTATGATGCACAGCCG
AGCCATGGGTGCGGGCAAGCGCCAGGCCACTGCCCACTGCCCAACAGCTATGTTAATAA
CAATGACATCTACCAGGTCACTCCTGTCTACCGCCTCGGCTCCAACGACAAGGAGATCAT
GAAGGAGCTGATGGAGAATGGCCCTGTCCAAGCCCTCATGGAGGTGCATGAGGACTTCTT
CCTATAACAAGGAGGCATCTACAGCCACACGCCAGTGAGCCTTGGGAGGCCAGAGAGATA
CCGCCGGCATGGGACCCACTCAGTCAAGATCACAGGATGGGGAGAGGAGACGCTGCCAGA
TGGAAGGACGCTCAAATACTGGACTGCGGCCAACTCCTGGGGCCAGCCTGGGGCGAGAG
GGGCCACTTCCGCATCGTGCGCGGCGTCAATGAGTGCGACATCGAGAGCTTCGTGCTGGG
CGTCTGGGGCCGCGTGGGCATGGAGGACATGGGTCACTGAGGCTGCGGGCACCACGC
GGGGTCCGGCCTGGGATCCAGGCTAAGGGCCGGCGGAAGAGGCCCAATGGGGCGGTGAC
CCCAGCCTCGCCCGACAGAGCCCGGGCGCAGGCGGGCGCCAGGGCGCTAATCCCGGCGC
GGGTTCGCTGACGCGAGCGCCCCGCTGGGAGCCGCGGGCAGGCGAGACTGGCGGAGCCC
CCAGACCTCCCAGTGGGGACGGGGCAGGGCCTGGCCTGGGAAGAGCACAGCTGCAGATCC
CAGGCCTCTGGCGCCCCCACTCAAGACTACCAAAGCCAGGACACCTCAAGTCTCCAGCCC
CAATACCCCAACCCCAATCCCGTATTCTTTTTTTTTTTTTTTTTTTAGACAGGGTCTTGCTCCG
TTGCCAGGTTGGAGTGAGTGAGTGGCCCATCAGGGCTCACTGTAACCTCCGACTCCTGGGTT
CAAGTGACCCTCCACCTCAGCCTCTCAAGTAGCTGGGACTACAGGTGCACCACCACACC
TGGCTAATTTTTGTATTTTTTGTAAAGAGGGGGTCTCACTGTGTTGCCCAGGCTGGTTT
CGAACTCCTGGGCTCAAGCGGTCCACCTGCCTCCGCCTCCCAAAGTGCTGGGATTGCAGG
CATGAGCCACTGCACCCAGCCCTGTATTCTTATTCTTCAGATATTTATTTTTCTTTTAC
TGTTTTAAATAAAACCAAAGTATTGATAAAAAAAA

26/392

FIGURE 26

MWRCPLGLLLLLPLAGHLALGAQQGRGRRELAPGLHLRGIRDAGGRYCQEQLCCRGRAD
DCALPYLGAIICYCDLFCNRTVSDCCPDFWDFCLGVPPFPPIQGCMHGGRIYPVLGTYWD
NCNRCTCQENRQWHGGSRHDQSHQPGQLWLAGWEPQRLLGHDPG

N-glycosylation site:

amino acids 78-82, 161-165

Casein kinase II phosphorylation site:

amino acids 80-84, 117-121, 126-130, 169-173, 205-209, 296-300,
411-415

N-myristoylation site:

amino acids 21-27, 39-45, 44-50, 104-110, 160-164, 224-230,
269-275, 378-384, 442-448

Amidation site:

amino acids 26-30, 318-322

Eukaryotic thiol (cysteine) proteases histidine active site:

amino acids 398-409

27/392

FIGURE 27

CCCACGCGTCCGGCAGGTTTTCTTCAAGCCAAGAAGGACACGGATTGGCTGAAGGAGAA
AGTGCAGAGCTTGCAGACACTGGCTGCCAACAACCTCTGCGTTGGCCAAAGCCAACAACGA
CACCTTGGAGGATATGAACAGCCAGCTCAACTCATTACAGGTCAGATGGAGAACATCAC
CACTATCTCTCAAGCCAACGAGCAGAACCTGAAAGACCTGCAGGACTTACACAAAGATGC
AGAGAATAGAACAGCCATCAAGTTCAACCAACTGGAGGAACGCTTCCAGCTCTTTGAGAC
GGATATTGTGAACATCATTAGCAATATCAGTTACACAGCCCACCACCTGCGGACGCTGAC
CAGCAATCTAAATGAAGTCAGGACCACTTGCACAGATACCTTACCAAACACACAGATGAT
CTGACCTCCTTGAATAATACCCTGGCCAACATCCGTTTGGATTCTGTTTCTCTCAGGATG
CAACAAGATTTGATGAGGTCGAGGTTAGACACTGAAGTAGCCAACCTTATCAGTGATTATG
GAAGAAATGAAGCTAGTAGACTCCAAGCATGGTCAGCTCATCAAGAATTTTACAATACTA
CAAGGTCCACCGGGCCCCAGGGGTCCAAGAGGTGACAGAGGATCCCAGGGACCCCCTGGC
CCAAGTGGCAACAAGGGACAGAAAGGAGAGAAGGGGGAGCCTGGACCACCTGGCCCTGCG
GGTGAGAGAGGCCCAATTGGACCAGCTGGTCCCCCGGAGAGCGTGGCGGCAAAGGATCT
AAAGGCTCCCAGGGCCCCAAAGGCTCCCGTGGTTCCCCTGGGAAGCCCGGCCCTCAGGGC
CCCAGTGGGGACCCAGGCCCCCCGGGGCCACCAGGCAAAGAGGGACTCCCCGGCCCTCAG
GGCCCTCCTGGCTTCCAGGGACTTCAGGGCACCGTTGGGGAGCCTGGGGTGCCTGGACCT
CGGGGACTGCCAGGCTTGCTTGGGGTACCAGGCATGCCAGGCCCAAGGGCCCCCCCCGGC
CCTCCTGGCCCATCAGGAGCGGTGGTGGCCCTGGCCCTGCAGAATGAGCCAACCCCGGCA
CCGGAGGACAATAGCTGCCCGCCTCACTGGAAGAACTTCACAGACAAATGCTACTATTTT
TCAGTTGAGAAAGAAATTTTTGAGGATGCAAAGCTTTTCTGTGAAGACAAGTCTTCACAT
CTTGTTTTTCATAAACACTAGAGAGGAACAGCAATGGATAAAAAAACAGATGGTAGGGAGA
GAGAGCCACTGGATCGGCCTCACAGACTCAGAGCGTGAAAATGAATGGAAGTGGCTGGAT
GGGACATCTCCAGACTACAAAAATTGGAAAGCTGGACAGCCGGATAACTGGGGTCATGGC
CATGGGCCAGGAGAAGACTGTGCTGGGTTGATTTATGCTGGGCAGTGGAACGATTTCCAA
TGTGAAGACGTCAATAACTTCATTTGCGAAAAAGACAGGGAGACAGTACTGTCATCTGCA
TTATAACGGACTGTGATGGGATCACATGAGCAAATTTTCAGCTCTCAAAGGCAAAGGACA
CTCCTTTCTAATTGCATCACCTTCTCATCAGATTGAAAAAAAAAAAAAGCACTGAAAACCAA
TTACTGAAAAAAAAAATTGACAGCTAGTGTTTTTTTACCATCCGTCATTACCCAAAGACTTGG
GAACTAAATGTTCCCCAGGGTGATATGCTGATTTTCATTGTGCACATGGACTGAATCAC
ATAGATTCTCCTCCGTGAGTAACCGTGCGATTATACAAATTATGTCTTCCAAAGTATGGA
ACACTCCAATCAGAAAAAGGTTATCATTGGTCGTTGAGTTATGGGAAGAACTTAAGCATA
TACTGTGTAAACAGTGCCATACATTTCTAAAATCCCAAGTGTAGGAAAAATATGCAGACA
TACAGATATATAGGCCAACTATTAGTAATAATATGAAATATACTTAAAGAGCTTTTAAAA
CTTTGTATTTTTGTACAAAAAAA

28/392

FIGURE 28

MQQDLMRSRLDTEVANLSVIMEEMKLVDSKHGQLIKNFTILQGPPGPRGPRGDRGSQGPP
GPTGNKGQKGEKGEPGPPGPAGERGPAGPPGERGGKGSKGSQGPKGSRGSPGKPGPQ
GPSGDPGPPGPPGKEGLPGPQGPPGFQGLQGTVEPGVPGPRGLPGLPGVPGMPGPKGPP
GPPGPPSGAVVPLALQNEPTPAPEDNSCPPHWKNFDTKCYFYSVEKEIFEDAKLFCEDKSS
HLVFINTREEQQWIKKQMVGREHIGLTDSERENKWLWDGTSPDYKNWKAGQPDNWDGH
GHGPGEDCAGLIYAGQWDFQCEDVNNFICEKDRETVLSSAL

Signal sequence:

None

Transmembrane domain:

None

N-glycosylation site:

16-19, 37-40, 213-216

Tyrosine kinase phosphorylation site:

212-220

N-myristoylation site:

97-102, 100-105, 148-153, 267-272, 293-298, 310-315

Cell attachment sequence:

51-53

C-type lectin domain signature:

308-330

Lectin C-type domain:

233-330

Collagen triple helix repeat:

43-102, 127-186

29/392

FIGURE 29

GGACTAATCTGTGGGAGCAGTTTATTCCAGTATCACCCAGGGTGCAGCCACACCAGGACT
GTGTTGAAGGGTGTGTTTTTTCTTTTAAATGTAATACCTCCTCATCTTTTCTTCTTACAC
AGTGTCTGAGAACATTTACATTATAGATAAGTAGTACATGGTGGATAACTTCTACTTTTA
GGAGGACTACTCTCTTCTGACAGTCCTAGACTGGTCTTCTACACTAAGACACCCATGAAGG
AGTATGTGCTCCTATTATTCTGGCTTTGTGCTCTGCCAAACCCCTTCTTTAGCCCTTCAC
ACATCGCACTGAAGAATATGATGCTGAAGGATATGGAAGACACAGATGATGATGATGATG
ATGATGATGATGATGATGATGATGAGGACAACCTCTCTTTTCCAACAAGAGAGCCAAGAA
GCCATTTTTTTCCATTTGATCTGTTTCCAATGTGTCCATTTGGATGTCAGTGCTATTTCAC
GAGTTGTACATTGCTCAGATTTAGGTTTGACCTCAGTCCCAACCAACATTCCATTTGATA
CTCGAATGCTTGATCTTCAAAACAATAAAATTAAGGAAATCAAAGAAAATGATTTTAAAG
GACTCACTTCACTTTATGGTCTGATCCTGAACAACAACAAGCTAACGAAGATTACCCAA
AAGCCTTTCTAACCACAAAGAAGTTGCGAAGGCTGTATCTGTCCCACAATCAACTAAGTG
AAATACCACTTAATCTTCCCAAATCATTAGCAGAACTCAGAATTCATGAAAATAAAGTTA
AGAAAATACAAAAGGACACATTCAAAGGAATGAATGCTTTACACGTTTTGGAAATGAGTG
CAAACCCCTCTTGATAATAATGGGATAGAGCCAGGGGCATTTGAAGGGGTGACGGTGTTCC
ATATCAGAATTGCAGAAGCAAACTGACCTCAGTTCCTAAAGGCTTACCACCAACTTTAT
TGGAGCTTCACTTAGATTATAATAAAATTTCAACAGTGGAACCTTGAGGATTTTAAACGAT
ACAAAGAACTACAAAGGCTGGGCCTAGGAAACAACAAATCACAGATATCGAAAATGGGA
GTCTTGCTAACATAACCACGTGTGAGAGAAATACATTTGGAAAACAATAAACTAAAAA
TCCCTTCAGGATTACCAGAGTTGAAATACCTCCAGATAATCTTCCTTCATTCTAATTCAA
TTGCAAGAGTGGGAGTAAATGACTTCTGTCCAACAGTGCCAAAGATGAAGAAATCTTTAT
ACAGTGCAATAAGTTTATTCAACAACCCGGTGAAATACTGGGAAATGCAACCTGCAACAT
TTCGTTGTGTTTTGAGCAGAATGAGTGTTAGCTTGGGAACCTTTGGAATGTAATAATTAG
TAATTGGTAATGTCCATTTAATATAAGATTCAAAAATCCCTACATTTGGAATACTTGAAC
TCTATTAATAATGGTAGTATTATATATACAAGCAAATATCTATTCTCAAGTGGAAGTCC
ACTGACTTATTTTATGACAAGAAATTTCAACGGAATTTTGCCAAACTATTGATACATAAG
GGGTTGAGAGAAACAAGCATCTATTGCAGTTTCCTTTTTTGCGTACAAATGATCTTACATA
AATCTCATGCTTGACCATTCCTTTCTTCATAACAAAAAGTAAGATATTCGGTATTTAAC
ACTTTGTTATCAAGCACATTTTAAAAAGAACTGTACTGTAAATGGAATGCTTGACTTAGC
AAAATTTGTGCTCTTTTCATTTGCTGTTAGAAAAACAGAATTAACAAAGACAGTAATGTGA
AGAGTGCAATTACACTATTCTTATTCTTTAGTAACTTGGGTAGTACTGTAATATTTTTTAAT
CATCTTAAAGTATGATTTGATATAATCTTATTGAAATTACCTTATCATGTCTTAGAGCCC
GTCTTTATGTTTAAACTAATTTCTTAAATAAAGCCTTCAGTAAATGTTTATTACCAAC
TTGATAAATGCTACTCATAAGAGCTGGTTTGGGGCTATAGCATATGCTTTTTTTTTTTTA
ATTATTACCTGATTTAAAAATCTCTGTAAAAACGTGTAGTGTTTCATAAAATCTGTAACT
CGCATTTTAATGATCCGCTATTATAAGCTTTTAATAGCATGAAAATTGTTAGGCTATATA
ACATTGCCACTTCAACTCTAAGGAATATTTTTGAGATATCCCTTTGGAAGACCTTGCTTG
GAAGAGCCTGGACACTAACAATCTACACCAAATTTGTCTCTTCAAATACGTATGGACTGG
ATAACTCTGAGAAACACATCTAGTATAACTGAATAAGCAGAGCATCAAATTAAACAGACA
GAAACCGAAAGCTCTATATAAATGCTCAGAGTTCTTTATGTATTTCTTATTGGCATTCAA
CATATGTAAATCAGAAAACAGGGAAATTTTCATTAATAATATTGGTTTGAAAT

30/392

FIGURE 30

MKEYVLLLFLALCSAKPFFSPSHIALKNMMLKDMEDTDDDDDDDDDDDDDEDNSLFPTRE
PRSHFFPFDLFPMPFCQCYSRVVHCSDLGLTSVPTNIPFDTRMLDLQNNKIKEIKEND
FKGLTSLYGLILNNNKLTKIHPKAFLTTKKLRRLYLSHNQLSEIPLNLPKSLAELRIHEN
KVKKIQKDTFKGMNALHVLEMSANPLDNNGIEPGAEGVTVFHIRIAEAKLTSVPKGLPP
TLLELHLDYNKISTVELEDFKRYKELQRLGLGNNKITDIENGLANI PRVREIHLENNKL
KKIPSGLPKYLQIIFLHSNSIARVGVNDFCPTVPKMKKSLYSAISLFNNPVKYWEMQP
ATFRCVLSRMSVQLGNFGM

Signal sequence:
amino acids 1-15

N-glycosylation site:
amino acids 281-285

N-myristoylation sites:
amino acids 129-135, 210-216, 214-220, 237-243, 270-276,
282-288

Leucine zipper pattern:
amino acids 154-176

31/392

FIGURE 31

AGCAGGGAAATCCGGATGTCTCGGTTATGAAGTGGAGCAGTGAGTGTGAGCCTCAACATA
GTTCCAGAACTCTCCATCCGGACTAGTTATTGAGCATCTGCCTCTCATATCACCAGTGGC
CATCTGAGGTGTTTTCCCTGGCTCTGAAGGGGTAGGCACGATGGCCAGGTGCTTCAGCCTG
GTGTTGCTTCTCACTTCCATCTGGACCACGAGGCTCCTGGTCCAAGGCTCTTTGCGTGCA
GAAGAGCTTTCCATCCAGGTGTCTATGCAGAATTATGGGGATCACCTTGTGAGCAAAAAG
GCGAACCAGCAGCTGAATTTACAGAAGCTAAGGAGGCCTGTAGGCTGCTGGGACTAAGT
TTGGCCGGCAAGGACCAAGTTGAAACAGCCTTGAAAGCTAGCTTTGAAACTTGCAGCTAT
GGCTGGGTTGGAGATGGATTTCGTGGTTCATCTCTAGGATTAGCCCAAACCCCAAGTGTGGG
AAAAATGGGGTGGGTGTCTGATTGGAAGGTTCCAGTGAGCCGACAGTTTGCAGCCTAT
TGTTACAACCTCATCTGATACTTGGACTAACTCGTGCATTCCAGAAATTATCACCACCAA
GATCCCATATTCAACACTCAAACGCAACACAAACAGAAATTTATTGTGAGTGACAGT
ACCTACTCGGTGGCATCCCCTTACTCTACAATACCTGCCCCCTACTACTCTCCTCTGCT
CCAGCTTCCACTTCTATTCCACGGAGAAAAAATTGATTTGTGTACAGAAAGTTTTTATG
GAACTAGCACCATGTCTACAGAACTGAACCATTTGTTGAAAATAAAGCAGCATTCAAG
AATGAAGCTGCTGGGTTTGGAGGTGTCCCCACGGCTCTGCTAGTGCTTGCTCTCCTCTTC
TTTGGTGCTGCAGCTGGTCTTGGATTTTGCTATGTCAAAGGTATGTGAAGGCCTTCCCT
TTTACAAACAAGAATCAGCAGAAGGAAATGATCGAAACCAAAGTAGTAAAGGAGGAGAAG
GCCAATGATAGCAACCCTAATGAGGAATCAAAGAAAACCTGATAAAAACCCAGAAGAGTCC
AAGAGTCCAAGCAAAACTACCGTGCGATGCCTGGAAGCTGAAGTTTAGATGAGACAGAAA
TGAGGAGACACACCTGAGGCTGGTTTTCTTTCATGCTCCTTACCCTGCCCCAGCTGGGGAA
ATCAAAAGGGCCAAAGAACCAGAAAGAAAGTCCACCCTTGGTTCCCTAACTGGAATCAGC
TCAGGACTGCCATTGGACTATGGAGTGCACCAAGAGAATGCCCTTCTCCTTATTGTAAC
CCTGTCTGGATCCTATCCTCCTACCTCCAAAGCTTCCACGGCCTTTCTAGCCTGGCTAT
GTCCTAATAATATCCCACTGGGAGAAAGGAGTTTTGCAAAGTGCAAGGACCTAAACATC
TCATCAGTATCCAGTGGTAAAGGCCTCCTGGCTGTCTGAGGCTAGGTGGGTTGAAAGC
CAAGGAGTCACTGAGACCAAGGCTTTCTCTACTGATTCCGCAGCTCAGACCCCTTTCTTCA
GCTCTGAAAGAGAAACACGTATCCACCTGACATGTCTTCTGAGCCCGGTAAGAGCAAA
AGAATGGCAGAAAAGTTTAGCCCCTGAAAGCCATGGAGATTCTCATAACTTGAGACCTAA
TCTCTGTAAAGCTAAAATAAAGAAATAGAACAAGGCTGAGGATACGACAGTACACTGTCA
GCAGGGACTGTAAACACAGACAGGGTCAAAGTGTTTTCTCTGAACACATTGAGTTGGAAT
CACTGTTTAGAACACACACACTTACTTTTTCTGGTCTCTACCACTGCTGATATTTTCTCT
AGGAAATATACTTTTACAAGTAACAAAAATAAAAACTCTTATAAATTTCTATTTTTTATCT
GAGTTACAGAAATGATTACTAAGGAAGATTACTCAGTAATTTGTTTAAAAAGTAATAAAA
TTCAACAAACATTTGCTGAATAGCTACTATATGTCAAGTGCTGTGCAAGGTATTACACTC
TGTAATTGAATATTATTCTCAAAAAATTGCACATAGTAGAACGCTATCTGGGAAGCTAT
TTTTTTCAGTTTTGATATTTCTAGCTTATCTACTTCCAACTAATTTTTTATTTTTGCTGA
GACTAATCTTATTCATTTTCTCTAATATGGCAACCATTATAACCTTAATTTATTATTAAC
ATACCTAAGAAGTACATTGTTACCTCTATATACCAAAGCACATTTTAAAAGTGCCATTAA
CAAATGTATCACTAGCCCTCCTTTTTTCCAACAAGAGGGACTGAGAGATGCAGAAATATT
TGTGACAAAAAATTAAAGCATTTAGAAAACCTT

32/392

FIGURE 32

MARCFSLVLLLLTSIWTTTRLLVQGSLRAEELSIQVSCRIMGITLVSKKANQQQLNFTEAKEA
CRLGLSLAGKDQVETALKASFETCSYGWVGDFVVISRISPNPKCGKNGVGVLWKVPV
SRQFAAYCYNSSDTWTNSCIPEIITTKDPIFNTQTATQTTEFIVSDSTYSVASPYSTIPA
PTTTPPAPASTSIPRRKKLICVTEVFMETSTMSTETEPFVENKAAFKNAAAGFGGVPTAL
LVLALLFFGAAAGLGFCYVKRYVKAFPFTNKNQQKEMIETKVVKEEKANDSNPNEESKKT
DKNPEESKSPSKTTVRCLEAEV

Signal sequence:

amino acids 1-16

Transmembrane domain:

amino acids 235-254

N-glycosylation site:

amino acids 53-57, 130-134, 289-293

Casein kinase II phosphorylation site:

amino acids 145-149, 214-218

Tyrosine kinase phosphorylation site:

amino acids 79-88

N-myristoylation site:

amino acids 23-29, 65-71, 234-240, 235-239, 249-255, 253-259

33/392

FIGURE 33

[illegible]

34/392

TCTACTTATGTTGGACACTTGGCAGAAGGACCGTGCCCGGCGGCCTCATTTTGACCAGCT
GGTGGCTGCATTTGACAAGATGATCCGCAAGCCAGATACCCTGCAGGCTGGCGGGGACCC
AGGGGAAAGGCCTTCCCAGGCCCTTCTGACCCCTGTGGCCCTGGACTTTCCTTGTCTGGA
CTCACCCCAGGCCTGGCTTTCAGCCATTGGACTGGAGTGCTACCAGGACAACTTCTCCAA
GTTTGGCCTCTGTACCTTCAGTGATGTGGCTCAGCTCAGCCTAGAAGACCTGCCTGCCCT
GGGCATCACCTTGGCTGGCCACCAGAAGAAGCTGCTGCACCACATCCAGCTCCTTCAGCA
ACACCTGAGGCAGCAGGGCTCAGTGGAGGTCTGAGAATGACGATACCCGTGACTCAGCCC
TGGACACTGGTCCGAGAAGGGACATGTGGGACGTGAGCCGGGCTCCAACAGCCTCTGTGA
GAGATGCCCCACACCAAACCCAACCCTCCGATGGCTGCATTCCCTGGTCTCCGCTTTTC
CACCAGCCCCCTCCTCATTAAGGGAAAGAAGGGAATTTGCAAAAAAAAAAAAAAAAAAAAA
AAAAAA

35/392

FIGURE 34

MATEGAAQLGNRVAGMVC SLWVLLLVSSVLAL EEVLLD TTGETSEIGWLTYP PGGWDEV S
VLDDQRRLTRTFEACHVAGAPP GTGQDNWLQ THFVERRGAQRAHIRLHFSVRACSSSLGVS
GGTCRETFTLYYRQAE EPDSPDSVSSWHLKRWTKVD TIAADESFPSSSSSSSSSSSSSSA AW
AVGPHGAGQ RAGLQLNVKERSFGPLTQRGFYVAFQDTGAC LALVAVRLFSYTCPAVLRSF
ASFPETQASGAGGASLVA AVGTCVAHAEP EEDGVGGQAGGSP PRLHCN GEGKWMVAVGGC
RCQPGYQPARGDKACQACPRGLYKASAGNAPCSPCPARSHAPNPAAPVCPCLEGFYRASS
DPPEAPCTGPPSAPQELWFEVQGSALMLHWRLPRELGGRGDLLFN VVCKECEGRQEPASG
GGGTCHRCRDEVHFDPRQRGLTESRVLVGGLRAHVPIILEVQAVNGVSELSPDPPQAAAI
NVSTSHEVPSAVPVVHQVSRASNSITVSWPQPDQTNGNILDYQLRYDQAEDESHSFTLT
SETNTATVTQLSPGHIYGFQVRARTAAGHGPGYGGKVYFQTL PQGELSSQLPERLSLVIGS
TLGALAFLLLLAAITVLAVVFQRKRRTGYTEQLQQYSSPGLGVKYYIDPSTYEDPCQAIR
ELAREVDPAYIKIEEVIGTGSFGEVRQGR LQPRGRREQTVAIQALWAGGAESLQMTFLGR
AAVLGQFQHPNILRLEGVVTKSRLMLVLT EFMELGPLDSFLRQREGQFSSSLQLVAMQRGV
AAAMQYLSSFAFVHRSLSAHSVLVNSHLVCKVARLGHSPQGPSCLLRWAAPEVIAHGKHT
HVGSDDELWRTALLGHE

signal sequence:

Amino acids 1-31

Transmembrane domains:

Amino acids 217-234;598-618

N-glycosylation site:

Amino acids 481-485

Glycosaminoglycan attachment sites:

Amino acids 249-253;419-423

cAMP- and cGMP-dependent protein kinase phosphorylation sites:

Amino acids 66-70;150-154;624-628

Tyrosine kinase phosphorylation sites:

Amino acids 644-673;664-671

N-myristoylation sites:

Amino acids 10-16;15-21;79-85;99-105;118-124;188-194;
192-198;218-224;250-256;261-267;275-281;276-282;298-304;321-
327;328-334;420-426;421-427;440-446;449-455;599-605;626-632;
708-714;766-772;779-785

Amidation site:

Amino acids 693-697

Cell attachment sequences:

Amino acids 310-313;399-402

36/392

FIGURE 35

GGGGTCTCCCTCAGGGCCGGGAGGCACAGCGGTCCCTGCTTGCTGAAGGGCTGGATGTAC
GCATCCGCAGGTTCCCGCGGACTTGGGGGCGCCGCTGAGCCCCGGCGCCCGCAGAAGAC
TTGTGTTTGCCTCCTGCAGCCTCAACCCGGAGGGCAGCGAGGGCCTACCACCATGATCAC
TGGTGTGTTTCTGAGCATGCGCTTGTGGACCCAGTGGGCGTCCTGACCTCGCTGGCGTACTG
CCTGCACCAGCGGCGGGTGGCCCTGGCCGAGCTGCAGGAGGCGGATGGCCAGTGTCCGGT
CGACCGCAGCCTGCTGAAGTTGAAAATGGTGCAGGTCGTGTTTCGACACGGGGCTCGGAG
TCCTCTCAAGCCGCTCCCGCTGGAGGAGCAGGTAGAGTGGAACCCCCAGCTATTAGAGGT
CCCACCCCAAATCAGTTTGATTACACAGTCACCAATCTAGCTGGTGGTCCGAAACCATA
TTCTCCTTACGACTCTCAATACCATGAGACCACCCTGAAGGGGGGCATGTTTGCTGGGCA
GCTGACCAAGGTGGGCATGCAGCAAATGTTTGCCTTGGGAGAGAGACTGAGGAAGAACTA
TGTGGAAGACATTCCCTTTCTTTCACCAACCTTCAACCCACAGGAGGTCTTTATTTCGTTT
CACTAACATTTTTTCGGAATCTGGAGTCCACCCGTTGTTTCTGGCTGGGCTTTTCCAGTG
TCAGAAAGAAGGACCCATCATCATCCACACTGATGAAGCAGATTTCAGAAGTCTTGTATCC
CAACTACCAAAGCTGCTGGAGCCTGAGGCAGAGAACCAGAGGCCGAGGCAGACTGCCTC
TTTACAGCCAGGAATCTCAGAGGATTTGAAAAGGTGAAGGACAGGATGGGCATTGACAG
TAGTGATAAAGTGGAATCTTTCATCCTCCTGGACAACGTGGCTGCCGAGCAGGCACACAA
CCTCCCAAGCTGCCCCATGCTGAAGAGATTTGCACGGATGATCGAACAGAGAGCTGTGGA
CACATCCTTGTACATACTGCCCAAGGAAGACAGGGAAAGTCTTCAGATGGCAGTAGGCCC
ATTCTCCACATCCTAGAGAGCAACCTGCTGAAAGCCATGGACTCTGCCACTGCCCCCGA
CAAGATCAGAAAGCTGTATCTCTATGCGGCTCATGATGTGACCTTCATACCGCTCTTAAT
GACCCTGGGGATTTTTGACCACAAATGGCCACCGTTTGCTGTTGACCTGACCATGGAAT
TTACCAGCACCTGGAATCTAAGGAGTGGTTTGTGCAGCTCTATTACCACGGGAAGGAGCA
GGTGCCGAGAGGTGCCCCTGATGGGCTCTGCCCGCTGGACATGTTCTTGAATGCCATGTC
AGTTTATACCTTAAGCCCAGAAAAATACCATGCACTCTGCTCTCAAATCAGGTGATGGA
AGTTGGAAATGAAGAGTAACTGATTTATAAAAGCAGGATGTGTTGATTTTAAATAAAGT
GCCTTTATACAATG

37/392

FIGURE 36

MITGVFSMRLWTPVGVLTSLAYCLHQRRVALAELQEADGQCPVDRSLLKLKMVQVVFRHG
ARSPLKPLPLEEQVEWNPQLLEVPPQTQFDYTVTNLAGGPKPYSPYDSQYHETTLKGGMF
AGQLTKVGMQQMFALGERLRKNYVEDIPFLSPTFNPQEVFIRSTNIFRNLESTRCLLAGL
FQCQKEGPIIIHTDEADSEVLYPNYQSCWSLRQTRGRRTASLQPGISEDLKKVKDRMG
IDSSDKVDFILLDNVAEQAHNLPSCPMLKRFAFMIEQRAVDTSLYILPKEDRESLQMA
VGPFLHILESNNLLKAMDSATAPDKIRKLYLYAAHDVTFIPLLMTLGIFDHKWPPFAVDLT
MELYQHLESKEWFVQLYYHGKEQVPRGCPDGLCPLDMFLNAMS VYTLSP EKYHALCSQTQ
VMEVGNEE

Signal sequence:

amino acids 1-23

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 218-222

Casein kinase II phosphorylation site:

amino acids 87-91, 104-108, 320-324

Tyrosine kinase phosphorylation site:

amino acids 280-288

N-myristoylation site:

amino acids 15-21, 117-123, 118-124, 179-185, 240-246, 387-393

Amidation site:

amino acids 216-220

Leucine zipper pattern:

amino acids 10-32

Histidine acid phosphatases phosphohistidine signature:

amino acids 50-65

38/392

FIGURE 37

ACTGCACTCGGTTCTATCGATTGAATTCCCCGGGGATCCTCTAGAGATCCCTCGACCTCG
ACCCACGCGTCCGCGGACGCGTGGGCGGACGCGTGGGCCGGCTACCAGGAAGAGTCTGCC
GAAGGTGAAGGCCATGGACTTCATCACCTCCACAGCCATCCTGCCCCCTGCTGTTTCGGCTG
CCTGGGCGTCTTCGGCCTCTTCCGGCTGCTGCAGTGGGTGCGCGGGAAGGCCTACCTGCG
GAATGCTGTGGTGGTGATCACAGGCGCCACCTCAGGGCTGGGCAAAGAATGTGCAAAAGT
CTTCTATGCTGCGGGTGCTAAACTGGTGCTCTGTGGCCGGAATGGTGGGGCCCTAGAAGA
GCTCATCAGAGAACTTACCGCTTCTCATGCCACCAAGGTGCAGACACACAAGCCTTACTT
GGTGACCTTCGACCTCACAGACTCTGGGGCCATAGTTGCAGCAGCAGCTGAGATCCTGCA
GTGCTTTGGCTATGTGACATACTTGTCAACAATGCTGGGATCAGCTACCGTGGTACCAT
CATGGACACCACAGTGGATGTGGACAAGAGGGTCTATGGAGACAACTACTTTGGCCCAGT
TGCTCTAACGAAAGCACTCCTGCCCTCCATGATCAAGAGGAGGCAAGGCCACATTGTCGC
CATCAGCAGCATCCAGGGCAAGATGAGCATTCCTTTTCGATCAGCATATGCAGCCTCCAA
GCACGCAACCCAGGCTTTCTTTGACTGTCTGCGTGCCGAGATGGAACAGTATGAAATTGA
GGTGACCGTCATCAGCCCCGGCTACATCCACACCAACCTCTCTGTAAATGCCATCACCGC
GGATGGATCTAGGTATGGAGTTATGGACACCACCACAGCCCAGGGCCGAAGCCCTGTGGA
GGTGGCCCAGGATGTTCTTGCTGCTGTGGGGAAGAAGAAGAAAGATGTGATCCTGGCTGA
CTTACTGCCTTCCTTGGCTGTTTATCTTCGAACTCTGGCTCCTGGGCTCTTCTTCAGCCT
CATGGCCTCCAGGGCCAGAAAAGAGCGGAAATCCAAGAACTCCTAGTACTCTGACCAGCC
AGGGCCAGGGCAGAGAAGCAGCACTCTTAGGCTTGCTTACTCTACAAGGGACAGTTGCAT
TTGTTGAGACTTTAATGGAGATTTGTCTCACAAGTGGGAAAGACTGAAGAAACACATCTC
GTGCAGATCTGCTGGCAGAGGACAATCAAAAACGACAACAAGCTTCTTCCCAGGGTGAGG
GGAAACACTTAAGGAATAAATATGGAGCTGGGGTTTAACACTAAAACTAGAAATAAACA
TCTCAAACAGTAAAAAAAAAAAAAAAAAGGGCGGCCGCGACTCTAGAGTCGACCTGCAGAAG
CTTGGCCGCCATGGCCCAACTTGTTTATTGTCAGCTTATAATGGTTAC

39/392

FIGURE 38

MDFITSTAILPLLFGCLGVFGLFRLLQWVRGKAYLRNAVVVITGATSGLGKECAKVFYAA
GAKLVLCGRNGGALEELIRELTASHATKVQTHKPYLVTFDLTDSGAIVAAAAEILQCFGY
VDILVNNAGISYRGTIMDTTVDDVKRVMETNYFGPVALTKALLPSMIKRRQGHIVAIS
QGKMSIPFRSAYAASKHATQAFFDCLRAEMEQYEIEVTVISPGYIHTNLSVNAITADGSR
YGVMDTTTAQGRSPVEVAQDVLAAVGKKKKDVILADLLPSLAVYLRTLAPGLFFSLMASR
ARKERKSKNS

Signal sequence:

amino acids 1-21

Transmembrane domain:

amino acids 104-120, 278-292

N-glycosylation site:

amino acids 228-232

Glycosaminoglycan attachment site:

amino acids 47-51

Casein kinase II phosphorylation site:

amino acids 135-139, 139-143, 253-257

Tyrosine kinase phosphorylation site:

amino acids 145-153, 146-153

N-myristoylation site:

amino acids 44-50, 105-111, 238-244, 242-248, 291-297

Amidation site:

amino acids 265-269

Prokaryotic membrane lipoprotein lipid attachment site:

amino acids 6-17

40/392

FIGURE 39

GCAAGCCAAGGCGCTGTTTGAGAAGGTGAAGAAGTTCGGACCCATGTGGAGGAGGGGACATTGT
GTACCGCCTCTACATGCGGCAGACCATCATCAAGGTGATCAAGTTCATCCTCATCATCTGCTACAC
CGTCTACTACGTGCACAACATCAAGTTCGACGTGGACTGCACCGTGGACATTGAGAGCCTGACGGG
CTACCGCACCTACCGCTGTGCCACCCCTGGCCACACTCTTCAAGATCCTGGCGTCTTCTACAT
CAGCCTAGTCATCTTCTACGGCCTCATCTGCATGTACACACTGTGGTGGATGTCTACGGCGCTCCCT
CAAGAAGTACTCGTTTGAGTCGATCCGTGAGGAGAGCAGCTACAGCGACATCCCCGACGTCAAGAA
CGACTTCGCCTTCATGCTGCACCTCATTGACCAATACGACCCGCTCTACTCCAAGCGCTTCGCCGT
CTTCTGTGCGAGGTGAGTGAGAACAAAGCTGCGGCAGCTGAACCTCAACAACGAGTGGACGCTGGA
CAAGCTCCGGCAGCGGCTCACCAAGAACGCGCAGGACAAGCTGGAGCTGCACCTGTTTCATGCTCAG
TGGCATCCCTGACACTGTGTTTGACCTGGTGGAGCTGGAGGTCTCAAGCTGGAGCTGATCCCCGA
CGTGACCATCCCGCCAGCATTGCCAGCTCACGGGCCTCAAGGAGCTGTGGCTTACCACACAGC
GGCCAAGATTGAAGCGCCTGCGCTGGCCTTCTGCGCGAGAACCTGCGGGCGCTGCACATCAAGTT
CACCGACATCAAGGAGATCCCGCTGTGGATCTATAGCCTGAAGACACTGGAGGAGCTGCACCTGAC
GGGCAACCTGAGCGCGGAGAACCAACCGCTACATCGTCATCGACGGGCTGCGGGAGCTCAAACGCCT
CAAGGTGCTGCGGCTCAAGAGCAACCTAAGCAAGCTGCCACAGGTGGTCAAGATGTGGGCGTGCA
CCTGCAGAAGCTGTCCATCAACAATGAGGGCACCAAGCTCATCGTCTCAACAGCCTCAAGAAGAT
GGCGAACCTGACTGAGCTGGAGCTGATCCGCTGCGACCTGGAGCGCATCCCCCACTCCATCTTCAG
CTTCCACAACCTGCAGGAGATTGACCTCAAGGACAACAACCTCAAGACCATCGAGGAGATCATCAG
CTTCCAGCACCTGCACCGCCTCACCTGCCTTAAGCTGTGGTACAACCACATCGCCTACATCCCAT
CCAGATCGGCAACCTCACCAACCTGGAGCGCCTCTACCTGAACCGCAACAAGATCGAGAAGATCCC
CACCCAGCTCTTCTACTGCCGCAAGCTGCGCTACCTGGACCTCAGCCACAACAACCTGACCTTCTCT
CCCTGCCGACATCGGCCCTCTGCAGAACCTCCAGAACCTAGCCATCACGGCCAACCGGATCGAGAC
GTCCCTCCCGAGCTCTTCCAGTGCCGGAAGCTGCGGGCCCTGCACCTGGGCAACAACGTGTGCA
GTCACTGCCCTCCAGGGTGGGCGAGCTGACCAACCTGACGCGAGATCGAGCTGCGGGGCAACCGGCT
GGAGTGCCTGCCTGTGGAGCTGGGCGAGTGCCCACTGCTCAAGCGCAGCGGCTTGGTGGTGGAGGA
GGACCTGTTCAACACACTGCCACCCGAGGTGAAGGAGCGGCTGTGGAGGGCTGACAAGGAGCAGGC
CTGAGCGAGGCCGGCCAGCACAGCAAGCAGCAGGACCGCTGCCAGTCTCAGGCCCGGAGGGGC
AGGCCTAGCTTCTCCAGAACTCCCGGACAGCCAGGACAGCCTCGCGGCTGGGCGAGGAGCTGGGG
CCGCTTGTGAGTCAGGCCAGAGCGAGAGGACAGTATCTGTGGGGCTGGCCCTTTTCTCCCTCTGA
GACTCACGTCCCCCAGGGCAAGTGCTTGTGGAGGAGAGCAAGTCTCAAGAGCGCAGTATTTGGATA
ATCAGGGTCTCCTCCCTGGAGGCCAGCTCTGCCCCAGGGGCTGAGCTGCCACCAGAGGTCCTGGGA
CCCTCACTTTAGTTCTTGGTATTTATTTTCTCCATCTCCACCTCCTTCATCCAGATAACTTATA
CATTTCCCAAGAAAGTTTCAGCCAGATGGAAGGTGTTTCAGGGAAAGGTGGGCTGCCTTTTCCCTTG
TCCTTATTTAGCGATGCCGCCGGGCATTTAACACCCACCTGGACTTCAGCAGAGTGGTCCGGGGCG
AACCAGCCATGGGACGGTCACCCAGCAGTGCCGGGCTGGGCTCTGCGGTGCGGTCCACGGGAGAGC
AGGCCTCCAGCTGGAAAGGCCAGGCCTGGAGCTTGCCCTTTTCAGTTTTTGTGGCAGTTTTAGTTTT
TTGTTTTTTTTTTTTTTAATCAAAAAACAATTTTTTTTTAAAAAAGCTTTGAAATGGATGGTTT
GGGTATTAAGAAAGAAAAAAACTTAAAAAAGACACTAACGGCCAGTGAGTTGGAGTCTC
AGGGCAGGGTGGCAGTTTCCCTTGAGCAAAGCAGCCAGACGTTGAACGTGTGTTTCTTCCCTGGG
CGCAGGGTGCAGGGTGTCTTCCGATCTGGTGTGACCTTGGTCCAGGAGTTCTATTTGTTCTTGG
GAGGGAGGTTTTTTTGTGTTGTTTTTGGGTTTTTTTGGTGTCTTGTCTTCTCTCTCCATGTGT
CTTGGCAGGCACTATTCTGTGGCTGTGCGCCAGAGGAATGTTCTGGAGCTGCCAAGGAGGGAG
GAGACTCGGGTTGGCTAATCCCGGATGAACGGTGCTCCATTGCGACCTCCCTCCTCGTGCCTGC
CCTGCCTCTCCACGCACAGTGTTAAGGAGCCAAGAGGAGCCACTTCGCCCAGACTTTGTTTTCCCA
CCTCCTGCGGCATGGGTGTGTCCAGTGCCACCGCTGGCCTCCGCTGCTTCCATCAGCCCTGTGCC
ACCTGGTCTTTCATGAAGAGCAGACACTTAGAGGCTGGTGGGAATGGGGAGGTGCGCCCTGGGAG
GGCAGGCGTTGGTTCCAAGCCGGTTCCCGTCCCTGGCGCCTGGAGTGACACAGCCAGTCGGCAC
CTGGTGGCTGGAAGCCAACCTGCTTTAGATCACTCGGGTCCCCACCTTAGAAGGGTCCCCGCCTTA
GATCAATCACGTGGACACTAAGGCAGTTTTAGAGTCTCTTGTCTTAATGATTATGTCCATCCGTC
TGTCCGTCCATTTGTGTTTTCTGCGTCGTGTCATTGGATATAATCCTCAGAAATAATGCACACTAG
CCTCTGACAACCATGAAGCAAAAATCCGTTACATGTGGGTCTGAACTTGTTAGACTCGGTACAGTA
TCAAATAAATCTATAACAGAAAAA

41/392

FIGURE 40

MRQTIKVIKFILIIICYTVYYVHNIKFDVDCTVDIESLTGYRTYRCAHPLATLKFILASF
YISLVIFYGLICMYTLWWMLRRSLKKYSFESIREESSYSDIPDVKNDFAFMLHLIDQYDP
LYSKRFAVFLSEVSENKLRQLNLNNEWTLDKLRQLTKNAQDKLELHLFMLSIGIPDTVFD
LVELEVLKLELIPDVTIPPSIAQLTGLKELWLYHTAAKIEAPALAFLRENLRALHIKFTD
IKEIPLWIYSLKTLEELHLTGNLSAENNRIVIDGLRELKRLKVLRLKSNLSKLPQVVTD
VGVHLQKLSINNEGTKLIVLNSLKKMANLTELELIRCDLERIPHSIFSLHNLQEIDLKDN
NLKTIEEIIISFQHLHRLTCLKLWYNHIAIPIQIGNLTNLERLYLNRNKIEKIPTQLFYC
RKLRYLDLSHNNLTFLPADIGLLQNLQNLAITANRIETLPPELFCRKLRLALHLGNNVLQ
SLPSRVGELTNLTQIELRGNRLECLPVELGECPLLKRSGLVVEEDLFNTLPPEVKERLWR
ADKEQA

Transmembrane domain:

amino acids 51-75 (type II)

N-glycosylation site:amino acids 262-266, 290-294, 328-332, 396-400, 432-436,
491-495**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

amino acids 85-89

Casein kinase II phosphorylation site:amino acids 91-95, 97-101, 177-181, 253-257, 330-334, 364-368,
398-402, 493-497**N-myristoylation site:**

amino acids 173-179, 261-267, 395-401, 441-447

42/392

FIGURE 41

GGGGGAGAAGGCGGCCGAGCCCCAGCTCTCCGAGCACCGGGTCGGAAGCCGCGACCCGAG
CCGCGCAGGAAGCTGGGACCGGAACCTCGGCGGACCCGGCCCCACCCAACCTACCTGCGC
AGGTCACCAGCACCCCTCGGAACCCAGAGGCCCGCGCTCTGAAGGTGACCCCCCTGGGGAG
GAAGGCGATGGCCCCCTGCGAGGACGATGGCCCCGCGCCCGCCTCGCCCCGGCCGGCATCCC
TGCCGTGCGCTTGTGGCTTCTGTGCACGCTCGGCCTCCAGGGCACCCAGGCCGGGCCACC
GCCCCGCGCCCCCTGGGCTGCCCCGCGGGAGCCGACTGCCTGAACAGCTTTACCGCCGGGGT
GCCTGGCTTCGTGCTGGACACCAACGCCTCGGTGAGCAACGGAGCTACCTTCCTGGAGTC
CCCCACCGTGCGCCGGGGCTGGGACTGCGTGCGCGCCTGCTGCACCACCCAGAACTGCAA
CTTGGCGCTAGTGGAGCTGCAGCCCCGACCGCGGGGAGGACGCCATCGCCGCGCTGCTTCCT
CATCAACTGCCTCTACGAGCAGAACTTCGTGTGCAAGTTTCGCGCCCAGGGAGGGCTTCAT
CAACTACCTCACGAGGGAAGTGTACCGCTCCTACCGCCAGCTGCGGACCCAGGGCTTTGG
AGGGTCTGGGATCCCCAAGGCCTGGGCAGGCATAGACTTGAAGGTACAACCCAGGAACC
CCTGGTGCTGAAGGATGTGGAAAACACAGATTGGCGCCTACTGCGGGGTGACACGGATGT
CAGGGTAGAGAGGAAAGACCCAAACCAGGTGGAAGTGTGGGGACTCAAGGAAGGCACCTA
CCTGTTCCAGCTGACAGTGAAGTACTGCTCAGACCACCCAGAGGACACGGCCAACGTACAGT
CACTGTGCTGTCCACCAAGCAGACAGAAGACTACTGCCTCGCATCCAACAAGGTGGGTGCG
CTGCCGGGGCTCTTTCCACGCTGGTACTATGACCCACCGAGCAGATCTGCAAGAGTTT
CGTTTATGGAGGCTGCTTGGGCAACAAGAACAACCTACCTTCGGAAGAAGAGTGCATTCT
AGCCTGTGCGGGTGTGCAAGGTGGGCCTTTGAGAGGCAGCTCTGGGGCTCAGGCGACTTT
CCCCCAGGGCCCCCTCCATGGAAAGGCGCCATCCAGTGTGCTCTGGCACCTGTCAGCCAC
CCAGTTCCGCTGCAGCAATGGCTGCTGCATCGACAGTTTCCTGGAGTGTGACGACACCCC
CAACTGCCCCGACGCCTCCGACGAGGCTGCCTGTGAAAAATACACGAGTGGCTTTGACGA
GCTCCAGCGCATCCATTTCCCAGTGACAAAGGGCACTGCGTGGACCTGCCAGACACAGG
ACTCTGCAAGGAGAGCATCCCGCGCTGGTACTACAACCCCTTCAGCGAACACTGCGCCCCG
CTTTACCTATGGTGGTTGTTATGGCAACAAGAACAACCTTTGAGGAAGAGCAGCAGTGCCT
CGAGTCTTGTGCGGGCATCTCCAAGAAGGATGTGTTTGGCCTGAGGCGGGAAATCCCCAT
TCCCAGCACAGGCTCTGTGGAGATGGCTGTACAGTGTTCCTGGTCATCTGCATTGTGGT
GGTGGTAGCCATCTTGGGTTACTGCTTCTTCAAGAACCAGAGAAAGGACTTCCACGGACA
CCACCACCACCCACCCACCCCTGCCAGCTCCACTGTCTCCACTACCGAGGACACGGA
GCACCTGGTCTATAACCACACCACCCGGCCCCCTCTGAGCCTGGGTCTCACCGGCTCTCAC
CTGGCCCTGCTTCCTGCTTGCCAAGGCAGAGGCCTGGGCTGGGAAAAACTTTGGAACAG
ACTCTTGCTGTTTCCCAGGCCCACTGTGCCTCAGAGACCAGGGCTCCAGCCCCCTTTGG
AGAAGTCTCAGCTAAGCTCACGTCTGAGAAAGCTCAAAGGTTTGAAGGAGCAGAAAAC
CCTTGGGCCAGAAGTACCAGACTAGATGGACCTGCCTGCATAGGAGTTTGGAGGAAGTTG
GAGTTTTGTTTCTCTGTTCAAAGCTGCCTGTCCCTACCCCATGGTGCTAGGAAGAGGAG
TGGGGTGGTGTGAGACCCCTGGAGGCCCAACCCTGTCTCCTCCCGAGCTCCTCTTCCATGCT
GTGCGCCAGGGCTGGGAGGAAGGACTTCCCTGTGTAGTTTGTGCTGTAAAGAGTTGCTT
TTTGTTTATTTAATGCTGTGGCATGGGTGAAGAGGAGGGGAAGAGGCCTGTTTGGCCTCT
CTGTCTCTCTTCTTCCCCAAGATTGAGCTCTCTGCCCTTGATCAGCCCCACCCCTG
GCCTAGACCAGCAGACAGAGCCAGGAGAGGCTCAGCTGCATTCCGCAGCCCCCACCCTG
AGGTTCTCCAACATCACAGCCAGCCACCCACTGGGTAATAAAAAGTGGTGTGGAAAA
AAAAAAAAAAAAAAAAAAAAA

43/392

FIGURE 42

MAPARTMARARLAPAGIPAVALWLLCTLGLQGTQAGPPPAPPGLPAGADCLNSFTAGVPG
 FVLDTNASVSNGATFLESPTVRRGWDCVRACCTTQNCNLALVELQPDRGEDAIAACFLIN
 CLYEQNFVCKFAPREGFINYLTRVYRSYRQLRTQGFGGSGIPKAWAGIDLKVQPQEPLV
 LKDVENTDWRLLRGDTDVRRVERKDPNQVELWGLKEGTYLFQLTVTSSDHPEDTANVTVTV
 LSTKQTEDYCLASNKVGRCRGSFPRWYYDPTEQICKSFVYGGCLGNKNNYLREEECILAC
 RGVQGGPLRGSSGAQATFPQGSPMERHPVCSGTCQPTQFRCSNGCCIDSFLECDTPNC
 PDASDEAAKEYTSGFDELQRIHFPSDKGHCVDLPDTGLCKESI PRWYYNPFSEHCARFT
 YGGCYGNKNNFEEEEQQCLESCRGISKKDVFGRLRREIPI PSTGSEMAVTVFLVICIVVVV
 AILGYCFFKNQRKDFHGHHPPTPASSTVSTTEDTEHLVYNHTTRPL

signal sequence:

Amino acids 1-35

transmembrane domain:

Amino acids 466-483

N-glycosylation sites:

Amino acids 66-70;235-239;523-527

N-myristoylation sites:

A m i n o a c i d s
 29-35;43-49;161-167;212-218;281-287;282-288;285-291;
 310-316;313-319;422-428;423-429;426-432

Cell attachment sequence:

Amino acids 193-199

Pancreatic trypsin inhibitor (Kunitz) family signatures:

Amino acids 278-298;419-438

44/392

FIGURE 43

CCCACGCGTCCGCACCTCGGCCCCGGGCTCCGAAGCGGCTCGGGGGCGCCCTTTTCGGTCA
ACATCGTAGTCCACCCCCTCCCCATCCCCAGCCCCCGGGGATTTCAGGCTCGCCAGCGCCC
AGCCAGGGAGCCGGCCGGGAAGCGCGATGGGGGCCCCAGCCGCTCGCTCCTGCTCCTGC
TCCTGCTGTTTCGCCTGCTGCTGGGCGCCCCGGCGGGGCCAACCTCTCCAGGACGACAGCC
AGCCCTGGACATCTGATGAAACAGTGGTGGCTGGTGGCACCGTGGTGCTCAAGTGCCAAG
TGAAAGATCACGAGGACTCATCCCTGCAATGGTCTAACCTGCTCAGCAGACTCTCTACT
TTGGGGAGAAGAGAGCCCTTCGAGATAATCGAATTCAGCTGGTTACCTCTACGCCCCACG
AGCTCAGCATCAGCATCAGCAATGTGGCCCTGGCAGACGAGGGCGAGTACACCTGCTCAA
TCTTCACTATGCCTGTGCGAACTGCCAAGTCCCTCGTCACTGTGCTAGGAATTCCACAGA
AGCCCATCATCACTGGTTATAAATCTTCATTACGGGAAAAAGACACAGCCACCCTAAACT
GTCAGTCTTCTGGGAGCAAGCCTGCAGCCCGGCTCACCTGGAGAAAGGGTGACCAAGAAC
TCCACGGAGAACCAACCCGCATACAGGAAGATCCCAATGGTAAAACCTTCACTGTCAGCA
GCTCGGTGACATTCCAGGTTACCCGGGAGGATGATGGGGCGAGCATCGTGTGCTCTGTGA
ACCATGAATCTCTAAAGGGAGCTGACAGATCCACCTCTCAACGCATTGAAGTTTTATACA
CACCAACTGCGATGATTAGGCCAGACCCTCCCCATCCTCGTGAGGGCCAGAAGCTGTTGC
TACACTGTGAGGGTCGCGGCAATCCAGTCCCCCAGCAGTACCTATGGGAGAAGGAGGGCA
GTGTGCCACCCCTGAAGATGACCCAGGAGAGTGCCCTGATCTTCCCTTTCTCAACAAGA
GTGACAGTGGCACCTACGGCTGCACAGCCACCAGCAACATGGGCAGCTACAAGGCCCTACT
ACACCCTCAATGTTAATGACCCAGTCCGGTGCCCTCCTCCTCCAGCACCTACCACGCCA
TCATCGGTGGGATCGTGGCTTTCATTGTCTTCTGCTGCTCATCATGCTCATCTTCCTTG
GCCACTACTTGATCCGGCACAAAGGAACCTACCTGACACATGAGGCAAAAGGCTCCGACG
ATGCTCCAGACGCGGACACGGCCATCATCAATGCAGAAGGCGGGCAGTCAGGAGGGGACG
ACAAGAAGGAATATTTTCATCTAGAGGCGCCTGCCACTTCCTGCGCCCCCAGGGGCCCC
GTGGGGACTGCTGGGGCCGTCACCAACCCGGACTTGTACAGAGCAACCGCAGGGCCGCCC
CTCCCGCTTGCTCCCCAGCCACCCACCCCTGTACAGAATGTCTGCTTTGGGTGCGGT
TTTGTACTCGGTTTGAATGGGGAGGGAGGAGGGCGGGGGAGGGGAGGGTTGCCCTCAG
CCCTTTCGCTGGCTTCTCTGCATTGGGGTATTATTATTTTGTAAACAATCCCAATCAA
ATCTGTCTCCAGGCTGAGAGGCAGGAGCCCTGGGGTGAGAAAAGCAAAAACAAACAAA
AAACA

45/392

FIGURE 44

MGAPAASLLLLLLLLFACCWAPGGANLSQDDSQPWTSDETVVAGGTVVLKCQVKDHEDSSL
QWSNPAQQTLYFGEKRALRDNRIQLVTSTPHELSSISNVALADEGEYTCSTFTMPVRTA
KSLVTVLGIPQKPIITGYKSSLREKDTATLNCQSSGSKPAARLTWRKGDQELHGEPTRIQ
EDPNGKTFTVSSSVTFQVTRREDDGASIVCSVNHESLKGADRSTSQRIEVLYTPTAMIRPD
PPHPREGQKLLHCEGRGNPVPQQYLWEKEGSVPPLKMTQESALIFPFLNKSDSGTYGCT
ATSNMGSYKAYYTLNVNDPSPVPSSSSTYHAIIGGIVAFIVFLLLIIMLIFLGHYLIIRHKG
TYLTHEAKGSDDAPDADTAIINAEGGQSGGDDKKEYFI

Signal sequence:

amino acids 1-20

Transmembrane domain:

amino acids 331-352

N-glycosylation site:

amino acids 25-29, 290-294

Casein kinase II phosphorylation site:

amino acids 27-31, 35-39, 89-93, 141-145, 199-203, 388-392

N-myristoylation site:amino acids 2-8, 23-29, 156-162, 218-224, 295-301, 298-304,
306-310, 334-340, 360-364, 385-389, 386-390**Prokaryotic membrane lipoprotein lipid attachment site:**

amino acids 7-18

46/392

FIGURE 45

ACTTGCCATCACCTGTTGCCAGTGTGGAAAAATTCTCCCTGTTGAATTTTTTGCACATGG
AGGACAGCAGCAAAGAGGGCAACACAGGCTGATAAGACCAGAGACAGCAGGGAGATTATT
TTACCATACGCCCTCAGGACGTTCCCTCTAGCTGGAGTTCTGGACTTCAACAGAACCCCA
TCCAGTCATTTTGATTTTGCTGTTTTATTTTTTTTTTTCTTTTTCTTTTTCCCACCACATTG
TATTTTATTTCCGTACTTCAGAAATGGGCCTACAGACCACAAAGTGGCCAGCCATGGGG
CTTTTTTCTGGAAGTCTTGGCTTATCATTTCCCTGGGGCTCTACTCACAGGTGTCCAAAC
TCCTGGCCTGCCCTAGTGTGTGCCGCTGCGACAGGAACCTTGTCTACTGTAATGAGCGAA
GCTTGACCTCAGTGCCTCTTGGGATCCCGGAGGGCGTAACCGTACTCTACCTCCACAACA
ACCAAATTAATAATGCTGGATTTCCTGCAGAACTGCACAATGTACAGTCGGTGCACACGG
TCTACCTGTATGGCAACCAACTGGACGAATCCCCATGAACCTTCCCAAGAATGTCAGAG
TTCTCCATTTGCAGGAAAACAATATTAGACCATTTACGGGCTGCTCTTGCCCAGCTCT
TGAAGCTTGAAGAGCTGCACCTGGATGACAACTCCATATCCACAGTGGGGGTGGAAGACG
GGGCTTCCGGGAGGCTATTAGCCTCAAATTGTTGTTTTGTCTAAGAATCACCTGAGCA
GTGTGCCTGTTGGGCTTCTGTGGACTTGCAAGAGCTGAGAGTGGATGAAAATCGAATTG
CTGTCATATCCGACATGGCCTTCCAGAATCTCACGAGCTTGGAGCGTCTTATTGTGGACG
GGAACCTCCTGACCAACAAGGGTATCGCCGAGGGCACCTTCAGCCATCTCACCAAGCTCA
AGGAATTTTCAATTGTACGTAATTCGCTGTCCACCCTCCTCCCGATCTCCCAGGTACGC
ATCTGATCAGGCTCTATTTGCAGGACAACCAGATAAACACATTCCTTTGACAGCCTTCT
CAAATCTGCGTAAGCTGGAACGGCTGGATATATCCAACAACCAACTGCGGATGCTGACTC
AAGGGGTTTTTGATAATCTCTCCAACCTGAAGCAGCTCACTGCTCGGAATAACCCTTGGT
TTTGTGACTGCAGTATTAAATGGGTACAGAATGGCTCAAATATATCCCTTCATCTCTCA
ACGTGCGGGGTTTCATGTGCCAAGGTCTGAACAAGTCCGGGGGATGGCCGTGAGGGAAT
TAAATATGAATCTTTTGTCTGTCCACCACGACCCCCGGCCTGCCTCTCTTACCCCCAG
CCCCAAGTACAGCTTCTCCGACCACTCAGCCTCCCACCCTCTCTATTCCAAACCCTAGCA
GAAGCTACACGCCTCCAACCTCCTACCACATCGAACTTCCCACGATTCTCTGACTGGGATG
GCAGAGAAAGAGTGACCCACCTATTTCTGAACGGATCCAGCTCTCTATCCATTTTGTGA
ATGATACTTCCATTCAAGTCAGCTGGCTCTCTCTCTTACCCTGATGGCATACAAACCTCA
CATGGGTGAAAATGGGCCACAGTTTAGTAGGGGGCATCGTTTCAGGAGCGCATAGTCAGCG
GTGAGAAGCAACACCTGAGCCTGGTTAACTTAGAGCCCCGATCCACCTATCGGATTTGTT
TAGTGCCACTGGATGCTTTTAACTACCGCGCGGTAGAAGACACCATTTGTTTCAGAGGCCA
CCACCCATGCCTCCTATCTGAACAACGGCAGCAACACAGCGTCCAGCCATGAGCAGACGA
CGTCCCACAGCATGGGCTCCCCCTTTCTGCTGGCGGGCTTGATCGGGGGCGCGGTGATAT
TTGTGCTGGTGGTCTTGCTCAGCGTCTTTTGTGCTGGCATATGCACAAAAGGGGCGCTACA
CCTCCAGAAAGTGGAATAACAACGGGGCGGGCGGAAAGATGATTATTGCGAGGCAGGCA
CCAAGAAGGACAACTCCATCCTGGAGATGACAGAAACCAGTTTTTCAGATCGTCTCTTAA
ATAACGATCAACTCCTTAAAGGAGATTTAGACTGCAGCCCATTTACACCCCAAATGGGG
GCATTAATTACACAGACTGCCATATCCCCAACACATGCGATACTGCAACAGCAGCGTGC
CAGACCTGGAGCACTGCCATACGTGACAGCCAGAGGCCAGCGTTATCAAGGCGGACAAT
TAGACTCTTGAGAACACACTCGTGTGTGCACATAAAGACACGCAGATTACATTTGATAAA
TGTTACACAGATGCATTTGTGCATTTGAATACTCTGTAATTTATACGGTGTACTATATAA
TGGGATTTAAAAAAGTGCTATCTTTTCTATTTCAAGTTAATTACAAACAGTTTTGTAA
TCTTTGCTTTTAAATCTT

47/392

FIGURE 46

MGLQTTKWPSHGAFFLKSWLIISLGLYSQVSKLLACPSVCRCDRNFVYCNERSLTSVPLG
IPEGVTVLVYLHNNQINNAGFPAELHNVQSVHTVYLYGNQLDEFPMNLPKNVRVLHLQENN
IQTISRALAQLLKLEELHLDDNSISTVGVEDGAFREAI SLKLLFLSKNHLSSVPVGLPV
DLQELRV DENRIAVISDMAFQNLTSLERLIVDGNLLTNKGIAEGTFSHLTKLKEFSIVRN
SLSHPPPDLPGTHLIRLYLQDNQINHIPLTAFSNLRKLERLDISNNQLRMLTQGVFDNLS
NLKQLTARNNPWFCDCSI KWVTEWLKYIPSSLNVRGFMCGPEQVRGMVAVRELNMNLLSC
PTTTPGLPLFTPAPSTASPTTQPPTLSIPNPSRSYTPPTPTTSKLPTIPDWDGRERVTPP
ISERIQLSIHFVNDTSIQVSWLSLFTVMAYKLTWVKMGHSLVGGIVQERIVSGEKQHLSL
VNLEPRSTYRICLVPLDAFN YRAVEDTICSEATTHAS YLNNGSNTASSHEQTTSHSMGSP
FLLAGLIGGAVIFVLVVLLSVFCWHMHKKGRTYSQWKYNRGRRKDDYCEAGTKKDNSIL
EMTETSFQIVSLNNDQLLKGD FRLQPIYTPNGGIN YTDCHI PNNMRYCNSSVPDLEHCHT

Signal peptide:

amino acids 1-42

Transmembrane domain:

amino acids 542-561

N-glycosylation site:amino acids 202-206, 298-302, 433-437, 521-525, 635-639,
649-653**Casein kinase II phosphorylation site:**amino acids 204-208, 407-411, 527-531, 593-597, 598-602,
651-655**Tyrosine kinase phosphorylation site:**

amino acids 319-328

N-myristoylation site:amino acids 2-8, 60-66, 149-155, 213-219, 220-226, 294-300,
522-528, 545-551, 633-639**Amidation site:**

amino acids 581-585

Leucine zipper pattern:

amino acids 164-186

Phospholipase A2 aspartic acid active site:

amino acids 39-50

48/392

FIGURE 47

GCAGCGAGCGCCGGGTGCGGCCCTGCCGCCGCAGGGATGTGACCTTCACCGTCGCTTAGC
CAGGATGACCGGAGCCCGTGTCTCGCGCGTCCGCGCCTCGCTTCAGCCTCCCGGGTGCT
CTGACCGCACGCTCCCGGCTGCTAGGCTCCCCGGCACCGGCCTCGCCATGCCGCCACCGC
CCGGGCCCGCCGCCGCCCTGGGCACTGCGCTTCTGCTGCTCCTGCTGGCTTCCGAGTCTT
CTCACACTGTGCTGTTGCGGGCGCGTGAGGCGGCGCAGTTTCTGCGGCCAGGCAGCGCC
GCGCCTACCAAGTCTTCGAGGAGGCCAAGCAGGGCCACCTGGAACGGGAGTGCGTGGAGG
AGGTGTGCAGCAAAGAGGAGGCCAGAGAGGTGTTGAGAACGACCCCGAGACGGAGTATT
TCTATCCACGATATCAAGAGTGCATGAGAAAATATGGCAGGCCTGAAGAAAAAACCAG
ATTTCCGCAAATGTGTTCAGAACCTGCCTGACCAGTGCACCCCAAACCCTTGTGATAAGA
AGGGTACTCATATCTGCCAAGACCTCATGGGCAACTTCTTCTGCGTGTGCACAGATGGCT
GGGGAGGCCGGCTCTGTGACAAAGATGTCAATGAGTGTGTCCAGAAGAATGGGGGCTGCA
GCCAGGTCTGCCACAACAAACCAGGAAGCTTCCAATGTGCCTGCCATAGTGGCTTCTCGC
TTGCATCAGACGGCCAGACCTGCCAAGATATCGATGAATGCACAGACTCAGACACCTGTG
GGGACGCGCGATGCAAGAACTTGCCAGGCTCCTACTCTTGCCCTCTGCGATGAGGGATATA
CATACAGCTCCAAGGAGAAGACCTGCCAAGATGTGGACGAGTGCCAGCAGGATCGCTGTG
AGCAGACCTGTGTCAACTCCCAGGCAGCTATACCTGCCACTGTGATGGGCGAGGGGGCC
TAAACTATCCCCAGACATGGATACTTGTGAGGACATCTTACCATGTGTGCCCTTCAGCA
TGGCCAAGAGCGTGAAGTCCTTGTACCTGGGCCGCATGTTTCAGCGGGACCCCCGTGATTA
GACTACGCTTCAAGAGGCTTCAGCCTACCAGGCTGCTGGCTGAATTTGACTTCCGCACTT
TTGACCCTGAAGGAGTCCTCTTCTTCGCTGGAGGCCGTTTCAGACAGCACCTGGATTGTCC
TGGGCCTAAGAGCTGGGCGGCTTGAGCTGCAGCTTCGGTACAATGGCGTTGGGCGCATCA
CCAGCAGCGGGCCAACCATCAACCACGGCATGTGGCAAACCTATCTCCGTGGAAGAGCTGG
AACGTAACCTTGTCAATCAAGGTCAACAAAGATGCTGTAATGAAGATCGCGGTAGCTGGGG
AGCTGTTTCAGCTGGAGAGGGGCTCTATCACCTGAATCTCACCGTGGGCGGCATTCCCT
TCAAGGAGAGTGAGCTCGTCCAGCCGATTAACCCTCGCCTGGATGGGTGCATGAGGAGTT
GGAACCTGGCTGAACGGGGAAGACAGCGCCATCCAGGAGACAGTCAAGGCAAACACAAAA
TGCAGTGCTTCTCTGTGACAGAAAGGGGCTCCTTCTTCCCGGGGAATGGATTGTCTACCT
ACAGGCTCAACTACACCCGAACATCGCTGGATGTGGCACGGAAACCACCTGGGAAGTTA
AAGTTGTGGCTCGGATCCGCCCTGCCACGGACACGGGGGTGCTGCTGGCGCTGGTGGGGG
ACGACGATGTCGTCTCTGTGGCCCTAGTCGACTACCACTCTACAAAGAAGCTCAAGA
AGCAGTTGGTGGTCTTGGCAGTTGAGGATGTTGCCCTGGCACTGATGGAAATCAAGGTGT
GCGACAGCCAGGAACACACGGTCACTGTCTCCCTGCGGGAGGGTGAGGCCACCCTAGAAG
TGGATGGCACAAAGGGCCAGAGTGAAGTGAGCACTGCCAGCTGCAGGAGCGACTGGACA
CACTTAAGACACATCTGCAAGGCTCTGTGCACACCTATGTTGGAGGCCTGCCAGAAGTAT
CGGTGATTTCTGCACCCGTCCTGCGTTCTACCGCGGATGCATGACTCTGGAGGTAAACG
GGAAAATCCTGGACCTGGATACGGCCTCGTACAAGCACAGTGACATCACCTCCCACTCCT
GCCCCGCTGTGGAGCATGCCACCCCCTAGACCGAGCTGCAAGAGGGCTCCACACCTAAAG
ACAAAAATGAAGCAGGGTTTGGACACACAGCACTGGCTCCTCTCGCATGGTCTCTGCAACA
CTGGAGCAGCGTGGACCGCCCTTGTGGTTTTTTTTTTCTTGAGATCTTCTTTTTTGCCTTG
TAACATATCTGTACATAATGGACGGGTGTGGGTACACGGCTGCTCAGAGAGAGCCACGT
GACCTGGTGGGAGCTGGCTGGAAGGGGCTGGGCTAGAGGGGCTGGCAGTTTGCAGCAGAA
CGGATGTGAAGAAAATAATTCTCTATTATTTTATTACTACATGCTTCTTTCTGACTCTA
AAATATGGAATAAAATATTTACAGAAACCTTTTAAAAA

49/392

FIGURE 48

MPPPPGPAAALGTALLLLLLLASESSHTVLLRAREAAQFLRPRQRRAYQVFEEAKQGHLE
ECVVEVCSKEEAREVFENDPETEYFYPRYQECMRKYGRPEEKNPDFAKCVQNLDPDQCTPN
PCDKKGTHICQDLMGNFFCVCTDGGWGGRLCDKDVNECVQKNGGCSQVCHNKPGSFQCACH
SGFSLASDGQTCQDIDECTDSDTCDGARDCKNLPGSYSLCDEGYTYSSKEKTCQDVDECQ
QDRCEQTCVNSPGSYTCHCDGRGGLKLSPEMDTCEDILPCVPFMAKSVKSLYLGRMFSG
TPVIRLRFKRLQPTRLLAEFDFTFDPEGVLFFAGGRSDSTWIVLGLRAGRLELQLRYNG
VGRITSSGPTINHGMWQTISVEELERNLVIKVNKDAVMKIAVAGELFQLERGLYHLNLT
GGIPFKESELVQPINPRLDGCMRSWNWLNGEDSAIQETVKANTKMQCFSVTERGSFFPGN
GFATYRLNYTRTSLDVGTETTWEVKVVARIRPATDTGVLLALVGDDDVVISVALVDYHST
KKLKKQLVVLAVEDVALALMEIKVCDSEHTVTVSLREGEATLEVDGTKGQSEVSTAQLQ
ERLDTLKTHLQGSVHTYVGGLPEVSVISAPVTAFYRGCMTELVNGKILDLDTASYKHSDI
TSHSCPPVEHATP

50/392

FIGURE 49

CGCCGCGCTCCCGCACCCGCGGCCCGCCACCGCGCCGCTCCCGCATCTGCACCCGCAGC
CCGGCGGCCTCCCGGCGGGAGCGAGCAGATCCAGTCCGGCCCGCAGCGCAACTCGGTCCA
GTCGGGGCGGCGGCTGCGGGCGCAGAGCGGAGATGCAGCGGCTTGGGGCCACCCTGCTGT
GCCTGCTGCTGGCGGGCGGCGGTCCCCACGGCCCCCGCGCCCGCTCCGACGGCGACCTCGG
CTCCAGTCAAGCCCGGCCCGGCTCTCAGCTACCCGCAGGAGGAGGCCACCCTCAATGAGA
TGTTCCGCGAGGTTGAGGAACTGATGGAGGACACGCAGCACAAATTGCGCAGCGCGGTGG
AAGAGATGGAGGCAGAAGAAGCTGCTGCTAAAGCATCATCAGAAGTGAACCTGGCAAACCT
TACCTCCAGCTATCACAATGAGACCAACACAGACACGAAGGTTGGAAATAATACCATCC
ATGTGCACCGAGAAATTCACAAGATAACCAACAACCAGACTGGACAAATGGTCTTTTCAG
AGACAGTTATCACATCTGTGGGAGACGAAGAAGGCAGAAGGAGCCACGAGTGCATCATCG
ACGAGGACTGTGGGCCCAGCATGTACTGCCAGTTTGCCAGCTTCCAGTACACCTGCCAGC
CATGCCGGGGCCAGAGGATGCTCTGCACCCGGGACAGTGAGTGCTGTGGAGACCAGCTGT
GTGTCTGGGGTCACTGCACCAAAATGGCCACCAGGGGCAGCAATGGGACCATCTGTGACA
ACCAGAGGGACTGCCAGCCGGGGCTGTGCTGTGCCTTCCAGAGAGGCCCTGCTGTTCCCTG
TGTGCACACCCCTGCCCGTGGAGGGCGAGCTTTGCCATGACCCCGCCAGCCGGCTTCTGG
ACCTCATCACCTGGGAGCTAGAGCCTGATGGAGCCTTGGACCGATGCCCTTGTGCCAGTG
GCCTCCTCTGCCAGCCCCACAGCCACAGCCTGGTGTATGTGTGCAAGCCGACCTTCGTGG
GGAGCCGTGACCAAGATGGGGAGATCCTGCTGCCCAGAGAGGTCCCCGATGAGTATGAAG
TTGGCAGCTTCATGGAGGAGGTGCGCCAGGAGCTGGAGGACCTGGAGAGGAGCCTGACTG
AAGAGATGGCGCTGGGGGAGCCTGCGGCTGCCGCCGCTGCACTGCTGGGAGGGGAAGAGA
TTTAGATCTGGACCAGGCTGTGGGTAGATGTGCAATAGAAATAGCTAATTTATTTCCCCA
GGTGTGTGCTTTAGGCGTGGGCTGACCAGGCTTCTTCTTACATCTTCTTCCAGTAAGTT
TCCCTCTGGCTTGACAGCATGAGGTGTTGTGCAATTTGTTTTCAGTCCCCCAGGCTGTTCT
CCAGGCTTCACAGTCTGGTGCTTGGGAGAGTCAGGCAGGGTTAAACTGCAGGAGCAGTTT
GCCACCCCTGTCCAGATTATTGGCTGCTTTGCCTCTACCAGTTGGCAGACAGCCGTTTGT
TCTACATGGCTTTGATAATTGTTTGAGGGGAGGAGATGGAAACAATGTGGAGTCTCCCTC
TGATTGGTTTTTGGGAAATGTGGAGAAGAGTGCCCTGCTTTGCAAACATCAACCTGGCAA
AAATGCAACAAATGAATTTTCCACGCAGTTCTTTCCATGGGCATAGGTAAGCTGTGCCTT
CAGCTGTTGCAGATGAAATGTTCTGTTCAACCTGCATTACATGTGTTTATTTCATCCAGCA
GTGTTGCTCAGCTCCTACCTCTGTGCCAGGGCAGCATTTTTCATATCCAAGATCAATTCCC
TCTCTCAGCACAGCCTGGGGAGGGGGTCATTGTTCTCCTCGTCCATCAGGGATCTCAGAG
GCTCAGAGACTGCAAGCTGCTTGCCCAAGTCACACAGCTAGTGAAGACCAGAGCAGTTTC
ATCTGGTTGTGACTCTAAGCTCAGTGCTCTCTCCACTACCCACACCAGCCTTGGTGCCA
CCAAAAGTGCTCCCCAAAAGGAAGGAGAATGGGATTTTTCTTGAGGCATGCACATCTGGA
ATTAAGGTCAAACCTAATTCTCACATCCCTCTAAAAGTAACTACTGTTAGGAACAGCAGT
GTTCTCACAGTGTGGGGCAGCCGTCTTCTAATGAAGACAATGATATTGACACTGTCCCT
CTTTGGCAGTTGCATTAGTAACTTTGAAAGGTATATGACTGAGCGTAGCATAACAGGTTAA
CCTGCAGAAACAGTACTTAGGTAATTGTAGGGCGAGGATTATAAATGAAATTTGCAAAAT
CACTTAGCAGCAACTGAAGACAATTATCAACCACGTGGAGAAAATCAAACCGAGCAGGGC
TGTGTGAAACATGGTTGTAATATGCGACTGCGAACACTGAACTCTACGCCACTCCACAAA
TGATGTTTTTCAGGTGTGATGGACTGTTGCCACCATGTATTTCATCCAGAGTTCTTAAAGTT
TAAAGTTGCACATGATTGTATAAGCATGCTTTCTTTGAGTTTTAAATTATGTATAAACAT
AAGTTGCATTTAGAAATCAAGCATAAATCACTTCAACTGCAAAAAAAAAAAAAAAAAAAAA
AAAAAA

51/392

FIGURE 50

MQRLGATLLCLLLAAVPTAPAPAPTATSAPVKPGPALSYPQEEATLNEMFREVEELMED
TQHKLRSAVEEMEAEEAAAKASSEVNLANLPSPYHNETNTDTKVGNNNTIHVHREIHKITN
NQTGQMFSETVITSVGDEEGRRSHECIIDEDCGPSMYCQFASFQYTCQPCRGQRMCTR
DSECCGDQLCVWGHCTKMATRGSNGTICDNQRDCQPGLCCAFQRGLLFPVCTPLPVEGEL
CHDPASRLLDLITWELEPDGALDRCPASGLLCQPHSHSLVYVCKPTFVGSRDQDGEILL
PREVPDEYEVGSFMEEVRQELEDLERSLTEEMALGEPAAAAAALLGGEET

Signal sequence:

amino acids 1-19

N-glycosylation site:

amino acids 96-100, 106-110, 121-125, 204-208

Casein kinase II phosphorylation site:amino acids 46-50, 67-71, 98-102, 135-139, 206-210, 312-316,
327-331**N-myristoylation site:**

amino acids 202-208, 217-223

Amidation site:

amino acids 140-144

52/392

FIGURE 51

GCCTGTTGCTGATGCTGCCGTGCGGTACTTGTCATGGGAGCTGGCACTGCGGCGCTCTCCC
GTCCCGCGGTGGTTGCTGCTGCTGCCGTGCTGCTGGGCCTGAACGCAGGAGCTGTCATT
GACTGGCCCCACAGAGGAGGGCAAGGAAGTATGGGATTATGTGACGGTCCGCAAGGATGCC
TACATGTTCTGGTGGCTCTATTATGCCACCAACTCCTGCAAGAACTTCTCAGAACTGCCC
CTGGTCATGTGGCTTCAGGGCGGTCCAGGCGGTTCTAGCACTGGATTGGAACCTTTGAG
GAAATTGGGCCCCCTTGACAGTGATCTCAAACCACGGAAAACCACTGGCTCCAGGCTGCC
AGTCTCCTATTTGTGGATAATCCCGTGGGCACTGGGTTCAAGTTATGTGAATGGTAGTGGT
GCCTATGCCAAGGACCTGGCTATGGTGGCTTCAGACATGATGGTTCTCCTGAAGACCTTC
TTCAGTTGCCACAAAGAATTCAGACAGTTCCATTCTACATTTTCTCAGAGTCCTATGGA
GGAAAAATGGCAGCTGGCATTGGTCTAGAGCTTTATAAGGCCATTACAGCGAGGGACCATC
AAGTGCAACTTTGCGGGGGTTGCCCTGGGTGATTCTCTGGATCTCCCCTGTTGATTGCGGTG
CTCTCCTGGGGACCTTACCTGTACAGCATGTCTCTTCTCGAAGACAAAGGTCTGGCAGAG
GTGTCTAAGGTTGCAGAGCAAGTACTGAATGCCGTAAATAAGGGGCTCTACAGAGAGGCC
ACAGAGCTGTGGGGGAAAGCAGAAATGATCATTGAACAGAACACAGATGGGGTGAACCTTC
TATAACATCTTAATAAAAGCACTCCACGTCTACAATGGAGTCGAGTCTAGAATTCACA
CAGAGCCACCTAGTTTGTCTTTGTGTCAGCGCCACGTGAGACACCTACAACGAGATGCCTTA
AGCCAGCTCATGAATGGCCCCATCAGAAAGAAGCTCAAAATTATTCCTGAGGATCAATCC
TGGGGAGGCCAGGCTACCAACGTCTTTGTGAACATGGAGGAGGACTTCATGAAGCCAGTC
ATTAGCATTGTGGACGAGTTGCTGGAGGCAGGGATCAACGTGACGGTGTATAATGGACAG
CTGGATCTCATCGTAGATAACCATGGGTGAGGAGGCTGGGTGCGGAACTGAAGTGGCCA
GAACTGCCTAAATTCAGTCAGCTGAAGTGAAGGCCCTGTACAGTGACCCTAAATCTTTG
GAAACATCTGCTTTTGTCAAGTCTTACAAGAACCTTGCTTTCTACTGGATTCTGAAAGCT
GGTCATATGGTTCCTTCTGACCAAGGGGACATGGCTCTGAAGATGATGAGACTGGTGA
CAGCAAGAATAGGATGGATGGGGCTGGAGATGAGCTGGTTTGGCCTTGGGGCACAGAGCT
GAGCTGAGGCGCGCTGAAGCTGTAGGAAGCGCCATTCTTCCCTGTATCTAACTGGGGCTGT
GATCAAGAAGGTTCTGACCAGCTTCTGCAGAGGATAAAATCATTGTCTCTGGAGGCAATT
TGGAATTTATTTCTGCTTCTTAAAAAACCTAAGATTTTTTAAAAAATTGATTTGTTTTG
ATCAAAATAAAGGATGATAATAGATATTAA

53/392

FIGURE 52

MELALRRSPVPRWLLLLPLLLGLNAGAVIDWPTEEGKEVWDYVTVRKDAYMFWWLYYATN
SCKNFSELPLVMWLQGGPGGSSTGFGNFEEIGPLDSLKPRKTTWLQAASLLFVDNPVGT
GFSYVNGSGAYAKDLAMVASDMMVLLKTFFSCHKEFQTVPFYIFSESYGGKMAAGIGLEL
YKAIQRGTIKCNFAGVALGDSWISPVDSVLSWGPYLYSMSLLEDKGLAEVSKVAEQVLNA
VNKGLYREATELWGKAEMIEQNTDGVNFYNILTKSTPTSTMESSLEFTQSHLVCLCQRH
VRHLQORDALSQLMNGPIRKCLKIIPEDQSWGGQATNVFVNMEEDFMKPVISIVDELLEAG
INVTVYNGQLDLIVDTMGQEAWVRKLKWPELPKFSQLKWKALYSDPKSLETSAFVKSYKN
LAFYWILKAGHMVPSDQGDMAKMMRLVTQQE

Signal sequence:

amino acids 1-25

N-glycosylation site:

amino acids 64-68, 126-130, 362-366

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 101-105

Casein kinase II phosphorylation site:amino acids 204-208, 220-224, 280-284, 284-288, 351-355,
449-453**N-myristoylation site:**amino acids 22-28, 76-82, 79-85, 80-86, 119-125, 169-175,
187-193, 195-201, 331-337, 332-338, 360-366

54/392

FIGURE 53

GTCTGTTCCCAGGAGTCCTTCGGCGGCTGTTGTGTCA GTGGCCTGATCGCGATGGGGACA
AAGGCGCAAGTCGAGAGGAACTGTTGTGCCTCTTCATATTGGCGATCCTGTTGTGCTCC
CTGGCATTGGGCAGTGTTACAGTGCCTCTTCTGAACCTGAAGTCAGAATTCCTGAGAAT
AATCCTGTGAAGTTGTCTGTGCCTACTCGGGCTTTTCTTCTCCCCGTGTGGAGTGGAAG
TTTGACCAAGGAGACACCACCAGACTCGTTTGCTATAATAACAAGATCACAGCTTCCTAT
GAGGACCGGGTGACCTTCTTGCCAACTGGTATCACCTTCAAGTCCGTGACACGGGAAGAC
ACTGGGACATACACTTGTATGGTCTCTGAGGAAGGCGGCAACAGCTATGGGGAGGTCAAG
GTCAAGCTCATCGTGCTTGTGCCTCCATCCAAGCCTACAGTTAACATCCCCCTCCTCTGCC
ACCATTTGGGAACCGGGCAGTGCTGACATGCTCAGAACAAGATGGTTCCCCACCTTCTGAA
TACACCTGGTTCAAAGATGGGATAGTGATGCCTACGAATCCCAAAGCACCCGTGCCTTC
AGCAACTCTTCTATGTCTGAATCCCACAACAGGAGAGCTGGTCTTTGATCCCCCTGTCA
GCCTCTGATACTGGAGAATACAGCTGTGAGGCACGGAATGGGTATGGGACACCCATGACT
TCAAATGCTGTGCGCATGGAAGCTGTGGAGCGGAATGTGGGGGTCTCGTGGCAGCCGTC
CTTGTAACCCTGATTCTCCTGGGAATCTTGTTTTTGGCATCTGGTTTGCCTATAGCCGA
GGCCACTTTGACAGAACAAAGAAAGGGACTTCGAGTAAGAAGGTGATTTACAGCCAGCCT
AGTGCCCGAAGTGAAGGAGAATTCAAACAGACCTCGTCATTCTGGTGTGAGCCTGGTCG
GCTCACCGCCTATCATCTGCATTTGCCTTACTCAGGTGCTACCGGACTCTGGCCCCTGAT
GTCTGTAGTTTACAGGATGCCTTATTTGTCTTCTACACCCACAGGGCCCCCTACTTCT
TCGGATGTGTTTTTAATAATGTCAGCTATGTGCCCCATCCTCCTTCATGCCCTCCCTCCC
TTTCTTACCACTGCTGAGTGGCCTGGAACCTGTTTAAAGTGTTTATTCCCCATTTCTTTG
AGGGATCAGGAAGGAATCCTGGGTATGCCATTGACTTCCCTTCTAAGTAGACAGCAAAAA
TGGCGGGGGTTCGAGGAATCTGCACTCAACTGCCACCTGGCTGGCAGGGATCTTTGAAT
AGGTATCTTGAGCTTGGTTCTGGGCTCTTTCCTTGTGTACTGACGACCAGGGCCAGCTGT
TCTAGAGCGGGAATTAGAGGCTAGAGCGGCTGAAATGGTTGTTTGGTGATGACACTGGGG
TCCTTCCATCTCTGGGGCCCACTCTCTTCTGTCTTCCCATGGGAAGTGCCACTGGGATCC
CTCTGCCCTGTCTCCTGAATACAAGCTGACTGACATTGACTGTGTCTGTGGAAAATGGG
AGCTCTTGTTGTGGAGAGCATAGTAAATTTTCAGAGAACTTGAAGCCAAAAGGATTTAAA
ACCGCTGCTCTAAAGAAAAGAAAACCTGGAGGCTGGGCGCAGTGGCTCACGCCTGTAATCC
CAGAGGCTGAGGCAGGCGGATCACCTGAGGTGGGAGTTCTGGGATCAGCCTGACCAACAT
GGAGAAACCCTACTGGAAATACAAAGTTAGCCAGGCATGGTGGTGCATGCCTGTAGTCCC
AGCTGCTCAGGAGCCTGGCAACAAGAGCAAACTCCAGCTCAAAAAAAAAAAAAAAAAA

55/392

FIGURE 54

MGTKAQVERKLLCLFILAILLCSLALGSVTVHSSEPEVRIPENNPVKLS CAYSGFSSPRV
EWKFDQGD TTRLVCYNNKITASYEDRV TFLPTGITFKSVTREDTGT YTCMVSEEGNSYG
EVKVKLIVLVPPSKPTVNI PSSATIGNRAVLTCSEQDGSPPSEYTWFKDGI VMPTNPKST
RAFSNSSYVLNPTTGELVFDPLSASDTGEYSCEARNGYGTPMTSNAVRMEAVERNVGVIV
AAVLVTLILLGILVFGIWFAYS RGHFDR TKKGTSSKKVIYSQPSARSEGEFKQTSSFLV

Signal sequence:
amino acids 1-27

Transmembrane domain:
amino acids 238-255

N-glycosylation site:
amino acids 185-189

cAMP- and cGMP-dependent protein kinase phosphorylation site:
amino acids 270-274

Casein kinase II phosphorylation site:
amino acids 34-38, 82-86, 100-104, 118-122, 152-156,
154-158, 193-197, 203-207, 287-291

N-myristoylation site:
amino acids 105-111, 116-122, 158-164, 219-225, 237-243,
256-262

56/392

FIGURE 55

GTTGTGTCCTTCAGCAAAACAGTGGATTTAAATCTCCTTGCACAAGCTTGAGAGCAACAC
AATCTATCAGGAAAGAAAGAAAGAAAAAACCGAACCTGACAAAAAGAAGAAAAAGAAG
AAGAAAAAAATCATGAAAACCATCCAGCCAAAAATGCACAATTCTATCTCTTGGGCAAT
CTTCACGGGGCTGGCTGCTCTGTGTCTCTTCCAAGGAGTGCCCGTGCGCAGCGGAGATGC
CACCTTCCCCAAAGCTATGGACAACGTGACGGTCCGGCAGGGGGAGAGCGCCACCCTCAG
GTGCACTATTGACAACCGGGTCACCCGGGTGGCCTGGCTAAACCGCAGCACCATCCTCTA
TGCTGGGAATGACAAGTGGTGCCTGGATCCTCGCGTGGTCCTTCTGAGCAACACCCAAAC
GCAGTACAGCATCGAGATCCAGAACGTGGATGTGTATGACGAGGGCCCTTACACCTGCTC
GGTGCAGACAGACAACCACCCAAAGACCTCTAGGGTCCACCTCATTGTGCAAGTATCTCC
CAAATTGTAGAGATTTCTTCAGATATCTCCATTAATGAAGGGAACAATATTAGCCTCAC
CTGCATAGCAACTGGTAGACCAGAGCCTACGGTTACTTGGAGACACATCTCTCCAAAGC
GGTTGGCTTTGTGAGTGAAGACGAATACTTGGAATTCAGGGCATCACCCGGGAGCAGTC
AGGGGACTACGAGTGCAGTGCCTCCAATGACGTGGCCGCGCCCGTGGTACGGAGAGTAAA
GGTCACCGTGAACCTATCCACCATACATTTTCAGAAGCCAAGGGTACAGGTGTCCCCGTGGG
ACAAAAGGGGACACTGCAGTGTGAAGCCTCAGCAGTCCCCTCAGCAGAATTCCAGTGGTA
CAAGGATGACAAAAGACTGATTGAAGGAAAGAAAGGGGTGAAAGTGGAAAACAGACCTTT
CCTCTCAAACTCATCTTCTTCAATGTCTCTGAACATGACTATGGGAACTACACTTGCCT
GGCCTCCAACAAGCTGGGCCACACCAATGCCAGCATCATGCTATTTGGTCCAGGCGCCGT
CAGCGAGGTGAGCAACGGCACGTGAGGAGGGCAGGCTGCGTCTGGCTGCTGCCTCTTCT
GGTCTTGCACCTGCTTCTCAAATTTTGATGTGAGTGCCACTTCCCCACCCGGGAAAGGCT
GCCGCCACCACCACCACCAACACAACAGCAATGGCAACACCGACAGCAACCAATCAGATA
TATACAAATGAAATTAGAAGAAACACAGCCTCATGGGACAGAAATTTGAGGGAGGGGAAC
AAAGAATACTTTGGGGGGAAAAGAGTTTTTAAAAAAGAAATTGAAAATTGCCTTGCAGATA
TTTAGGTACAATGGAGTTTTCTTTTCCCAAACGGGAAGAACACAGCACACCCGGCTTGGA
CCCACTGCAAGCTGCATCGTGCAACCTCTTTGGTGCCAGTGTGGGCAAGGGCTCAGCCTC
TCTGCCCACAGAGTGCCCCACGTGGAACATTCTGGAGCTGGCCATCCCAAATTCATCA
GTCCATAGAGACGAACAGAATGAGACCTTCCGGCCCAAGCGTGGCGCTGCGGGCACTTTG
GTAGACTGTGCCACCACGGCGTGTGTTGTGAAACGTGAAATAAAAAGAGCAAAAAAAA

57/392

FIGURE 56

MKTIQPKMHNSISWAI FTGLAALCLFQGVPRSGDATFPKAMDNVTVRQGESATLRCTID
NRVTRVAWLNRSTILYAGNDKWCLDPRVVLLSNTQTQYSIEIQNV DVYDEGPYTCSVQTD
NHPKTSRVHLIVQVSPKIVEISSDISINEGNNISLT CIATGRPEPTVTWRHISPKAVGFV
SEDEYLEIQGITREQSGDYEC SASNDVAAPVVRVKVT VNYPPYISEAKGTGVPVGQKGT
LQCEASAVPSAEFQWKDDKRLIEGKKGVKVENRPFLSKLI FFNVSEHDYGN YTCVASNK
LGHTNASIMLFGPGAVSEVSNGTSRRAGCVWLLPLLVLHLL LKF

58/392

FIGURE 57

GCTGCGCCGGCTGCGGCTGCAGGGGAATCCGCTGTGGTGCGGCTGCCAGGCGCGGCCCT
ACTCGAGTGGCTGGCGCGGGCGCGCGTGCCTCGGACGGCGCGTGCCAGGGGCGCGGCG
CCTGCGGGGCGAGGCTCTGGACGCCCTGCGGCCCTGGGACCTGCGCTGCCCTGGGGACGC
GGCGCAGGAAGAGGAAGAGCTGGAAGAGCGGGCTGTGGCCGGGCCCCGCGCCCCCTCCGCG
CGGCCCTCCGCGCGGCCCCGGGGAGGAGCGGGCAGTCGCGCCTTGCCCTCGCGCTGCGT
GTGCGTCCCCGAGTCCCCGGCACAGCAGCTGCGAGGGCTGCGGCCTGCAGGCGGTGCCCCG
CGGCTTCCCCAGCGACACCAGCTCCTGGACCTGAGGCGGAACCACTTCCCCTCGGTGCC
CCGAGCGGCCTTCCCCGGNCTGGGCCACCTGGTGTGCTGCACCTGCAGCACTGCGGCAT
CGCGGAGCTGGAAGCGGGCGCCCTGGCCGGGCTGGGCCCGCTGATCTACCTGTACCTCTC
CGACAACCAGCTCGCAGGCCTCAGCGCTGCTGCCCTTGAAGGGGCTCCCCGCTCGGCTA
CCTGTACCTAGAACGCAACCGTTTCTGTCAGGTGCCAGGGGCTGCCNTGCGCGCCCTGCC
CAGCCTCTTCTCCCTGCACCTGCAGGACAACGCTGTGGACCGCCTGGCACCTGGGGACCT
GGGGAGAACACGGGCCTTGCGCTGGGTCTACCTGAGTGGAACCGCATCACCGAAGTGTC
CCTTGGGGCGCTGGGCCAGCTCGGGAGCTGGAGAAGCTGCACCTGGACAGGAATCAGCT
GCGAGAGGTGCCCACTGGGGCCTTGAGGGGCTGCCTGCCCTCCTGGAGCTGCAGCTCTC
GGGCAACCCACTCAGGGCCTTGCGTGACGGAGCCTTCCAGCCTGTGGGCAGGTGCTGCA
GCACCTCTTCTGAACAGCAGTGGCCTGGAGCAGATTTGTCTTGGGGCCTTTTCAGGCCT
GGGGCCCCGGGCTCCAGAGCCTGCACCTGCAGAAGAACCAGCTTCGGGCCCTGCCTGCCCT
GCCCAGTCTCAGCCAGCTGGAGCTCATCGACCTCAGCAGCAATCCCTTCCCCTGTGACTG
CCAGCTGCTTCCGCTGCACAGGTGGCTTACTGGGCTGAACCTGCGGGTGGGGGCCACCTG
CGCCACCCCTCCCAATGCCCGTGGCCAGAGGGTGAAGGCTGCAGCTGCTGTCTTTGAAGA
CTGCCCCGGGCTGGGCTGCCAGAAAGGCCAAGCGGACACCAGCCTCCAGGCCAGTGCCAG
GAGAACCCCCATCAAAGGAAGACAGTGTGGAGCAGATAAGAACATCCTCTTCCCCACATG
GTACCACACTGTGGAGCCCACCTCGCTGTCATAGGCCTGCGGCTCTGAAGGATGGCTTTG
CCCGCTCCCGCTCTGCCCCCTCAAGTGGAACCCAAGCTGGGCTCAGAATCTGTAGAGTGAG
GCCCCACCAAGGGAAACGACACCCACGGCCTGAGAGCCAGGTGGAGTCTGCCACTCAGC
TGCCTGCCTTTGCTCCCACCCTCTCCACCCCTCAAAGAGGTCTCGAGGGGACACTCTGAA
GGCACCTGGCTCAGAACCCTGCCATCCAAGGAGCGAGGAGTCCAGGGCTGAGCAAATG
CAGCGGGGAGGTGCGCAGTCCCCCTGCTTCCCGATCCTCATTTTCTGCTTCACTTGACTC
CTCCAGATAGGAGCTGCTCTCACTGCCACACTGCTG

59/392

FIGURE 58

LRRLRLQGNPLWCGCQARPLLEWLLARARVRS DGACQGPRRLRGEALDALRPWDLRCPGDA
AQEEEELEERAVAGPRAPPRGPPRGPGEEERAVAPCPACVCVPESRHSSCEGCGLQAVPR
GFPSDTQLLDLRRNHFPSPRAAFPGLGHLVSLHLQHCGIAELEAGALAGLGRLIYLYLS
DNQLAGLSAAALEGAPRLGYLYLERNRFLQVPGAAXRALPSLFSLHLQDNAVDRLAPGDL
GRTRALRWVYLSGNRITEVSLGALGPARELEKLHLDNRNQLREVPTGALEGLPALLELQLS
GNPLRALRDGAFQPVGRSLQHLFLNSSGLEQICPGAFSGLGPGQLSLHLQKNQLRALPAL
PSLSQLELIDLSSNPFPCDCQLLPLHRWLTGLNLRVGATCATPPNARGQRVKAAA VFED
CPGWAARKAKRTPASRPSARRTPIKGRQCGADKNILFPTWYHTVEPTSLS

Signal sequence:

None

Transmembrane domain:

None

N-glycosylation site:

325-328

Glycosaminoglycan attachment site:

338-341

Protein kinase C phosphorylation site:

438-440

N-myristoylation site:

166-171, 186-191, 253-258, 286-291, 335-340, 339-344, 450-455

Leucine rich repeat N-terminal domain:

94-123

Leucine Rich Repeat:

125-148, 149-172, 173-196, 197-220, 221-244, 245-268, 269-292, 293-316, 318-341, 343-364, 365-386

Leucine rich repeat C-terminal domain:

374-422

60/392

FIGURE 59

CTCCACGGTGTCCAGCGCCAGAAATGCGGCTTCTGGTCCTGCTATGGGGTTGCCTGCTG
CTCCAGGTTATGAAGCCCTGGAGGGCCCAGAGGAAATCAGCGGGTTCGAAGGGGACACT
GTGTCCCTGCAGTGCACCTACAGGGAAGAGCTGAGGGACCACCGGAAGTACTGGTGCAGG
AAGGGTGGGATCCTCTTCTCTCGCTGCTCTGGCACCATCTATGCAGAAGAAGAAGGCCAG
GAGACAATGAAGGGCAGGGTGTCCATCCGTGACAGCCGCCAGGAGCTCTCGCTCATTGTG
ACCCTGTGGAACCTCACCTGCAAGACGCTGGGGAGTACTGGTGTGGGGTCGAAAAACGG
GGCCCCGATGAGTCTTTACTGATCTCTCTGTTCTGTTCTTTCCAGGACCCTGCTGTCTCTCC
TCCCTTCTCCCACCTTCCAGCCTCTGGCTACAACACGCCTGCAGCCCCAAGGCAAAAGCT
CAGCAAACCCAGCCCCCAGGATTGACTTCTCCTGGGCTCTACCCGGCAGCCACCACAGCC
AAGCAGGGGAAGACAGGGGCTGAGGCCCTCCATTGCCAGGGACTTCCAGTACGGGCAC
GAAAGGACTTCTCAGTACACAGGAACCTCTCCTCACCAGCGACCTCTCCTCCTGCAGGG
AGCTCCCGCCCCCCCCATGCAGCTGGACTCCACCTCAGCAGAGGACACCAGTCCAGCTCTC
AGCAGTGGCAGCTCTAAGCCCAGGGTGTCCATCCCGATGGTCCGCATACTGGCCCCAGTC
CTGGTGTCTGCTGAGCCTTCTGTGAGCCGACGGCCTGATCGCCTTCTGCAGCCACCTGCTC
CTGTGGAGAAAGGAAGCTCAACAGGCCACGGAGACACAGAGGAACGAGAAGTTCTGGCTC
TCACGCTTGACTGCGGAGGAAAAGGAAGCCCCCTTCCAGGCCCTGAGGGGGACGTGATC
TCGATGCCTCCCCCTCCACACATCTGAGGAGGAGCTGGGCTTCTCGAAGTTTGTCTCAGCG
TAGGGCAGGAGGCCCTCCTGGCCAGGCCAGCAGTGAAGCAGTATGGCTGGCTGGATCAGC
ACCGATTCCCGAAAGCTTTCACCTCAGCCTCAGAGTCCAGCTGCCCGGACTCCAGGGCT
CTCCCCACCCTCCCCAGGCTCTCCTCTTGCACTGTTCCAGCCTGACCTAGAAGCGTTTGTCT
AGCCCTGGAGCCCAGAGCGGTGGCCTTGCTCTTCCGGCTGGAGACTGGGACATCCCTGAT
AGGTTACATCCCTGGGCAGAGTACCAGGCTGCTGACCCTCAGCAGGGCCAGACAAGGCT
CAGTGGATCTGGTCTGAGTTTCAATCTGCCAGGAACCTCCTGGGCCTCATGCCCAGTGTCTG
GACCCTGCCTTCCCTCCCACTCCAGACCCACCTTGTCTTCCCTCCCTGGCGTCTCTCAGAC
TTAGTCCCACGGTCTCCTGCATCAGCTGGTGATGAAGAGGAGCATGCTGGGGTGAGACTG
GGATTCTGGCTTCTCTTTGAACCACCTGCATCCAGCCCTTCCAGGAAGCCTGTGAAAAACG
TGATTCTGGCCCCACCAAGACCCACCAAAACCATCTCTGGGCTTGGTGCAGGACTCTGA
ATTCTAACAATGCCCAGTGACTGTGCACTTGAGTTTGAGGGCCAGTGGGCCTGATGAAC
GCTCACACCCCTTCCAGCTTAGAGTCTGCATTTGGGCTGTGACGTCTCCACCTGCCCAAT
AGATCTGCTCTGTCTGCGACACCAGATCCACGTGGGGACTCCCCTGAGGCCTGCTAAGTC
CAGGCCTTGGTGCAGGTGAGGTGCACATTGCAGGATAAGCCCAGGACCGGCACAGAAGTGG
TTGCCTTTNCCATTGCCCCCTCCCTGNNCCATGCCTTCTTGCCCTTGGAAAAAATGATGAA
GAAAACCTTGGCTCCTTCTTGTCTGGAAAGGGTTACTTGCCCTATGGGTTCTGGTGGCTA
GAGAGAAAAGTAGAAAACAGAGTGCACGTAGGTGTCTAACACAGAGGAGAGTAGGAACA
GGCGGATACCTGAAGGTGACTCCGAGTCCAGCCCCCTGGAGAAGGGGTGCGGGGTGGTG
GTAAAGTAGCACAACTACTATTTTTTTTCTTTTCCATTATTATTGTTTTTTAAGACAGA
ATCTCGTGTCTGCTGCCAGGCTGGAGTGCAGTGGCACGATCTGCAAACCTCCGCCTCCTGG
GTTCAAGTGATTCTTCTGCCTCAGCCTCCCGAGTAGCTGGGATTACAGGCACGCACCACC
ACACCTGGCTAATTTTTGTACTTTTAGTAGAGATGGGGTTTACCAGTGTGGCCAGGCTG
GTCTTGAACCTCTGACCTCAAATGAGCCTCCTGCTTCAGTCTCCCAAATTGCCGGGATTA
CAGGCATGAGCCACTGTGTCTGGCCCTATTTCTTTTAAAAAGTGAAATTAAGAGTTGTTT
AGTATGCAAACTTGGAAAGATGGAGGAGAAAAAGAAAAGGAAGAAAAAATGTCACCCA
TAGTCTCACCAGAGACTATCATTATTTCTGTTTGTGTTGTTTCTTCTTCCACTCTTTCTTC
TTCACATAATTTGCCGGTGTCTTTTTTACAGAGCAATTATCTTGTATATACAACCTTTGTA
TCCTGCCTTTTCCACCTTATCGTTCCATCACTTTATTCCAGCACTTCTCTGTGTTTTACA
GACCTTTTTTATAAATAAAATGTTTCATCAGCTGCATAAAAAAAAAAAAAA

61/392

FIGURE 60

MRLLVLLWGCLLLPGYEALLEGPEEISGFEGDTVSLQCTYREELRDHRKYWCRKGGILFSR
CSGTIYAEEEGQETMKGRVSIRDSRQELSLIVTLWNLTLDAGEYWCGVEKRGPDSESLI
SLFVFPGPCCPPSPSPPTFQPLATTRLQPKAKAQQTQPPGLTSPGLYPAATTAKQKGTGAE
APPLPGTSQYGHERTSQYTGTSPHPATSPAGSSRPPMQLDSTSAEDTSPALSSGSSKPR
VSIPMVRILAPVLVLLSLLSAAGLIAFCSHLLLRKEAQQATETQRNEKFWLSRLTAEK
EAPSQAPEGDVISMPLHTSEEEELGFSKFVSA

Important features:**Signal peptide:**

amino acids 1-17

Transmembrane domain:

amino acids 248-269

N-glycosylation site:

amino acids 96-99

Fibrinogen beta and gamma chains C-terminal domain:

amino acids 104-113

Ig like V-type domain:

amino acids 13-128

62/392

FIGURE 61

CGGGCCAGCCTGGGGCGGCCGGCCAGGAACCAACCCGTTAAGGTGTCTTCTCTTTAGGGAT
GGTGAGGTTGGAAAAAGACTCCTGTAACCCTCCTCCAGGATGAACCACCTGCCAGAAGAC
ATGGAGAACGCTCTCACCGGGAGCCAGAGCTCCCATGCTTCTCTGCGCAATATCCATTCC
ATCAACCCACACAACCTCATGGCCAGGATTGAGTCCATGAAGGAAGGGAAAAAGAAAGGC
ATATCTGATGTCAGGAGGACTTTCTGTTTGTGTTGTCACCTTTGACCTCTTATTCGTAACA
TTACTGTGGATAATAGAGTTAAATGTGAATGGAGGCATTGAGAACACATTAGAGAAGGAG
GTGATGCAGTATGACTACTATTCTTCATATTTTGGATATATTTCTTCTGGCAGTTTTTTCGA
TTTAAAGTGTTAATACTTGCATATGCTGTGTGCAGACTGCGCCATTGGTGGGCAATAGCG
TTGACAACGGCAGTGACCAGTGCCTTTTTACTAGCAAAAGTGATCCTTTTCGAAGCTTTTC
TCTCAAGGGGCTTTTGGCTATGTGCTGCCCATCATTTTCATTCATCCTTGCCCTGGATTGAG
ACGTGGTTCCCTGGATTTCAAAGTGTTACCTCAAGAAGCAGAAGAAGAAAACAGACTCCTG
ATAGTTCAGGATGCTTCAGAGAGGGCAGCACTTATACCTGGTGGTCTTTCTGATGGTCAG
TTTTATTTCCCTCCTGAATCCGAAGCAGGATCTGAAGAAGCTGAAGAAAACAGGACAGT
GAGAAACCACTTTTAGAACTATGAGTACTACTTTTGTTAAATGTGAAAAACCTTCACAGA
AAGTCATCGAGGCAAAAAGAGGCAGGCAGTGAGTCTCCCTGTCGACAGTAAAGTTGAAA
TGGTGACGTCCACTGCTGGCTTTATTGAACAGCTAATAAAGATTTATTTATTGTAATACC
TCACAAACGTTGTACCATATCCATGCACATTTAGTTGCCTGCCTGTGGCTGGTAAGGTAA
TGTCATGATTCATCCTCTCTTCAGTGAGACTGAGCCTGATGTGTAAACAAATAGGTGAAG
AAAGTCTTGTGCTGTATTCCTAATCAAAAGACTTAATATATTGAAGTAACACTTTTTTTAG
TAAGCAAGATACCTTTTTTATTTCAATTCACAGAATGGAATTTTTTTTGTTCATGTCTCAG
ATTTATTTTGTATTTCTTTTTTAACACTCTACATTTCCCTTGTTTTTTAACTCATGCACA
TGTGCTCTTTGTACAGTTTTTAAAAAGTGTAATAAAATCTGACATGTCAATGTGGCTAGTT
TTATTTTTCTTGTTTTGCATTATGTGTATGGCCTGAAGTGTTGGACTTGCAAAAGGGGAA
GAAAGGAATTGCGAATACATGTAAAATGTCACCAGACATTTGTATTATTTTTATCATGAA
ATCATGTTTTTCTCTGATTGTTCTGAAATGTTCTAAATACTCTTATTTTGAATGCACAAA
ATGACTTAAACCATTCATATCATGTTTCCTTTGCGTTCAGCCAATTTCAATTAAAATGAA
CTAAATTAAAAA

63/392

FIGURE 62

MNHLPEDMENALTGSQSSHASLRNIHSINPTQLMARIESYEGREKKGISDVRRTFCLFVT
FDLLFVTLLWIIELNVNGGIENTLEKEVMQYDYYSSYFDIFLLAVFRFKVLILAYAVCRL
RHWWAIALTTAVTSAFLLAKVILSKLFSQGAFGYVLPPIISFILAWIETWFLDFKVLPOEA
EEENRLLIVQDASERAALIPGGLSDGQFYSPPESEAGSEEAEKQDSEKPLLEL

Important features of the protein:**Signal peptide:**

amino acids 1-20

Transmembrane domains:

amino acids 54-72, 100-118, 130-144, 146-166

N-myristoylation sites:

amino acids 14-20, 78-84, 79-85, 202-208, 217-223

64/392

FIGURE 63

GCGCCGGGAGCCCATCTGCCCCAGGGGCACGGGGCGCGGGGCCGGCTCCCGCCCGGCAC
ATGGCTGCAGCCACCTCGCGCGCACCCCGAGGCGCCGCGCCAGCTCGCCCGAGGTCCGT
CGGAGGCGCCCGGCCGCCCCGGAGCCAAGCAGCAACTGAGCGGGGAAGCGCCCGCGTCCG
GGGATCGGGATGTCCTCCTCCTTCTCCTCTTGCTAGTTTCTACTATGTTGGAACCTTG
GGGACTCACACTGAGATCAAGAGAGTGGCAGAGGAAAAGGTCACTTTGCCCTGCCACCAT
CAACTGGGGCTTCCAGAAAAAGACACTCTGGATATTGAATGGCTGCTCACCGATAATGAA
GGGAACCAAAAAGTGGTGATCACTTACTCCAGTCGTATGTCTACAATAACTTGACTGAG
GAACAGAAGGGCCGAGTGGCCTTTGCTTCCAATTTCTTGGCAGGAGATGCCTCCTTGAG
ATTGAACCTCTGAAGCCAGTGATGAGGGCCGGTACACCTGTAAGGTTAAGAATTCAGGG
CGCTACGTGTGGAGCCATGTCATCTTAAAAGTCTTAGTGAGACCATCCAAGCCCAAGTGT
GAGTTGGAAGGAGAGCTGACAGAAGGAAGTGACCTGACTTTGCAGTGTGAGTCATCCTCT
GGCACAGAGCCCATTTGTGTATTACTGGCAGCGAATCCGAGAGAAAGAGGGAGAGGATGAA
CGTCTGCCTCCCAAATCTAGGATTGACTACAACCACCCTGGACGAGTTCTGCTGCAGAAT
CTTACCATGTCTTACTCTGGACTGTACCAGTGACAGCAGGCAACGAAGCTGGGAAGGAA
AGCTGTGTGGTGCGAGTAACTGTACAGTATGTACAAAGCATCGGCATGGTTGCAGGAGCA
GTGACAGGCATAGTGGCTGGAGCCCTGCTGATTTTCTCTTGGTGTGGCTGCTAATCCGA
AGGAAAGACAAAGAAAGATATGAGGAAGAAGAGAGACCTAATGAAATTCGAGAAGATGCT
GAAGCTCCAAAAGCCCGTCTTGTGAAACCCAGCTCCTCTTCTCAGGCTCTCGGAGCTCA
CGCTCTGGTTCTTCTCCTCCTCGCTCCACAGCAAATAGTGCCTCACGCAGCCAGCGGACA
CTGTCAACTGACGCAGCACCCAGCCAGGGCTGGCCACCCAGGCATACAGCCTAGTGGGG
CCAGAGGTGAGAGGTTCTGAACCAAGAAAGTCCACCATGCTAATCTGACCAAAGCAGAA
ACCACACCCAGCATGATCCCCAGCCAGAGCAGAGCCTTCCAAACGGTCTGAATTACAATG
GACTTGACTCCCACGCTTTCTTAGGAGTCAGGGTCTTTGGACTCTTCTCGTCATTGGAGC
TCAAGTCACCAGCCACACAACCAGATGAGAGGTGATCTAAGTAGCAGTGAGCATTGCACG
GAACAGATTGAGATGAGCATTCTTCTTATACAATACCAAACAAGCAAAAGGATGTAAGCT
GATTCATCTGTAAAAGGCATCTTATTGTGCCTTTAGACCAGAGTAAGGGAAAGCAGGAG
TCCAAATCTATTTGTTGACCAGGACCTGTGGTGAGAAGGTTGGGGAAAGGTGAGGTGAAT
ATACCTAAAACCTTTTAATGTGGGATATTTTGTATCAGTGCTTTGATTCACAATTTTCAAG
AGGAAATGGGATGCTGTTTGTAAATTTTCTATGCATTTCTGCAAACTTATTGGATTATTA
GTTATTGAGACAGTCAAGCAGAACCCACAGCCTTATTACACCTGTCTACACCATGTACTG
AGCTAACCACTTCTAAGAACTCCAAAAAAGGAAACATGTGTCTTCTATTCTGACTTAAC
TTCATTTGTGATAAGGTTTGGATATTAATTTCAAGGGGAGTTGAAATAGTGGGAGATGGA
GAAGAGTGAATGAGTTTCTCCCACTCTATACTAATCTCACTATTTGTATTGAGCCCAAAA
TAACTATGAAAGGAGACAAAAATTTGTGACAAAGGATTGTGAAGAGCTTTCCATCTTCAT
GATGTTATGAGGATTGTTGACAAACATTAGAAATATATAATGGAGCAATTGTGGATTTC
CCTCAAATCAGATGCCCTCTAAGGACTTTCTTGCTAGATATTTCTGGAAGGAGAAAATACA
ACATGTCATTTATCAACGTCCTTAGAAAGAATTCTTCTAGAGAAAAAGGGATCTAGGAAT
GCTGAAAGATTACCCAACATAACATTATAGTCTCTTCTTTCTGAGAAAATGTGAAACCAG
AATTGCAAGACTGGGTGGACTAGAAAGGGAGATTAGATCAGTTTTCTCTTAATATGTCAA
GGAAGGTAGCCGGGCATGGTGCCAGGCACCTGTAGGAAAATCCAGCAGGTGGAGGTTGCA
GTGAGCCGAGATTATGCCATTGCACTCCAGCCTGGGTGACAGAGCGGGACTCCGTCTC

65/392

FIGURE 64

MSLLLLLLLLVSYYVGTGLGTHTEIKRVAEEKVTLPCHHQLGLPEKDTLDIEWLLTDNEGNQ
KVVITYSSRHVYNNLTTEEQGRVAFASNFLAGDASLQIEPLKPSDEGRYTCKVKNSGRYV
WSHVILKVLVRPSKPKCELEGELTEGSDLTQCESSSGTEPIVYYWQRIREKEGEDERLP
PKSRIDYNHPGRVLLQNLTMSYSGLYQCTAGNEAGKESCVVRVTVQYVQSIGMVAGAVTG
IVAGALLIFLLVWLLIRRKDKERYEEEEERPNEIREDAEAPKARLVKPPSSSSSGSRSSRSG
SSSTRSTANSASRSQRTLSTDAAPQPGLATQAYSLVGPEVRGSEPKKVHHANLTKAETTP
SMIPSQSRAFQTV

Signal sequence:
amino acids 1-16

Transmembrane domain:
amino acids 232-251

66/392

FIGURE 65

GTCTGGGGCTGCGCGACGGCGCAGGGGCTGCGGGGAGCGCCGCGCAGGCCGTGCAGTTCCT
AGCGAGGAGGCGCCGCGCCGCGCCATTGCCGCTCTCTCGGTGAGCGCAGCCCCGCTCTCCGGGC
CGGGCCTTCGCGGGCCACCGGCGCCATGGGGCCAGTGCGGCATCACCTCCTCCAAGACCGT
GCTGGTCTTTCTCAACCTCATCTTCTGGGGGGCAGCTGGCATTATGCTATGTGGGAGC
CTATGTCTTCATCACTTATGATGACTATGACCACTTCTTTGAAGATGTGTACACGCTCAT
CCCTGCTGTAGTGATCATAGCTGTAGGAGCCCTGCTTTTCATCATTGGGCTAATTGGCTG
CTGTGCCACAATCCGGGAAAGTCGCTGTGGACTTGCCACGTTTGTTCATCATCCTGCTCTT
GGTTTTTGTTCACAGAAGTTGTTGTAGTGGTTTTGGGATATGTTTACAGAGCAAAGGTGGA
AAATGAGGTTGATCGCAGCATTAGAAAGTGTATAAGACCTACAATGGAACCAACCCTGA
TGCTGCTAGCCGGGCTATTGATTATGTACAGAGACAGCTGCATTGTTGTGGAATTCACAA
CTACTCAGACTGGGAAAATACAGATTGGTTCAAAGAAACCAAAAACCAGAGTGTCCCTCT
TAGCTGCTGCAGAGAGACTGCCAGCAATTGTAATGGCAGCCTGGCCCACCCTTCCGACCT
CTATGCTGAGGGGTGTGAGGCTCTAGTAGTGAAGAAGCTACAAGAAATCATGATGCATGT
GATCTGGGCCGCACTGGCATTTCAGCTATTGAGCTGCTGGGCATGCTGTGTGCTTGCAT
CGTGTGTGCAGAAGGAGTAGAGATCCTGCTTACGAGCTCCTCATCACTGGCGGAACCTA
TGCATAGTTGACAACCTCAAGCCTGAGCTTTTTGGTCTTGTTCTGATTTGGAAGGTGAATT
GAGCAGGTCTGCTGCTGTTGGCCTCTGGAGTTCATTTAGTTAAAGCACATGTACACTGGT
GTTGGACAGAGCAGCTTGGCTTTTCATGTGCCACCTACTTACCTACTACCTGCGACTTT
CTTTTTCTTGTTCTAGCTGACTCTTCATGCCCCTAAGATTTTAAGTACGATGGTGAACG
TTCTAATTTCAGAACCAATTGCGAGTCATGTAGTGTGGTAGAATTAAAGGAGGACACGAG
CCTGCTTCTGTTACCTCCAAGTGGTAACAGGACTGATGCCGAAATGTCACCAGGTCTTT
CAGTCTTCACAGTGGAGAACTCTTGGCCAAAGTTTTTGCGGGGAGGAGGAGGAAACCAG
CTTTCTGGTTAAGGTTAACACCAGATGGTGCCCCCTCATTGGTGTCTTTTAAAAAATATT
TACTGTAGTCCAATAAGATAGCAGCTGTACAAAATGACTAAAATAGATTGTAGGATCATA
TGGCGTATATCTTGGTTTCATCTTCAAAATCAGAGACTGAGCTTTGAAACTAGTGGTTTTT
AATCAAAGTTGGCTTTATAGGAGGAGTATAATGTATGCACTACTGTTTTAAAGAATTAG
TGTGAGTGTGTTTTTGTATGAATGAGCCCATTGATGGTAAGTCTTAAGCTTGTGGAAAT
AATGTACCCATGTAGACTAGCAAAATAGTATGTAGATGTGATCTCAGTTGTAAATAGAAA
AATCTAATTCAATAAACTCTGTATCAGCCCCCAAAAAAAAAAAAAAAAAA

67/392

FIGURE 66

MGQCGITSSKTVLVFLNLI FWGAAGILCYVGAYVFITYDDYDHFFEDVYTLIPAVVIIAV
GALLFIIGLIGCCATIRESRCGLATFVIIILLVFVTEVVVVVLGYVYRAKVENEVDRSIQ
KVYKTYNGTNPDAASRAIDYVQRQLHCCGIHNYSDWENTDWFKETKNQSVPLSCCRETAS
NCNGSLAHPSDLYAEGCEALVVKKLQEIMMHVIWAALAFAAIQLLGMLCACIVLCRRSRD
PAYELLITGGTYA

Signal peptide:

none

Type II transmembrane domain:

11-38

Other transmembrane domains:

48-68, 87-107, 208-235

N-glycosylation site:

127-131, 152-156, 167-171, 183-187

Tyrosine kinase phosphorylation site:

236-244

N-myristoylation site:

5-11, 68-74, 71-77, 226-232

Prokaryotic membrane lipoprotein lipid attachment site:

62-73, 221-232

Transmembrane 4 family proteins:

7-35, 56-106

68/392

FIGURE 67

GCGGCACCTGGAAGATGCGCCCATTTGGCTGGTGGCCTGCTCAAGGTGGTGTTCGTGGTCT
TCGCCTCCTTGTGTGCCTGGTATTCGGGGTACCTGCTCGCAGAGCTCATTCCAGATGCAC
CCCTGTCCAGTGCTGCCTATAGCATCCGCAGCATCGGGGAGAGGCCTGTCCTCAAAGCTC
CAGTCCCCAAAAGGCAAAAATGTGACCACTGGACTCCCTGCCCATCTGACACCTATGCCT
ACAGGTTACTCAGCGGAGGTGGCAGAAGCAAGTACGCCAAAATCTGCTTTGAGGATAACC
TACTTATGGGAGAACAGCTGGGAAATGTTGCCAGAGGAATAAACATTGCCATTGTCAACT
ATGTAACTGGGAATGTGACAGCAACACGATGTTTTGATATGTATGAAGGCGATAACTCTG
GACCGATGACAAAGTTTTATTTCAGAGTGCTGCTCCAAAATCCCTGCTCTTCATGGTGACCT
ATGACGACGGAAGCACAAAGACTGAATAACGATGCCAAGAATGCCATAGAAGCACTTGGAA
GTAAAGAAATCAGGAACATGAAATTCAGGTCTAGCTGGGTATTTATTGCAGCAAAGGCT
TGGAACCTCCCTTCCGAAATTCAGAGAGAAAAGATCAACCACTCTGATGCTAAGAACACA
GATATTCTGGCTGGCCTGCAGAGATCCAGATAGAAGGCTGCATACCCAAAGAACGAAGCT
GACTGTCAGGGTCCTGAGTAAATGTGTTCTGTATAAAACAAATGCAGCTGGAATCGCTCA
AGAATCTTATTTTTCTAAATCCAACAGCCCATATTTGATGAGTATTTTGGGTTTGTGTGTA
AACCAATGAACATTTGCTAGTTGTATCAAATCTTGGTACGCAGTATTTTATACCAGTAT
TTTATGTAGTGAAGATGTCAATTAGCAGGAACTAAAATGAATGGAAATTCCTAAAAAAA
AAA

69/392

FIGURE 68

MRPLAGGLLKVVVFVVFASLCAWYSGYLLAELIPDAPLSSAAYSIRSIGERPVLKAPVPKR
QKCDHWTPCPSDTYAYRLLSGGGRSKYAKICFEDNLLMGEQLGNVARGINIAIVNYVTGN
VTATRCFDMYEGDNSGPMTKFIQSAAPKSLLFMVTYDDGSTRLNNDKNAIEALGSKEIR
NMKFRSSWVFIAAKGLELPSEIQREKINHSDAKNNRYSGWPAEIQIEGCI PKERS

Signal sequence:

amino acids 1-20

N-glycosylation sites:

amino acids 120-124, 208-212

Glycosaminoglycan attachment site:

amino acids 80-84

N-myristoylation sites:

amino acids 81-87, 108-114, 119-125

70/392

FIGURE 69

ACACAACTTTACACCTGAATGAACGCCAAACCTCTATGGATATATAAAGGGAAGCTTGAG
GAGGAATTTACAGTTACAGTGCAGAAGCAGAAGCAAAAGAATTAACCAGCTCTTCAGTC
AAGCAAATCCTCTACTCACCATGCCTTCCTCCTGCCATTCATTTCTATCTCCTTCCCCTTG
CATGCATCCTAATGAAAAGCTGTTTGGCTTTTAAAAATGATGCCACAGAAATCCTTTATT
CACATGTGGTTAAACCTGTTCCAGCACACCCCAGCAGCAACAGCACGTTGAATCAAGCCA
GAAATGGAGGCAGGCATTTTCAGTAACACTGGACTGGATCGGAACACTCGGGTTCAAGTGG
GTTGCCGGGAAGCTGCGTTCCACCAAATACATCTCTGATGGCCAGTGCACCAGCATCAGCC
CTCTGAAGGAGCTGGTGTGTGCTGGCGAGTGCTTGCCCCTGCCAGTGCTCCCTAACTGGA
TTGGAGGAGGCTATGGAACAAAGTACTGGAGCAGGAGGAGCTCCCAGGAGTGGCGGTGTG
TCAATGACAAAACCCGTACCCAGAGAATCCAGCTGCAGTGCCAAGATGGCAGCACACGCA
CCTACAAAATCACAGTAGTCACTGCCTGCAAGTGCAAGAGGTACACCCGGCAGCACAAACG
AGTCCAGTCACAACTTTGAGAGCATGTACCTGCCAAGCCAGTCCAGCATCACAGAGAGC
GGAAAAGAGCCAGCAAATCCAGCAAGCACAGCATGAGTTAGAACTCAGACTCCCATAACT
AGACTTACTAGTAACCATCTGCTTTACAGATTTGATTGCTTGGAAGACTCAAGCCTGCCA
CTGCTGTTTTCTCACTTGAAAGTATATGCTTTCTGCTTTGATCAAACCCAGCAAGCTGTC
TTAAGTATCAGGACCTTCTTTGGGAATAGTTTTTCTTTTAAAGTTTTTCAAGATGTAGG
TATATCCATGAATGCAATTTGCATTTAAATTCACGTATCCCTGTAGTTTTAAATTCCTCA
TTGGTCTTAAAAGACTGTTGATACTATAAACATCAGTGGAATCAATTATATTTTAAACA
GAAAAGGGCTT

71/392

FIGURE 70

MLPPAIHFYLLPLACILMKSCLA FKNDAT EILYSHVVKPVP AHPSSNSTLNQARNGGRHF
SNTGLDRNTRVQVGCRELRSTKYISDGQCT SISPLKELVCAGECLPLPVL PNWIGGGYGT
KYWSRRSSQEWRCVNDKTRTQRIQLQCQDGSTR TYKITVVTACKCKRYTRQHNESSHNFE
SMSPAKPVQHHRRERKRASKSSKHSMS

Signal sequence:

1-23

Transmembrane domain:

None

N-glycosylation site:

47-50, 173-176

cAMP- and cGMP-dependent protein kinase phosphorylation site:

125-128, 166-169, 195-198

N-myristoylation site:

64-69, 87-92, 115-120, 116-121, 150-155

72/392

FIGURE 71

CCCAGGCTCTAGTGCAGGAGGAGAAGGAGGAGGAGCAGGAGGTGGAGATTCCCAGTTAAA
AGGCTCCAGAATCGTGTACCAGGCAGAGAACTGAAGTACTGGGGCCTCCTCCACTGGGTC
CGAATCAGTAGGTGACCCCGCCCCTGGATTCTGGAAGACCTCACCATGGGACGCCCCCGA
CCTCGTGCGGCCAAGACGTGGATGTTCTGCTCTTGCTGGGGGGAGCCTGGGCAGGACAC
TCCAGGGCACAGGAGGACAAGGTGCTGGGGGGTTCATGAGTGCCAACCCCATTCGCAGCCT
TGGCAGGCGGCCTTGTTCCAGGGCCAGCAACTACTCTGTGGCGGTGTCCTTGTAAGGTGGC
AACTGGGTCTTACAGCTGCCCCACTGTAAAAAACCGAAATACACAGTACGCCTGGGAGAC
CACAGCCTACAGAATAAAGATGGCCCAGAGCAAGAAATACCTGTGGTTCAGTCCATCCCA
CACCCCTGCTACAACAGCAGCGATGTGGAGGACCACAACCATGATCTGATGCTTCTTCAA
CTGCGTGACCAGGCATCCCTGGGGTCCAAAGTGAAGCCCATCAGCCTGGCAGATCATTGC
ACCCAGCCTGGCCAGAAGTGCACCGTCTCAGGCTGGGGCACTGTCAACAGTCCCCGAGAG
AATTTTCTTGACACTCTCAACTGTGCAGAAGTAAAAATCTTTCCCCAGAAGAAGTGTGAG
GATGCTTACCCGGGGCAGATCACAGATGGCATGGTCTGTGCAGGCAGCAGCAAAGGGGCT
GACACGTGCCAGGGCGATTCTGGAGGCCCCCTGGTGTGTGATGGTGCCTCCAGGGCATC
ACATCCTGGGGCTCAGACCCCTGTGGGAGGTCCGACAAACCTGGCGTCTATACCAACATC
TGCCGCTACCTGGACTGGATCAAGAAGATCATAGGCAGCAAGGGCTGATTCTAGGATAAG
CACTAGATCTCCCTTAATAAACTCACAACCTCTCTGGTTC

73/392

FIGURE 72

MGRPRPRAAKTWMFLLLLGGAWAGHSRAQEDKVLGGHECQPHSQPWQAALFQGQQLLCGG
VLVGGNWLTAAHCKKPKYTVRLGDHSLQNKDGPEQEIPVVQSIHPHCYNSSDVEDHNHD
LMLLQLRDQASLGSKVKPISLADHCTQPGQKCTVSGWGTVTSPRENF PDTLNCAEVKIFP
QKKCEDAYPGQITDGMVCAGSSKGADTCQGDSSGGLVCDGALQGITSWGSDPCGRSDKPG
VYTNICRYLDWIKKIIGSKG

Important Features:**Signal peptide:**

amino acids 1-23

Transmembrane domain:

amino acids 51-71

N-glycosylation site:

amino acids 110-113

Serine proteases, trypsin family, histidine active site:

amino acids 69-74 and 207-217

Tyrosine kinase phosphorylation site:

amino acids 182-188

Kringle domain proteins motif:

amino acids 205-217

74/392

FIGURE 73

CTCGGGCGCGCACAGGCAGCTCGGTTTGCCTTGCGATTGAGCTGCGGGTCGCGGCCGGCG
CCGGCCTCTCCAATGGCAAATGTGTGTGGCTGGAGGCGAGCGCGAGGCTTTCGGCAAAGG
CAGTCGAGTGTTTGCAGACCGGGGCGAGTCTGTGAAAGCAGATAAAAGAAAACATTTAT
TAACGTGTCATTACGAGGGGAGCGCCCGGGCGGGGCTGTGCGACTCCCCGCGGAACATTT
GGCTCCCTCCAGCTCCGAGAGAGGAGAAGAAGAAAGCGGAAAAGAGGCAGATTACGTCG
TTTCCAGCCAAGTGGACCTGATCGATGGCCCTCCTGAATTTATCACGATATTTGATTTAT
TAGCGATGCCCCCTGGTTTGTGTGTTACGCACACACACGTGCACACAAGGCTCTGGCTCG
CTTCCCTCCCTCGTTTCCAGCTCCTGGGCGAATCCCACATCTGTTTCAACTCTCCGCCGA
GGGCGAGCAGGAGCGAGAGTGTGTGCAATCTGCGAGTGAAGAGGGACGAGGGAAAAGAAA
CAAAGCCACAGACGCAACTTGAGACTCCCGCATCCCAAAGAAGCACCAGATCAGCAAAA
AAAGAAGATGGGGCCCCCGAGCCTCGTGCTGTGCTTGCTGTCCGCAACTGTGTTCTCCCT
GCTGGGTGGAAGCTCGGCCTTCTGTGCGACCACCGCCTGAAAGGCAGGTTTTCAGAGGGA
CCGCAGGAACATCCGCCCCAACATCATCTGGTGCTGACGGACGACCAGGATGTGGAGCT
GGGTTCCATGCAGGTGATGAACAAGACCCGGCGCATCATGGAGCAGGGCGGGGCGCACTT
CATCAACGCCTTCGTGACCACACCCATGTGCTGCCCCCTCACGCTCCTCCATCCTCACTGG
CAAGTACGTCCACAACCACAACCTACACCAACAATGAGAACTGCTCCTCGCCCTCCTG
GCAGGCACAGCACGAGAGCCGCACCTTTGCCGTGTACCTCAATAGCACTGGCTACCGGAC
AGCTTTCTTCGGGAAGTATCTTAATGAATAACAACGGCTCCTACGTGCCACCCGGCTGGAA
GGAGTGGGTGCGACTCCTTAAAACTCCCGCTTTTATACTACACGCTGTGTGCGAACGG
GGTGAAAGAGAAGCACGGCTCCGACTACTCCAAGGATTACCTCACAGACCTCATCACCAA
TGACAGCGTGAGCTTCTTCCGCACGTCCAAGAAGATGTACCCGCACAGGCCAGTCCTCAT
GGTCATCAGCCATGCAGCCCCCACGGCCCTGAGGATTCAGCCCCACAATATTCAGGCCT
CTTCCCAAACGCATCTCAGCACATCACGCCGAGCTACAACCTACGCGCCCAACCCGGACAA
ACACTGGATCATGCGCTACACGGGGCCCATGAAGCCCATCCACATGGAATTCACCAACAT
GCTCCAGCGGAAGCGCTTGACAGCCCTCATGTGCGGTGGACGACTCCATGGAGACGATTTA
CAACATGCTGGTTGAGACGGGCGAGCTGGACAACACGTACATCGTATACACCGCCGACCA
CGGTTACCACATCGGCCAGTTTGGCCTGGTGAAAGGGAAATCCATGCCATATGAGTTTGA
CATCAGGGTCCCGTTCTACGTGAGGGGCCCCAACGTGGAAGCCGGCTGTCTGAATCCCCA
CATCGTCTCAACATTGACCTGGCCCCCACCATCCTGGACATTGCAGGCCTGGACATACC
TGCGGATATGGACGGGAAATCCATCCTCAAGCTGCTGGACACGGAGCGGCCGGTGAATCG
GTTTCACTTGAAAAAGAAGATGAGGGTCTGGCGGGACTCCTTCTTGGTGGAGAGAGGCAA
GCTGCTACACAAGAGAGACAATGACAAGGTGGACGCCAGGAGGAGAACTTTCTGCCCAA
GTACCAGCGTGTGAAGGACCTGTGTGACGCTGCTGAGTACCAGACGGCGTGTGAGCAGCT
GGGACAGAAGTGGCAGTGTGTGGAGGACGCCACGGGGAAGCTGAAGCTGCATAAGTGCAA
GGGCCCCATGCGCTGGGCGGCAGCAGAGCCCTCTCCAACCTCGTGCCCAAGTACTACGG
GCAGGGCAGCGAGGCCTGCACCTGTGACAGCGGGGACTACAAGCTCAGCCTGGCCGGACG
CCGAAAAAACTCTTCAAGAAGAAGTACAAGGCCAGCTATGTCCGCAGTCGCTCCATCCG
CTCAGTGGCCATCGAGGTGGACGGCAGGGTGTACCACGTAGGCCTGGGTGATGCCGCCCA
GCCCCGAAACCTCACCAAGCGGCACTGGCCAGGGGCCCCCTGAGGACCAAGATGACAAGGA
TGGTGGGGACTTCAGTGGCACTGGAGGCCTTCCCGACTACTCAGCCGCCAACCCCATTA
AGTGACACATCGGTGCTACATCCTAGAGAACGACACAGTCCAGTGTGACCTGGACCTGTA
CAAGTCCCTGCAGGCCTGGAAAGACCACAAGCTGCACATCGACCACGAGATTGAAACCT
GCAGAACAAAATTAAGAACCTGAGGGAAGTCCGAGGTCACCTGAAGAAAAAGCGGCCAGA
AGAATGTGACTGTCAAAAATCAGCTACCACACCCAGCACAAAGGCCGCTCAAGCACAG
AGGCTCCAGTCTGCATCCTTTCAGGAAGGGCCTGCAAGAGAAGGACAAGGTGTGGCTGTT
GCGGGAGCAGAAGCGCAAGAAGAACTCCGCAAGCTGCTCAAGCGCCTGCAGAACACGA
CACGTGCAGCATGCCAGGCCTCACGTGCTTACCCACGACAACCAAGCACTGGCAGACGGC

75/392

GCCTTTCTGGACACTGGGGCCTTTCTGTGCCTGCACCAGCGCCAACAATAACACGTACTG
GTGCATGAGGACCATCAATGAGACTCACAATTCCTCTTCTGTGAATTTGCAACTGGCTT
CCTAGAGTACTTTGATCTCAACACAGACCCCTACCAGCTGATGAATGCAGTGAACACACT
GGACAGGGATGTCCCTCAACCAGCTACACGTACAGCTCATGGAGCTGAGGAGCTGCAAGGG
TTACAAGCAGTGTAACCCCCGGACTCGAAACATGGACCTGGATGGAGGAAGCTATGAGCA
ATACAGGCAGTTTCAGCGTCGAAAGTGGCCAGAAATGAAGAGACCTTCTTCCAAATCACT
GGGACAACTGTGGGAAGGCTGGGAAGGTTAAGAAACAACAGAGGTGGACCTCCAAAAACA
TAGAGGCATCACCTGACTGCACAGGCAATGAAAAACCATGTGGGTGATTTCCAGCAGACC
TGTGCTATTGGCCAGGAGGCCCTGAGAAAGCAAGCACGCACTCTCAGTCAACATGACAGAT
TCTGGAGGATAACCAGCAGGAGCAGAGATAACTTCAGGAAGTCCATTTTTTGCCCTGCTT
TTGCTTTGGATTATACCTCACCAGCTGCACAAAATGCATTTTTTTCGTATCAAAAAGTCAC
CACTAACCCTCCCCAGAAGCTCACAAAGGAAAACGGAGAGAGCGAGCGAGAGAGATTTT
CTTGGAATTTCTCCAAGGGCGAAAGTCATTGGAATTTTAAATCATAGGGGAAAAGCA
GTCCTGTTCTAAATCCTCTTATTCTTTTGGTTTGTACAAAGAAGGAACCTAAGAAGCAGG
ACAGAGGCAACGTGGAGAGGCTGAAAACAGTGCAGAGACGTTTGACAATGAGTCAGTAGC
ACAAAAGAGATGACATTTACCTAGCACTATAAACCTGGTTGCCTCTGAAGAACTGCCT
TCATTGTATATATGTGACTATTTACATGTAATCAACATGGGAACTTTTAGGGGAACCTAA
TAAGAAATCCCAATTTTCAGGAGTGGTGGTGTCAATAAACGCTCTGTGGCCAGTGTA
AAAAA

76/392

FIGURE 74

MGPPSLVLCLLSATVFSLLGGSSAFLSHHRLKGRFQDRRNIRPNIILVLTDDQDVELGS
MQVMNKTRRIMEQGAHFVAFVTTMCCPSRSSILTGKYVHNHNTYTNNECSPSWQA
QHESRTFAVYLNSTGYRTAFFGKYLNEYNGSYVPPGWKEWVGLLKNSRFYNYTLCRNGVK
EKHGSDYSKDYLTDLITNDSVSFFRTSKKMPHRPVLVISHAAPHGPEDSAPQYSRLFP
NASQHITPSYNYAPNPDKHWIMRYTGPMKPIHMEFTNMLQKRLQTLMSVDDSMETIYNM
LVETGELDNTYIVYTADHGYHIGQFGLVKGKSMPEYFDIRVPFYVRGPNVEAGCLNPHIV
LNIDLAPTILDIAGLDIPADM DGKSILKLLDTERPVNRFHLKKKMRVWRDSFLVERGKLL
HKRDNDKVDAQEENFLPKYQRVKDLQRAEYQTACEQLGQKWQCVEDATGKLLKHKCKGP
MRLGGSRALSNLVPKYQGQSEACTCDSDYKLSLAGRRKKLFKKKYKASYVRSRSIRSV
AIEVDGRVYHVGLGDAAPRNLTKRHWPGAPEDQDDKDGGDFSGTGGLPDYSAANPIKVT
HRCYILENDTVQCDLDLYKSLOAWKDHKLHIDHEIETLQNKIKNLREVRGHLKKKRPEEC
DCHKISYHTQHKGRCLKHRGSSLHPPFRKGLQEKDKVWLLREQKRKKKLRKLLKRLQNNDTC
SMPGLTCFTHDNQHWQTAPFWTLGPFCACTSANNTYWCMTINETHNFLFCEFATGFLE
YFDLNTDPYQLMNAVNTLDRDVLNQLHVQLMELRSCGYKQCNPRTNMDLDGGSYEQYR
QFQRRKWPEMKRPSSKSLGQLWEGWEG

Important features:**Signal peptide:**

amino acids 1-17

Sulfatases signature 1:

amino acids 86-99

Homologous region to sulfatase:

amino acids 87-106, 133-146, 216-229, 291-320, 365-375

N-glycosylation sites:amino acids 65-69, 112-116, 132-136, 149-153, 171-175,
198-202, 241-245, 561-565, 608-612, 717-721, 754-758,
764-768

77/392

FIGURE 75

CCCACGCGTCCGCCCACGCGTCCGGTGGACTATGGGCCAGTTTTTGTGCAAGAACCAGAT
GATATTATTTTTTCCAAGTCTGATGAAAAGAAGGTAGCATTGAATTGTGAAGTTCGT
GGCAATCCAGTTCACAGATGGCTTCGAAATGGAACAGAAATAGATCTGGAAAGT
GATTATCGCTACAGTTTGATAGATGGCACCTTCATTATAAGCAATCCAAGTGAAGCAAAG
GATTCTGGTCATTATCAGTGTTTAGCAACCAACACTGTGGGGAGTATTCTTAGTAGAGAA
GCTACACTGCAGTTTGCCTATCTGGGAAATTTTAGTGGCCGACAAGAAGTGCAGTCTCT
GTGAGGGAAGGCCAGGGTGTCTGATGCTCTCCTCCGCCACATTCACCAGAGATC
ATCTATAGCTGGGTATTTAATGAGTTCCTTCCTTTGTGGCGGAAGACAGCCGCGGTTT
ATCTCCCAGGAGACAGGCAACCTTTATATTTCTAAAGTCAAACATCAGATGTTGGCAGC
TATATTTGTCTGGTGAAAAACACAGTGACGAATGCTAGAGTCCTTAGTCCTCCAACGCCA
CTCACTCTGCGTAATGATGGTGTGATGGGAGAATATGAGCCGAAAATTGAGGTCCATTTT
CCTTTACGGTTACAGCTGCTAAAGGAACAACCTGTTAAGATGGAATGCTTTGCACTTGGC
AACCCCGTTCCAACAATCACATGGATGAAGGTTAATGGTTATATTCTAGTAAGGCACGT
CTGCGGAAATCTCAGGCGGTGCTGGAAATACCGAATGTACAGCTGGATGATGCAGGCATT
TATGAGTGCAGAGCTGAAAACCTCACGTGGAAAAAATTCCTTTCGTGGACAATTACAAGTA
TACACCTACCCACACTGGGTAGAAAAACTGAATGATACTCAGTTAGACAGTGGGAGCCCT
CTCCGATGGGAATGTAAGGCTACTGGAAAACCCAGACCCACGTATCGTTGGCTGAAGAAT
GGAGTACCCCTCTCACCTCAGAGTAGGGTTGAGATGGTTAATGGAGTATTGATGATCCAC
AATGTGAATCAATCAGATGCTGGAATGTATCAGTGTTTGGCTGAAAATAAGTATGGAGCC
ATTTACGCTAGTGCTGAGCTGAAGATTCTAGCTTCAGCTCCCACTTTTGCAGTGAATCAA
CTGAAGAAAACAATAATTGTTACCAAAGACCAAGAAGTTGTCATAGAGTGCAAACCCCAA
GGCTCTCCAAAACCAACCATCTCTTGGAAGAAAGGAGACAGAGCAGTTAGAGAAAACAAA
AGAATAGCTATTCTTCCAGACGGGAGTCTACGGATCCTAAATGCTTCCAAATCAGACGAG
GGAAAGTACGTTTGCCGAGGGGAAAACGTCTTTGGTTCTGCTGAAAT

78/392

FIGURE 76

MCSPPPHSPEIIYSWVFNEFPSFVAEDSRRFISQETGNLYISKVQTSADVGSYICLVKNTV
TNARVLSPTPLTLRNDGVMGEYEPKIEVHFPTVTAAGTTVKMECFALGNPVPTITWM
KVNGYIPSKARLRKSQAVLEIPNVQLDDAGIYECRAENSRGKNSFRGQLQVYTYPHWVEK
LNDTQLDSGSPLRWECKATGKPRPTYRWLKNGVPLSPQSRVEMVNGVLMIHNVNQSDAGM
YQCLAENKYGAIIYASAELKILASAPTFALNQLKKTIIIVTKDQEVVIECKPQGSPKPTISW
KKGDRAVRENKRIAILPDGSLRILNASKSDEGKYVCRGENVFGSAE

Signal sequence:

None

Transmembrane domain:

None

N-glycosylation site:

182-185, 234-237, 325-328

Tyrosine kinase phosphorylation site:

328-334

N-myristoylation site:

50-55, 150-155, 239-244, 250-255

Immunoglobulin domain:

2-56, 100-156, 189-245, 281-338

79/392

FIGURE 77

GCTCCCAGCCAAGAACCTCGGGGCCGCTGCGCGGTGGGGAGGAGTTCCTCCGAAACCCGGC
CGCTAAGCGAGGCCTCCTCCTCCCGCAGATCCGAACGGCCTGGGCGGGGTACCCCGGCT
GGGACAAGAAGCCGCCGCTGCCTGCCCGGGCCCGGGAGGGGGCTGGGGCTGGGGCCGG
AGGCGGGGTGTGAGTGGGTGTGTGCGGGGGGCGGAGGCTTGATGCAATCCCGATAAGAAA
TGCTCGGGTGTCTTGGGCACCTACCCGTGGGGCCCGTAAGGCGCTACTATATAAGGCTGC
CGGCCCGGAGCCGCCGCGCCGTGAGAGCAGGAGCGCTGCGTCCAGGATCTAGGGCCACGA
CCATCCCAACCCGGCACTCACAGCCCCGAGCGCATCCCGGTGCGCCGCCAGCCTCCCGC
ACCCCATCGCCGGAGCTGCGCCGAGAGCCCCAGGGAGGTGCCCATGCGGAGCGGGTGTGT
GGTGGTCCACGTATGGATCCTGGCCGGCCTCTGGCTGGCCGTGGCCGGGCGCCCCCTCGC
CTTCTCGGACGCGGGGCCCCACGTGCACTACGGCTGGGGCGACCCCATCCGCCTGCGGCA
CCTGTACACCTCCGGCCCCCACGGGCTCTCCAGCTGCTTCCTGCGCATCCGTGCCGACGG
CGTCGTGGACTGCGCGCGGGGCCAGAGCGCGCACAGTTTGCTGGAGATCAAGGCAGTCGC
TCTGCGGACCGTGGCCATCAAGGGCGTGACAGCGTGCGGTACCTCTGCATGGGCGCCGA
CGGCAAGATGCAGGGGCTGCTTCAGTACTCGGAGGAAGACTGTGCTTTCGAGGAGGAGAT
CCGCCCAGATGGCTACAATGTGTACCGATCCGAGAAGCACCGCCTCCCGGTCTCCCTGAG
CAGTGCCAAACAGCGGCAGCTGTACAAGAACAGAGGCTTTCTTCCACTCTCTCATTTCT
GCCCATGCTGCCATGGTCCCAGAGGAGCCTGAGGACCTCAGGGGCCACTTGGAATCTGA
CATGTTCTCTTCGCCCCCTGGAGACCGACAGCATGGACCCATTTGGGCTTGTCACCGGACT
GGAGGCCGTGAGGAGTCCAGCTTTGAGAAGTAACTGAGACCATGCCCGGGCCTCTTCAC
TGCTGCCAGGGGCTGTGGTACCTGCAGCGTGGGGGACGTGCTTCTACAAGAACAGTCCTG
AGTCCACGTTCTGTTTAGCTTTAGGAAGAAACATCTAGAAGTTGTACATATTCAGAGTTT
TCCATTGGCAGTGCCAGTTTCTAGCCAATAGACTTGTCTGATCATAACATTGTAAGCCTG
TAGCTTGCCCAGCTGCTGCCTGGGCCCCCATTCTGCTCCCTCGAGGTTGCTGGACAAGCT
GCTGCACTGTCTCAGTTCTGCTTGAATACCTCCATCGATGGGGAACCTCACTTCCTTTGGA
AAAATTCTTATGTCAAGCTGAAATTCTCTAATTTTTTCTCATCACTTCCCCAGGAGCAGC
CAGAAGACAGGCAGTAGTTTTAATTTTCAAGAACAGGTGATCCACTCTGTAAACAGCAGG
TAAATTTCACTCAACCCCATGTGGGAATTGATCTATATCTCTACTTCCAGGGACCATTG
CCCTTCCCAAATCCCTCCAGGCCAGAACTGACTGGAGCAGGCATGGCCCACCAGGCTTCA
GGAGTAGGGGAAGCCTGGAGCCCCACTCCAGCCCTGGGACAACCTTGAGAATTTCCCTGA
GGCCAGTTCTGTATGGATGCTGTCTTGAGAATAACTTGCTGTCCCGGTGTCACCTGCTT
CCATCTCCAGCCACCAGCCCTCTGCCACCTCACATGCCCTCCCCATGGATTGGGGCCT
CCCAGGCCCCCACCTTATGTCAACCTGCACCTTCTTGTTCAAAAATCAGGAAAAGAAAAG
ATTTGAAGACCCCAAGTCTTGTCAATAACTTGCTGTGTGGAAGCAGCGGGGGAAGACCTA
GAACCCTTTCCCAGCACTTGTTTTTCCAACATGATATTTATGAGTAATTTATTTTGATA
TGTACATCTCTATTTTCTTACATTATTTATGCCCCCAAATTATATTTATGTATGTAAGT
GAGGTTTGTTTTGTATATTAAAATGGAGTTTGTTTGT

80/392

FIGURE 78

MRSGCVVHVWILAGLWLAVAGRPLAFSDAGPHVHYGWGDPIRLRHLYTSGPHGLSSCFL
RIRADGVVDCARGQSAHSLEIKAVLRRTVAIKGVHVRVRYLCMGADGKMQGLLQYSEEDC
AFEEEIRPDGYNVYRSEKHRLPVSLSSAKQRQLYKNRGFLPLSHFLPMLPMVPEEPEDLR
GHLESDMFSSPLETDSMDPFGLVTGLEAVRSPSF EK

Signal peptide:
amino acids 1-22

Casein kinase II phosphorylation site:
amino acids 78-82, 116-120, 190-194, 204-208

N-myristoylation site:
amino acids 15-21, 54-60, 66-72, 201-207

Prokaryotic membrane lipoprotein lipid attachment site:
amino acids 48-59

81/392

FIGURE 79

CGGACGCGTGGGCGGACGCGTGGGCCTGGGCAAGGGCCGGGGCGCCGGGCGGAGCCACCTCTTCCC
CTCCCCCGCTTCCCTGTCGCGCTCCGCTGGCTGGACGCGCTGGAGGAGTGGAGCAGCACCCGGCCG
GCCCTGGGGGCTGACAGTCGGCAAAGTTTGGCCCCAAGAGGAAGTGGTCTCAAACCCCGGCAGGTG
GCGACCAGGCCAGACCAGGGGCGCTCGCTGCCTGCGGGCGGGCTGTAGGCGAGGGCGCGCCCCAGT
GCCGAGACCCGGGGCTTCAGGAGCCGGCCCCCGGGAGAGAAGAGTGC GGCGGGCAGGAGAAAACA
ACTCCAAAGTTGGCGAAAGGCACCGCCCCCTACTCCCGGGCTGCCGCGCCTCCCCGCCCCAGCCC
TGGCATCCAGAGTACGGGTCGAGCCCGGGCCATGGAGCCCCCTGGGGAGGCGGCACACAGGGAGCC
TGGGCGCCCCGGGGCTCCGCGCGACCCCATCGGGTAGACACAGAAAGCTCCGGGACCTTCCGGCA
CCTCTGGACAGCCAGGATGCTGTTGGCCACCCTCCTCCTCCTCCTTGGAGGCGCTCTGGCCC
ATCCAGACCGGATTATTTTCCAAATCATGCTTGTGAGGACCCCCCAGCAGTGCTCTTAGAAGTGC
AGGGCACCTTACAGAGGCCCTGGTCCGGGACAGCCGCACCTCCCCTGCCAACTGCACCTGGCTCA
TCCTGGGCGAGCAAGGAACAGACTGTACCATCAGGTTCCAGAAGCTACACCTGGCCTGTGGCTCAG
AGCGCTTAACCCCTACGCTCCCCTCTCCAGCCACTGATCTCCTGTGTGAGGCACCTCCAGCCCTC
TGCAGCTGCCCGGGGGCAACGTACCATCACTTACAGCTATGCTGGGGCCAGAGCACCCATGGGCC
AGGGCTTCCTGCTCTCCTACAGCCAAGATTGGCTGATGTGCCTGCAGGAAGAGTTTCAGTGCCCTGA
ACCACCGCTGTGTATCTGCTGTCCAGCGCTGTGATGGGGTTGATGCCCTGTGGCGATGGCTCTGATG
AAGCAGGTTGCAGCTCAGACCCCTTCCCTGGCCTGACCCCAAGACCCGTCCCCTCCCTGCCTTGCA
ATGTCACCTTGGAGGACTTCTATGGGGTCTTCTCCTCTCCTGGATATACACACCTAGCCTCAGTCT
CCCACCCCCAGTCTTGCCATTGGCTGCTGGACCCCCATGATGGCCGGCGGCTGGCCGTGCGCTTCA
CAGCCCTGGACTTGGGCTTTGGAGATGCAGTGCACTGTGTATGACGGCCCTGGGCCCCCTGAGAGCT
CCCGACTACTGCGTAGTCTCACCCACTTCAGCAATGGCAAGGCTGTCACTGTGGAGACACTGTCTG
GCCAGGCTGTTGTGTCTTACCACACAGTTGCTTGGAGCAATGGTCGTGGCTTCAATGCCACCTACC
ATGTGCGGGGCTATTGCTTGCCTTGGGACAGACCCTGTGGCTTAGGCTCTGGCCTGGGAGCTGGCG
AAGGCCTAGGTGAGCGCTGCTACAGTGAGGCACAGCGCTGTGACGGCTCATGGGACTGTGCTGACG
GCACAGATGAGGAGGACTGCCCAGGCTGCCACCTGGACACTTCCCCTGTGGGGCTGCTGGCACCT
CTGGTGCCACAGCCTGTCTACCTGCCTGCTGACCGCTGCAACTACCAGACTTTCTGTGCTGATGGAG
CAGATGAGAGACGCTGTGCGCATTGCCAGCCTGGCAATTTCCGATGCCGGGACGAGAAGTGCGTGT
ATGAGACGTGGGTGTGCGATGGGCAGCCAGACTGTGCGGACGGCAGTGATGAGTGGGACTGCTCCT
ATGTTCTGCCCCGCAAGGTCATTACAGCTGCAGTCACTGGCAGCCTAGTGTGCGGCCTGCTCCTGG
TCATCGCCCTGGGCTGCACCTGCAAGCTCTATGCCATTGCAACCCAGGAGTACAGCATCTTTGCCC
CCCTCTCCCGGATGGAGGCTGAGATTGTGACAGCAGCAGGCACCCCTTCCCTACGGGCAGCTCATTG
CCCAGGCTGCCATCCACCTGTAGAAGACTTTCCTACAGAGAATCCTAATGATAACTCAGTGCTGG
GCAACCTGCGTTCTCTGCTACAGATCTTACGCCAGGATATGACTCCAGGAGGTGGCCAGGTGCC
GCCGTGCTCAGCGGGGCCGCTTGATGCGACGCCTGGTACGCCGTCTCCGCCGCTGGGGCTTGCTCC
CTCGAACCAACACCCCGGCTCGGGCCTCTGAGGCCAGATCCCAGGTCACACCTTCTGCTGCTCCCC
TTGAGGCCCTAGATGGTGGCACAGGTCCAGCCCGTGAGGGCGGGGCGAGTGGGTGGGCAAGATGGGG
AGCAGGCACCCCCACTGCCCATCAAGGCTCCCCCTCCCATCTGCTAGCACGTCTCCAGCCCCCACTA
CTGTCCCTGAAGCCCCAGGGCCACTGCCCTCACTGCCCTTAGAGCCATCACTATTGTCTGGAGTGG
TGCAGGCCCTGCGAGGCCGCTGTGCCCAGCCTGGGGCCCCCAGGACCAACCCGGAGCCCCCTG
GACCCACACAGCAGTCTTGCCCTGGAAGATGAGGACGATGTGCTACTGGTGCCACTGGCTGAGC
CGGGGTGTGGGTAGCTGAGGCAGAGGATGAGCCACTGCTTACCTGAGGGGACCTGGGGGCTCTAC
TGAGGCCCTCTCCCCTGGGGGCTCTACTCATAGTGGCACAACTTTTAGAGGTGGGTGAGCCTCCCC
TCCACCACTTCCCTTCCCTGTCCCTGGATTTAGGGACTTGGTGGGCCCTCCCGTTGACCCATATGTAG
CTGCTATAAAGTTAAGTGTCCCTCAGGCAGGGAGAGGCTCACAGAGTCTCCTCTGTACGTGGCCA
TGGCCAGACACCCAGTCCCTTACCACCACTGCTCCCCACGCCACCACCATTTGGGTGGCTGTT
TTTAAAAAGTAAAGTTCTTAGAGGATCATAGGTCCTGGACACTCCATCCTTGCCAAACCTCTACCCA
AAAGTGGCCTTAAGCACCGGAATGCCAATTAAGTAGAGACCCCTCCAGCCCCCAAGGGGAGGATTTG
GGCAGAACCTGAGGTTTGGCCATCCACAATCCCTCCTACAGGGCCTGGCTCACAAAAGAGTGCAA
CAATGCTTCTATTCCATAGCTACGGCATTGCTCAGTAAGTTGAGGTCAAAAATAAGGAATCATA
CATCTC

82/392

FIGURE 80

MLLATLLLLLLGGALAHPDRIIFPNHACEDPPAVLLEVQGTLQRPLVRDSRTSPANCTWL
ILGSKEQTVTIRFQKLHLACGSERLTLRSPLQPLISLCEAPPSPLQLPGGNVTITYSYAG
ARAPMGQGFLLSYSQDWLMCLQEEFQCLNHRCVSAVQRCDGVDACGDGSDEAGCSSDPFP
GLTPRPVPSLPCNVTLEDYGVFSSPGYTHLASVSHQPQCHWLLDPHDGRRRLAVRFTALD
LGFGDAVHVYDGPPESSRLRLSLTHFSNGKAVTVETLSGQAVVSYHTVAWSNGRGFNA
TYHVRGYCLPWDRPCGLGSGLGAGEGLGERCYSEAQRCDGSWDCADGTDEEDCPGCPPGH
FPCGAAGTSGATACYLPADRCNYQTFCADGADERRCRHCQPGNFRRCRDEKCVYETWVCDG
QPD CADGSDEWDCSYVLPRKVITA AVIGSLVCGLLLVI ALGCTCKLYAIRTQEYSIFAPL
SRMEAEIVQQQAPPSYGQLIAQGAIPPVEDFPTENPNDNSVLGNLRSLLQILRQDMTPGG
GPGARRRQRGRMLMRRLVRRRLRWGLLPRTNTPARASEARSQVTPSAAPLEALDGGTGPAR
EGGAVGGQDGEQAPPLPIKAPLPSASTSPAPTTVPEAPGPLPSLPLEPSLLSGVVQALRG
RLPSLGP PGPTRSPPGPHTAVLALEDEDDVLLVPLAEPGVWVAEAEDEPLLT

Important features:**Signal peptide:**

amino acids 1-16

Transmembrane domain:

amino acids 442-462

LDL-receptor class A (LDLRA) domain proteins:

amino acids 411-431, 152-171, 331-350 and 374-393

83/392

FIGURE 81

CTTCTGTGCTGTTCTTCTTGCCTCTAACTTGTAACAAGACGTACTAGGACGATGCTAA
TGGAAAGTCACAAACCGCTGGGTTTTTGAAGGATCCTTGGGACCTCATGCACATTTGTG
GAAACTGGATGGAGAGATTTGGGGAAGCATGGACTCTTTAGCCAGCTTAGTTCTCTGTGG
AGTCAGCTTGCTCCTTTCTGGAAGTGTGGAAGGTGCCATGGACTTGATCTTGATCAATTC
CCTACCTCTTGTATCTGATGCTGAAACATCTCTCACCTGCATTGCCTCTGGGTGGCGCCC
CCATGAGCCCATCACCATAGGAAGGGACTTTGAAGCCTTAATGAACCAGCACCAGGATCC
GCTGGAAGTTACTCAAGATGTGACCAGAGAATGGGCTAAAAAAGTTGTTTGGAAGAGAGA
AAAGGCTAGTAAGATCAATGGTGCTTATTTCTGTGAAGGGCGAGTTCGAGGAGAGGCAAT
CAGGATACGAACCATGAAGATGCGTCAACAAGCTTCCTTCCTACCAGCTACTTTAACTAT
GACTGTGGACAAGGGAGATAACGTGAACATATCTTTCAAAAAGGTATTGATTAAAGAAGA
AGATGCAGTGATTTACAAAAATGGTTCCTTCATCCATTGAGTCCCCCGGCATGAAGTACC
TGATATTCTAGAAGTACACCTGCCTCATGCTCAGCCCCAGGATGCTGGAGTGTAAGTCCGGC
CAGGTATATAGGAGGAAACCTCTTCACCTCGGCCTTCACCAGGCTGATAGTCCGGAGATG
TGAAGCCCAGAAAGTGGGGACCTGAATGCAACCATCTCTGTACTGCTTGTATGAACAATGG
TGTCTGCCATGAAGATACTGGAGAATGCATTTGCCCTCCTGGGTTTTATGGGAAGGACGTG
TGAGAAGGCTTGTGAAGTGCACACGTTTGGCAGAACTTGTAAGAAAGGTGCAGTGGACA
AGAGGGATGCAAGTCTTATGTGTTCTGTCTCCCTGACCCCTATGGGTGTTCTGTGCCAC
AGGCTGGAAGGGTCTGCAGTGCAATGAAGCATGCCACCCTGGTTTTTACGGGCCAGATTG
TAAGCTTAGGTGCAGCTGCAACAATGGGGAGATGTGTGATCGCTTCCAAGGATGTCTCTG
CTCTCCAGGATGGCAGGGGCTCCAGTGTGAGAGAGAAGGCATACCGAGGATGACCCCAA
GATAGTGGATTTGCCAGATCATATAGAAGTAAACAGTGGTAAATTTAATCCCATTTGCAA
AGCTTCTGGCTGGCCGCTACCTACTAATGAAGAAATGACCCTGGTGAAGCCGGATGGGAC
AGTGCTCCATCCAAAAGACTTTAACCATACGGATCATTTCTCAGTAGCCATATTCACCAT
CCACCGGATCCTCCCCCTGACTCAGGAGTTTGGGTCTGCAGTGTGAACACAGTGGCTGG
GATGGTGGAAAAGCCCTTCAACATTTCTGTAAAGTTCTTCCAAAGCCCCCTGAATGCCCC
AAACGTGATTGACACTGGACATAACTTTGCTGTCAACATCAGCTCTGAGCCTTACTT
TGGGGATGGACCAATCAAATCCAAGAAGCTTCTATACAAACCCGTTAATCACTATGAGGC
TTGGCAACATATTCAGTGACAAATGAGATTGTTACACTCAACTATTTGGAACCTCGGAC
AGAATATGAACTCTGTGTGCAACTGGTCCGTGCTGGAGAGGGTGGGGAAGGGCATCCTGG
ACCTGTGAGACGCTTCAACACAGCTTCTATCGGACTCCCTCCTCCAAGAGGTCTAAATCT
CCTGCCTAAAAGTCAGACCACTCTAAATTTGACCTGGCAACCAATATTTCCAAGCTCGGA
AGATGACTTTTATGTTGAAGTGGAGAGAAGGTCTGTGCAAAAAGTGATCAGCAGAATAT
TAAAGTTCAGGCAACTTGACTTCGGTGCTACTTAACAACCTTACATCCCAGGGAGCAGTA
CGTGGTCCGAGCTAGAGTCAACACCAAGGCCAGGGGAATGGAGTGAAGATCTCACTGC
TTGGACCCTTAGTGACATTCTTCCTCCTCAACCAGAAAACATCAAGATTTCCAACATTAC
ACACTCCTCGGCTGTGATTTCTTGACAATATTGGATGGCTATTCTATTTCTTCTATTAC
TATCCGTTACAAGGTTCAAGGCAAGAATGAAGACCAGCACGTTGATGTGAAGATAAAGAA
TGCCACCATCATTGAGTATCAGCTCAAGGGCCTAGAGCCTGAAACAGCATACCAGGTGGA
CATTTTTGTCAGAGAACAACATAGGGTCAAGCAACCCAGCCTTTTCTCATGAACTGGTGAC
CCTCCCAGAATCTCAAGCACCAGCGGACCTCGGAGGGGGGAAGATGCTGCTTATAGCCAT
CCTTGGCTCTGCTGGAATGACCTGCCTGACTGTGCTGTTGGCCTTTCTGATCATATTGCA
ATTGAAGAGGGCAAATGTGCAAAGGAGAATGGCCCAAGCCTTCCAAAACGTGAGGGGAAGA
ACCAGCTGTGCAGTTCAACTCAGGGACTCTGGCCCTAAACAGGAAGGTCAAAAACAACCC
AGATCCTACAATTTATCCAGTGCTTGACTGGAATGACATCAAATTTCAAGATGTGATTGG
GGAGGGCAATTTTGGCCAAGTTCTTAAGGCGCGCATCAAGAAGGATGGGTTACGGATGGA
TGCTGCCATCAAAAGAATGAAAGAATATGCCTCAAAGATGATCACAGGGACTTTGCAGG
AGAAGTGAAGTTCTTTGTAAACTTGACACCATCCAAACATCATCAATCTCTTAGGAGC

84/392

ATGTGAACATCGAGGCTACTTGTACCTGGCCATTGAGTACGCGCCCCATGGAAACCTTCT
GGACTTCCTTCGCAAGAGCCGTGTGCTGGAGACGGACCCAGCATTTGCCATTGCCAATAG
CACCGCGTCCACACTGTCTCCCAGCAGCTCCTTCACTTCGCTGCCGACGTGGCCCGGGG
CATGGACTACTTGAGCCAAAAACAGTTTATCCACAGGGATCTGGCTGCCAGAAACATTTT
AGTTGGTGAAAACTATGTGGCAAAAATAGCAGATTTTGGATTGTCCCGAGGTCAAGAGGT
GTACGTGAAAAAGACAATGGGAAGGCTCCAGTGCGCTGGATGGCCATCGAGTCACTGAA
TTACAGTGTGTACACAACCAACAGTGATGTATGGTCCTATGGTGTGTTACTATGGGAGAT
TGTTAGCTTAGGAGGCACACCCTACTGCGGGATGACTTGTGCAGAACTCTACGAGAAGCT
GCCCCAGGGCTACAGACTGGAGAAGCCCCCTGAAGTGTGATGATGAGGTGTATGATCTAAT
GAGACAATGCTGGCGGGAGAAGCCTTATGAGAGGCCATCATTTGCCCAGATATTGGTGTC
CTTAAACAGAATGTTAGAGGAGCGAAAGACCTACGTGAATACCACGCTTTATGAGAAGTT
TACTTATGCAGGAATTGACTGTTCTGCTGAAGAAGCGGCCTAGGACAGAACATCTGTATA
CCCTCTGTTTTCCCTTTCACTGGCATGGGAGACCCTTGACAACTGCTGAGAAAACATGCCT
CTGCCAAAGGATGTGATATATAAGTGTACATATGTGCTGGAATTCTAACAAGTCATAGGT
TAATATTTAAGACACTGAAAAATCTAAGTGATATAAATCAGATTCTTCTCTCTCATTTTA
TCCCTCACCTGTAGCATGCCAGTCCCGTTTCATTTAGTCATGTGACCACTCTGTCTTG TG
TTTCCACAGCCTGCAAGTTCAGTCCAGGATGCTAACATCTAAAAATAGACTTAAATCTCA
TTGCTTACAAGCCTAAGAATCTTTAGAGAAGTATACATAAGTTTAGGATAAAATAATGGG
ATTTTCTTTTCTTTTCTCTGGTAATATTGACTTGTATATTTTAAGAAATAACAGAAAGCC
TGGGTGACATTTGGGAGACATGTGACATTTATATATTGAATTAATATCCCTACATGTATT
GCACATTGTAAAAAGTTTTAGTTTTGATGAGTTGTGAGTTTACCTTGTATACTGTAGGCA
CACTTTGCACTGATATATCATGAGTGAATAAATGTCTTGCCCTACTCAAAAAAAAAAAAA

85/392

FIGURE 82

MDSLASLVLCGVSLLLSGTVEGAMDLILINSLPLVSDAETSLTCIASGWRPHEPITIGRD
FEALMNQHQDPLEVTQDVTREWAKKVWVKREKASKINGAYFCEGRVVRGEAIRIRTMKMRQ
QASFLPATLTMTVDKGDVNISFKKVLKEEDAVIYKNGSFIHSVPRHEVPDILEVHLPH
AQPQDAGVYSARYIGGNLFTSAFTRLIVRRCEAQKWGPECNHLCTACMNGVCHEDTGEC
ICPPGFMGRTCEKACELHTFGRTCKERCSCGQEGCKSYVFCCLPDYPYGCSCATGWKGLQCNE
ACHPGFYGPDCKLRCSNNGEMCDRFQGCCLSPGWQGLQCEREGIPRMTPKIVDLPDHIE
VNSGKFNPICKASGWPLPTNEEMTLVKPDGTVLHPKDFNHTDHFSAIFTIHRILPPDSG
VWVCSVNTVAGMVEKPFNISVKVLPKPLNAPNVIDTGHNFAVINISSEPYFGDGPIKSKK
LLYKPVNHYEAWQHIQVTNEIVTLNYLEPRTEYELCVQLVRRGEGGEGHPGPVRRFTTAS
IGLPPPRGLNLLPKSQTTLNLTWQPIFPSEDDFYVEVERRSVQKSDQQNIKVPGNLTSV
LLNNLHPREQYVVRARVNTKAQGEWSEDLTAWTLSDILPPQPENIKISNITHSSAVISWT
ILDGYSISSITIRYKVQGNEDQHVDVKIKNATIIQYQLKGLEPETAYQVDIFAENNIGS
SNPAFSHELVTLPESQAPADLGGGKMLLIAILGSAGMTCLTVLLAFLIILQLKRANVQRR
MAQAFQNVREEPAVQFNSGTLALNRKVKNPDPTIYPVLDWNDIKFQDVI GEGNFGQVLK
ARIKKDGLRMDAAIKRMKEYASKDDHRDFAGELEVLCKLGHPNIIINLLGACEHRGYLYL
AIEYAPHGNLLDFLRKSRVLETDPAFAIANSTASTLSSQQLHFAADVARGMDYLSQKQF
IHRDLAARNILVGENYVAKIADFGLSRGQEVYVKKTMGRLPVRWMAIESLNYSVYTTNSD
VWSYGVLLWEIVSLGGTPYCGMTCAELYEKLPGYRLEKPLNCDDEVYDLMRQCWREKPY
ERPSFAQILVSLNRMLEERKTYVNTTLYEKFTYAGIDCSAEAEA

Signal sequence:

1-38

Transmembrane domain:

750-770

N-glycosylation site:

140-143, 158-161, 399-402, 438-441, 464-467, 560-563, 596-599, 649-652, 691-694, 930-933, 1011-1014, 1104-1107

cAMP- and cGMP-dependent protein kinase phosphorylation site:

534-537

Tyrosine kinase phosphorylation site:

149-156, 808-816, 1094-1102

N-myristoylation site:

18-23, 98-103, 187-192, 196-201, 270-275, 286-291, 295-300, 420-425, 595-600, 984-989, 1036-1041, 1041-1046, 1115-1120

Prokaryotic membrane lipoprotein lipid attachment site:

882-892

86/392

EGF-like domain cysteine pattern signature:

240-251, 287-298, 329-340

Tyrosine protein kinases specific active-site signature:

960-972

Protein kinase domain:

824-1092

Fibronectin type III domain:

444-529, 543-626, 639-724

EGF-like domain:

220-251, 268-298

laminin_EGF Laminin EGF-like (Domains III and V):

219-268

Immunoglobulin domain:

156-193

Zinc finger:

295-313

Receptor tyrosine kinase:

844-868, 869-898, 936-982, 986-1024, 1025-1052, 1052-1088

87/392

FIGURE 83

CGCGCCGGGCGCAGGGAGCTGAGTGGACGGCTCGAGACGGCGGCGCGTGCAGCAGCTCCA
GAAAGCAGCGAGTTGGCAGAGCAGGGCTGCATTTCCAGCAGGAGCTGCGAGCACAGTGCT
GGCTCACAACAAGATGCTCAAGGTGTCAGCCGTACTGTGTGTGTGTGCAGCCGCTTGGTG
CAGTCAGTCTCTCGCAGCTGCCCGGGCGGTGGCTGCAGCCGGGGGGCGGTCCGACGGCGG
TAATTTTCTGGATGATAAACAATGGCTCACCACAATCTCTCAGTATGACAAGGAAGTCCG
ACAGTGGAACAAATTCGAGACGAAGTAGAGGATGATTATTTCCGCACTTGGAGTCCAGG
AAAACCCCTTCGATCAGGCTTTAGATCCAGCTAAGGATCCATGCTTAAAGATGAAATGTAG
TCGCCATAAAGTATGCATTGCTCAAGATTCTCAGACTGCAGTCTGCATTAGTCACCGGAG
GCTTACACACAGGATGAAAGAAGCAGGAGTAGACCATAGGCAGTGGAGGGGTCCCATATT
ATCCACCTGCAAGCAGTGCCCGAGTGGTCTATCCCAGCCCTGTTTGTGGTTTCAGATGGTCA
TACCTACTCTTTTCAGTGCAAAC TAGAATATCAGGCATGTGTCTTAGGAAAACAGATCTC
AGTCAAATGTGAAGGACATTGCCCATGTCTTCAGATAAGCCCACCAGTACAAGCAGAAA
TGTTAAGAGAGCATGCAGTGACCTGGAGTTCAGGGAAGTGGCAAACAGATTGCGGGACTG
GTTCAAGGCCCTTCATGAAAGTGGAAGTCAAAACAAGAAGACAAAAACATTGCTGAGGCC
TGAGAGAAGCAGATTCGATACCAGCATCTTGCCAATTTGCAAGGACTCACTTGGCTGGAT
GTTTAAACAGACTTGATACAAACTATGACCTGCTATTGGACCAGTCAGAGCTCAGAAGCAT
TTACCTTGATAAGAATGAACAGTGTACCAAGGCATTCTTCAATTCTTGTGACACATACAA
GGACAGTTTAAATATCTAATAATGAGTGGTGCTACTGCTTCCAGAGACAGCAAGACCCACC
TTGCCAGACTGAGCTCAGCAATATTCAGAAGCGGCAAGGGGTAAAGAAGCTCCTAGGACA
GTATATCCCCCTGTGTGATGAAGATGGTTACTACAAGCCAACACAATGTCATGGCAGTGT
TGGACAGTGCTGGTGTGTTGACAGATATGGAAATGAAGTCATGGGATCCAGAATAAATGG
TGTTGCAGATTGTGCTATAGATTTTGAGATCTCCGGAGATTTTGCTAGTGGCGATTTTCA
TGAATGGACTGATGATGAGGATGATGAAGACGATATTATGAATGATGAAGATGAAATTGA
AGATGATGATGAAGATGAAGGGGATGATGATGATGGTGGTGGTATGACCATGATGTATACAT
TTGATTGATGACAGTTGAAATCAATAAATTCTACATTTCTAATATTTACAAAAATGATAG
CCTATTTTAAATTTATCTTCTTCCCCAATAACAAAATGATTCTAAACCTCACATATATTTT
GTATAATTATTTGAAAAATTGCAGCTAAAGTTATAGAACTTTATGTTTAAATAAGAATCA
TTTGCTTTGAGTTTTTTATATTCCTTACACAAAAGAAAATACATATGCAGTCTAGTCAGA
CAAAATAAAGTTTTGAAGTGCTACTATAATAAATTTTTTACGAGAACAAACTTTGTAAAT
CTTCATAAGCAAAATGACAGCTAGTGCTTGGGATCGTACATGTTAATTTTTTGAAAGAT
AATTCTAAGTGAAATTTAAATAAATAAATTTTAAATGACCTGGGTCTTAAGGATTTAGG
AAAAATATGCATGCTTTAATTGCATTTCCAAAGTAGCATCTTGCTAGACCTAGATGAGTC
AGGATAACAGAGAGATACCACATGACTCCAAAAAAAAAAAAAAAAA

88/392

FIGURE 84

MLKVSAVLCVCAAAWCSQSLAAAAVAAGGRSDGGNFLDDKQWLTTISQYDKEVGQWNK
FRDEVEDDYFRTWSPGKPFQALDPAKDPCLKMKCSRHKVCIAQDSQTAVCISHRRLTHR
MKEAGVDHRQWRGPILSTCKQCPVVYPSPVCGSDGHTYSFQCKLEYQACVLGKQISVKCE
GHCPGPSDKPTSTSRNVKRACSDLEFREVANRLRDWFKALHESGSQNKKTLLRPERSR
FDTSILPICKDSLGMFMNRLDTNYDLLDQSELRSIYLDKNEQCTKAFFNSCDTYKDSL
SNNEWCYCFQRQQDPPCQTELSNIQKRQGVKKLLGQYIPLCDEDGYKPTQCHGSVGQCW
CVDRYGNEVMGSRINGVADCAIDFEISGDFASGDFHEWTDDEDDDDIMNDEDEIEDDDE
DEGDDDDGGDDHDVYI

Important features:

Signal peptide:

amino acids 1-16

Leucine zipper pattern:

amino acids 246-267

N-myristoylation sites:

amino acids 357-362, 371-376 and 376-381

Thyroglobulin type-1 repeat proteins:

amino acids 353-365 and 339-352

89/392

FIGURE 85

CCCACGCGTCCGGCACTGCAGTCTCCAGCCTGAGCCATGGGCCGCCGAGCCCTCCTGCTC
CTGCTTCTGTCTTTTCTGGCGCCCTGGGCCACCATAGCCCTCCGGCCGGCCTTAAGGGCC
CTCGGCAGCCTACACTTGCCAACCAACCCACATCCCTCCCGGCTGTAGCCAAGAACTAT
TCGGTTCCTACTTCCAACAGAAGGTTGATCATTTTGGATTTAATACTGTGAAAACTTTT
AATCAGCGGTACCTAGTAGCTGATAAATACTGGAAGAAAAATGGTGGATCAATACTTTTC
TACACTGGTAATGAAGGGGACATTATCTGGTTTTGTAATAACACGGGGTTCATGTGGGAT
GTGGCTGAGGAAGTGAAGCTATGTTGGTGTGTTGCTGAACATCGATACTATGGAGAGTCT
CTCCCCCTTTGGTGACAACCTCATTCAAGGATTCCAGACACTTGAATTTCTGACATCAGAA
CAAGCTCTGGCTGATTTTGCAGAGTTAATCAAACACTTGAAAAGAACAATCCCAGGAGCT
GAAAATCAACCTGTCAATTGCCATAGGAGGCTCCTATGGTGGCATGCTTGCCGCCTGGTTT
AGGATGAAATATCCTCATATGGTAGTTGGAGCTCTTGCACTTCTGCCCCCTATCTGGCAG
TTTGAGGATTTAGTACCTTGTGGTGTATTTATGAAGATCGTAACTACAGATTTTAGGAAA
AGCGGTCCACATTGTTTCAGAGAGCATCCACAGGTCTGGGATGCCATTAATCGACTCTCA
AATACTGGCAGTGGTTTGCAGTGGCTTACTGGAGCCCTTCACTTATGCAGCCCATTAACT
TCTCAGGACATCCAACATTTGAAAGACTGGATCTCTGAAACCTGGGTGAATCTGGCAATG
GTGGACTATCCTTATGCCTCTAACTTTTTTACAGCCTTGCCTGCTTGGCCTATCAAGGTA
GTGTGCCAGTATTTGAAAAATCCCAATGTATCTGATTCACTGCTGCTGCAGAATATTTTC
CAAGCTCTGAATGTATATTACAATTATTCGGGCCAGGTGAAATGCCTGAATATTTTCAGAG
ACAGCAACTAGCAGTCTGGGAACACTGGGTGGAGCTATCAGGCCTGCACAGAAGTAGTC
ATGCCCTTTTGTACTAATGGTGTGATGACATGTTTGAACCTCACTCATGGAACCTTAAAG
GAACTTTCTGATGACTGTTTTCAACAGTGGGGTGTGAGACCAAGGCCCTCCTGGATCACT
ACTATGTATGGAGGCAAAAACATTAGTTTACACACAAAACATTGTTTTTCAGCAATGGTGAA
CTAGACCCCTGGTCAGGAGGTGGAGTAACTAAGGATATCACAGACACTCTGGTTGCAGTC
ACCATCTCAGAGGGGGCCCACTTAGATCTCCGCACCAAGAATGCCTTGGATCCTATG
TCTGTGCTGTTAGCCCGCTCCTTGGAAGTTAGACATATGAAGAATTGGATCAGAGATTTTC
TATGACAGTGCGGGAAAGCAGCACTGAGAACTTTTGATTGTTTTCAATTTCTTCTTTTA
TGTTTACACCACCACATTCCCATTCACCTTGATTTTCTACATGTAATTACCTTCTTTTGT
TTATCATTAGATTTGATGGGGCCAAAGTTGAGATAGAATAGAGGGTGATGACGGTAAGAG
CAAGTGTCCCATGAATGTGATTTCTGGGTCTCACTGTCTTTGCACCACGTCTAGGAA
GAATCTTCTTGATAGCTCTCCACACCATCAGTGGCCCTCATAACTGGAGTAGAGTTCTT
GGTTGCTTTTCATAAGAGGGAGAGTTACTTTC

90/392

FIGURE 86

MGRRALLLLLLSFLAPWATIALRPALRALGSLHLPTNPTSLPAVAKNYSVLYFQQKVDHF
GFNTVKTFFNQRYLVADKYWKKNNGGSILFYTGNEGDIWFCNNTGFMWDVAEELKAMLVFA
EHRYYGESLPFGDNSFKDSRHLNFLTSEQALADFAELIKHLKRTIPGAENQPVIAIGGSY
GGMLAAWFRMKYPHVVVGALAASAPIWQFEDLVPCGVFMKIVTTDFRKSGPHCSESIHRS
WDAINRLSNTGSGLOWLTGALHLCSPLTSQDIQHLKDWISETWVNLAMVDYPYASNFLQP
LPAWPIKVVCQYLKNPNVSDSLLLQNIHQALNVYYNYSQGVKCLNISETATSSLGTLGWS
YQACTEVVMPFCTNGVDDMFEPHSWNLKELSDDCFQQWGVPRPSWITTMYGKKNISSHT
NIVFSNGELDPWSGGGVTKDITDTLVAVTISEGAHHLDLRTKNALDPMSVLLARSLEVRH
MKNWIRDFYDSAGKQH

Signal sequence:

1-18

Transmembrane domain:

None

N-glycosylation site:

47-50, 101-104, 317-320, 336-339, 345-348, 415-418

Glycosaminoglycan attachment site:

433-436

N-myristoylation site:

178-183, 181-186, 182-187, 198-203, 339-344, 434-439

Amidation site:

1-4

alpha/beta hydrolase fold:

115-372

91/392

FIGURE 87

GGCGGCGTCCGTGAGGGGCTCCTTTGGGCAGGGGTAGTGTTTGGTGTCCTGTCTTGCGT
GATATTGACAACTGAAGCTTTCCTGCACCACTGGACTTAAGGAAGAGTGTA CTCTCGTAGG
CGGACAGCTTTAGTGGCCGGCCGGCCGCTCTCATCCCCGTAAGGAGCAGAGTCCTTTGT
ACTGACCAAGATGAGCAACATCTACATCCAGGAGCCTCCACGAATGGGAAGGTTTTATT
GAAACTACAGCTGGAGATATTGACATAGAGTTGTGGTCCAAAGAAGCTCCTAAAGCTTG
CAGAAATTTTATCCAACCTTTGTTTGGAAGCTTATTATGACAATACCATTTTTTCATAGAGT
TGTGCCTGGTTTTCATAGTCCAAGCGGAGATCCTACTGGCACAGGGAGTGGTGGAGAGTC
TATCTATGGAGCGCCATTCAAAGATGAATTTTCATTACCGTTGCGTTTTAATCGGAGAGG
ACTGGTTGCCATGGCAAATGCTGGTTCTCATGATAATGGCAGCCAGTTTTTCTTCACACT
GGGTGAGCAGATGAACTTAACAATAAGCATAACCATCTTTGGAAGGTTACAGGGGATAC
AGTATATAACATGTTGCGACTGTCAGAAGTAGACATTGATGATGACGAAAGACCACATAA
TCCACACAAAATAAAAAGCTGTGAGGTTTTGTTTAATCCTTTTGATGACATCATTTCCAAG
GGAAATTAAAAGGCTGAAAAAGAGAAACCAGAGGAGGAAGTAAAGAAATTGAAACCCAA
AGGCACAAAAAATTTAGTTTACTTTTCAATTTGGAGAGGAAGCTGAGGAAGAAGAGGAGGA
AGTAAATCGAGTTAGTCAGAGCATGAAGGGCAAAAGCAAAAGTAGTCATGACTTGCTTAA
GGATGATCCACATCTCAGTTCTGTTCCAGTTGTAGAAAGTGAAAAAGGTGATGCACCAGA
TTTAGTTGATGATGGAGAAGATGAAAGTGCAGAGCATGATGAATATATTGATGGTGATGA
AAAGAACCTGATGAGAGAAAGAATTGCCAAAAAATTAAAAAAGGACACAAGTGCGAATGT
TAAATCAGCTGGAGAAGGAGAAGTGGAGAAGAAATCAGTCAGCCGCAGTGAAGAGCTCAG
AAAAGAAGCAAGACAATTAAAACGGGAACTCTTAGCAGCAAAACAAAAAAAAGTAGAAAA
TGCAGCAAAACAAGCAGAAAAAAGAAGTGAAGAGGAAGAAGCCCCTCCAGATGGTGCTGT
TGCCGAATACAGAAGAGAAAAAGCAAAAGTATGAAGCTTTGAGGAAGCAACAGTCAAAGAA
GGGAACTTCCCGGGAAGATCAGACCCTTGCACTGCTGAACCAGTTTAAATCTAAACTCAC
TCAAGCAATTGCTGAAACACCTGAAAATGACATTCTTGAAACAGAAGTAGAAGATGATGA
AGGATGGATGTCACATGTACTTTCAGTTTGGAGGATAAAAGCAGAAAAGTGAAAGATGCAAG
CATGCAAGACTCAGATACATTTGAAATCTATGATCCTCGGAATCCAGTGAATAAAAGAAG
GAGGGAAGAAAGCAAAAAGCTGATGAGAGAGAAAAAAGAAAGAAGATATAAATGAGAATAA
TGATAACCAGAACTTGCTGGAAATGTGCCTACAATGGCCTTGTAACAGCCATTGTTCCCA
ACAGCATCACTTAGGGGTGTGAAAAGAAGTATTTTTGAACCTGTTGTCTGGTTTTTGAAAA
ACAATTATCTTGTTTTGCAAATTGTGGAATGATGTAAGCAAATGCTTTTGGTTACTGGTA
CATGTGTTTTTTCCTAGCTGACCTTTTATATTGCTAAATCTGAAATAAAATAACTTTCTCT
TCCACAAAAA

92/392

FIGURE 88

MSNIYIQEPPTNGKVLLKTTAGDIDIELWSKEAPKACRNFIQLCLEAYYDNTIFHRVVP
FIVQGGDPTGTGSGGESIYGAPFKDEFHSRLRFNRRGLVAMANAGSHDNGSQFFFTLGRA
DELNNKHTIFGKVTGDTVYNMLRLSEVDIDDDERPHNPHKIKSCEVLFPFDDIIPREIK
RLKKEKPEEEVKKLKPKGTKNFSLLSFGEEAEVEVNRVSQSMKGKSKSSHDLKDDP
HLSSVPVVESEKGDAPDLVDDGEDESAEHDEYIDGDEKNLMRERIAKKLKKDTSANVKSA
GEGEVEKKSFSRSEELRKEARQLKRELLAAKQKKVENAAKQAEKRSEEEAAPPDGA
VAEYRREKQKYEALRKQQSKKGTSTREDQTLALLNQFKSKLTQAIAETPENDIPETE
VEDDEGWM SHVLQFEDKSRKVKDASMQSDTFEIIDPRNPVNKRREESKLMREKKERR

Important features:**Signal peptide:**

amino acids 1-21

N-glycosylation sites:

amino acids 109-112 and 201-204

Cyclophilin-type peptidyl-prolyl cis-trans isomerase**signature:**

amino acids 49-66

Homologous region to Cyclophilin-type peptidyl-prolyl cis-trans isomerase:

amino acids 96-140, 49-89 and 22-51

93/392

FIGURE 89

CCCGGCTCCGCTCCCTCTGCCCCCTCGGGGTCGCGCGCCACGATGCTGCAGGGCCCTGG
CTCGCTGCTGCTGCTCTTCCTCGCCTCGCACTGCTGCCTGGGCTCGGCGCGCGGGCTCTT
CCTCTTTGGCCAGCCGACTTCTCCTACAAGCGCAGCAATTGCAAGCCCATCCCGGTCAA
CCTGCAGCTGTGCCACGGCATCGAATACCAGAACATGCGGCTGCCCAACCTGCTGGGCCA
CGAGACCATGAAGGAGGTGCTGGAGCAGGCCGGCGCTTGGATCCCGCTGGTCATGAAGCA
GTGCCACCCGGACACCAAGAAGTTCCTGTGCTCGCTCTTCGCCCCCGTCTGCCTCGATGA
CCTAGACGAGACCATCCAGCCATGCCACTCGCTCTGCGTGCAGGTGAAGGACCGCTGCGC
CCCGGTCATGTCCGCCTTCGGCTTCCCCTGGCCCCGACATGCTTGAGTGCGACCGTTTCCC
CCAGGACAACGACCTTTGCATCCCCCTCGCTAGCAGCGACCACCTCCTGCCAGCCACCGA
GGAAGCTCCAAAGGTATGTGAAGCCTGCAAAAATAAAAATGATGATGACAACGACATAAT
GGAAACGCTTTGTAAAAATGATTTTGCCTGAAAATAAAAGTGAAGGAGATAACCTACAT
CAACCGAGATACCAAAATCATCCTGGAGACCAAGAGCAAGACCATTTACAAGCTGAACGG
TGTGTCCGAAAGGGACCTGAAGAAATCGGTGCTGTGGCTCAAAGACAGCTTGCAGTGCAC
CTGTGAGGAGATGAACGACATCAACGCGCCCTATCTGGTCATGGGACAGAAACAGGGTGG
GGAGCTGGTGATCACCTCGGTGAAGCGGTGGCAGAAGGGGCAGAGAGAGTTCAAGCGCAT
CTCCCGCAGCATCCGCAAGCTGCAGTGCTAGTCCCGGCATCCTGATGGCTCCGACAGGCC
TGCTCCAGAGCACGGCTGACCATTTCTGCTCCGGGATCTCAGCTCCCGTTCCCCAAGCAC
ACTCCTAGCTGCTCCAGTCTCAGCCTGGGCAGCTTCCCCCTGCCTTTTGCACGTTTGCAT
CCCCAGCATTTCTGAGTTATAAGGCCACAGGAGTGGATAGCTGTTTTACCTAAAGGAA
AAGCCCACCCGAATCTTGTAGAAATATTCAAATAATAAAATCATGAATATTTTAA

94/392

FIGURE 90

MLQGPGSLLLLFLASHCCLGSARGLFLFGQPDFS YKRSNCKPIPVNLQLCHGIEYQNMRL
PNLLGHETMKEVLEQAGAWIPLVMKQCHPDTKKFLCSLFAPVCLDDLDETIQPCHS LCVQ
VKDRCAPVMSAFGFPWPDMLECDRFPQDNDLCIPLASSDHLLPATEEAPKVCEACKNKND
DDNDIMETLCKNDFALKIKVKEITYINRDTKIILETKSKTIYKLNGVSE RDLKKSVLWLK
DSLQCTCEEMNDINAPYLVMGQKQG GELVITSVKRWQKGQREFKRISRSIRKLQC

Important features:

Signal peptide:

amino acids 1-20

Cysteine rich domain, homologous to frizzled N terminus:

amino acids 6-153

95/392

FIGURE 91

[illegible]

97/392

FIGURE 92

MT PQSLLQTTLFLLSLLFLVQGAHGRGHREDFRFCSQRNQTHRSSLHYKPTPDLRISIEN
SEEALTVHAPFPAAHPASRSFPDPRGLYHFCLYWNRHAGRLHLLYGKRD FLLSDKASSLL
CFQHQEESLAQGPPLLATSVTSWWSPQNISLPSAASFTFSFHSPHTAAHNASVDMCELK
RDLQLLSQFLKHPQKASRRPSAAPASQQLQSLESKLT SVRFMGDMVSFEEDRINATVWKL
QPTAGLQDLHIHSRQEEEQSEIMEYSVLLPRTL FQRTKGRSGEAEKRLLLVD FSSQALFQ
DKNSSQVLGEKVLGIVVQNTKVANLTEPVVLT FQHQLQPKNVTLQCVFWVEDPTLSSPGH
WSSAGCETVRRETQTSCFCNHLTYFAVLMVSSVEVD AVHKHYLSLLSYVGCVVVSALACL V
TIAAYLCSRVP LPCR KPRDYTIKVHMNLLLA VFLLDTSFLLSEPVALTGSEAGCRASAI
FLHFSLLTCLSWMGLEGYNLYRLVVEVFGTYVPGYLLKLSAMGWGFPIFLVTLVALVDVD
NYGPIILAVHRTPEGVIYPSMCWIRDSLVS YITNLGLFSLVFLFNMA MLATMVVQILRLR
PHTQKWSHVLTLGLSLVLGLPWALIFFS FASGTFQLVVL YLFSIITSFQGF LIFIWYWS
MRLQARGGPSPLKSN SDSARLPISSGSTSSRI

Important features:**Signal peptide:**

amino acids 1-25

Putative transmembrane domains:amino acids 382-398, 402-420, 445-468, 473-491, 519-537,
568-590 and 634-657**Microbodies C-terminal targeting signal:**

amino acids 691-693

cAMP- and cGMP-dependent protein kinase phosphorylation sites:

amino acids 198-201 and 370-373

N-glycosylation sites:amino acids 39-42, 148-151, 171-174, 234-237, 303-306, 324-
327 and 341-344**G-protein coupled receptors family 2 proteins:**

amino acids 475-504

98/392

FIGURE 93

CCCACGCGTCCGAAGGCAGACAAAGGTTCAATTTGTAAAGAAGCTCCTTCCAGCACCTCCT
CTCTTCTCCTTTTGCCCAAACCTCACCCAGTGAGTGTGAGCATTTAAGAAGCATCCTCTGC
CAAGACCAAAAGGAAAGAAGAAAAAGGGCCAAAAGCCAAAATGAAACTGATGGTACTTGT
TTTCACCATTGGGCTAACTTTGCTGCTAGGAGTTCAAGCCATGCCTGCAAATCGCCTCTC
TTGCTACAGAAAGATACTAAAAGATCACAACCTGTCAACCTTCCGGAAGGAGTAGCTGA
CCTGACACAGATTGATGTCAATGTCCAGGATCATTTCTGGGATGGGAAGGGATGTGAGAT
GATCTGTTACTGCAACTTCAGCGAATTGCTCTGCTGCCCAAAGACGTTTTCTTTGGACC
AAAGATCTCTTTTCGTGATTCTTGCAACAATCAATGAGAATCTTCATGTATTCTGGAGAA
CACCATTCTGATTTCCACAAACTGCACTACATCAGTATAACTGCATTTCTAGTTTCTA
TATAGTGCAATAGAGCATAGATTCTATAAATTCTTACTTGTCTAAGACAAGTAAATCTGT
GTTAAACAAGTAGTAATAAAAGTTAATTCAATCTAAAAAAAAAAAAA

99/392

FIGURE 94

MKLMVLVFTIGLTLLLGVQAMPANRLSCYRKILKDHNCHNLPEGVADLTQIDVNVQDHFWDGKGCEMICYCNFSELLCCPKDVFFGPKISFVIPCNNQ

Important features:

Signal peptide:

amino acids 1-20

N-glycosylation site:

amino acids 72-76

Tyrosine kinase phosphorylation site:

amino acids 63-71

100/392

FIGURE 95

GAATTCGGGGCCCCAGGATGCCAACTTTGAATAGGATGAAGACTACAACTTGTTCCCTTC
TCATCTGCATCTCCCTGCTCCAGCTGATGGTCCCAGTGAATACTGATGAGACCATAGAGA
TTATCGTGGAGAATAAGGTCAAGGAACTTCTTGCCAATCCAGCTAACTATCCCTCCACTG
TAACGAAGACTCTCTCTTGCACTAGTGTCAAGACTATGAACAGATGGGCCTCCTGCCCTG
CTGGGATGACTGCTACTGGGTGTGCTTGTGGCTTTGCCTGTGGATCTTGGGAGATCCAGA
GTGGAGATACTTGCAACTGCCTGTGCTTACTCGTTGACTGGACCACTGCCCCGCTGCTGCC
AACTGTCCTAAGAATGAAGAGGTGGAGAACCAGCTTTGATATGATGAATCTAACAAAAA
CTGCAGTCTCAATTTGGAAATCTGACTCATGTGCCTTTAAATGTGTTCATATTGCCCATT
TACCCTGCTTCTTGAAATGCTTCTTGAAAAATAAAGACAAATTTGCATGTG

101/392

FIGURE 96

MKTTTCSELLICISLLQLMVPVNTDETIEIIVENKVKELLANPANYPSTVTKTLSCTSVK
TMNRWASCPAGMTATGCACGFACGSWEIQSGDTCNCLCLLDWTTARCCQLS

102/392

FIGURE 97

GAGGCAGAAAGGCAGAAAGGAGAAAATTTCAGGATAACTCTCCTGAGGGGTGAGCCAAGCC
CTGCCATGTTAGTGCACGCAGGACATCAACAAACACAGATAACAGGAAATGATCCATTCCC
TGTGGTCACTTATTCTAAAGGCCCAACCTTCAAAGTTCAAGTAGTGATATGGATGACTC
CACAGAAAGGGAGCAGTCACGCCTTACTTCTTGCCTTAAGAAAAGAGAAGAAATGAAACT
GAAGGAGTGTGTTTTCCATCCTCCCACGGAAGGAAAGCCCCCTCTGTCCGATCCTCCAAAGA
CGGAAAGCTGCTGGCTGCAACCTTGCTGCTGGCACTGCTGTCTTGCTGCCTCACGGTGGT
GTCTTTCTACCAGGTGGCCGCCCTGCAAGGGGACCTGGCCAGCCTCCGGGCAGAGCTGCA
GGGCCACCACGCGGAGAAGCTGCCAGCAGGAGCAGGAGCCCCCAAGGCCGGCCTGGAGGA
AGCTCCAGCTGTACCGCGGGACTGAAAATCTTTGAACCACCAGCTCCAGGAGAAGGCAA
CTCCAGTCAGAACAGCAGAAATAAGCGTGCCGTTTCAGGGTCCAGAAGAAACAGTCACTCA
AGACTGCTTGCAACTGATTGCAGACAGTGAAACACCAACTATACAAAAAGGATCTTACAC
ATTTGTTCCATGGCTTCTCAGCTTTAAAAGGGGAAGTGCCCTAGAAGAAAAAGAGAATAA
AATATTGGTCAAAGAACTGGTTACTTTTTTATATATGGTCAGGTTTTATATACTGATAA
GACCTACGCCATGGGACATCTAATTCAGAGGAAGAAGGTCCATGTCTTTGGGGATGAATT
GAGTCTGGTGACTTTGTTTCGATGTATTCAAATATGCCTGAAACACTACCCAATAATTC
CTGCTATTCAGCTGGCATTGCAAACTGGAAGAAGGAGATGAACTCCAATTGCAATACC
AAGAGAAAATGCACAAATATCACTGGATGGAGATGTCACATTTTTTGGTGCATTGAACT
GCTGTGACCTACTTACACCATGTCTGTAGCTATTTTCCTCCCTTTCTCTGTACCTCTAAG
AAGAAAGAATCTAACTGAAAATACCAAAAAAAAAAAAAAAAAA

103/392

FIGURE 98

MDDSTEREQSRLTSCLKKREEMKLKECVSILPRKESPSVRSSKDGKLLAATLLLALLSCC
LTVVSFYQVAALQGD LASLRAELQGHHA EKLPAGAGAPKAGLEEAPAVTAGLKIFEPPAP
GEGNSSQNSRNKRAVQGPEETVTQDCLQLIADSETPTIQKGSYTFVPWLLSFKRGSAL EE
KENKILVKETGYFFIYGQVLYTDKTYAMGH LIQRKKVHVFGDELSLVTLFRCIQNMPETL
PNNSCYSAGIAKLEEGDELQLAIPRENAQISLDGDVTFFGALKLL

Transmembrane domain:
amino acids 47-72

N-glycosylation site:
amino acids 124-127, 242-245

**cAMP- and cGMP-dependent protein kinase phosphorylation
site:**
amino acids 33-36, 173-176

N-myristoylation site:
amino acids 96-101

TNF family proteins:
amino acids 172-206

104/392

FIGURE 99

CCGAGGTGGCGATCGCTGAGAGGCAGGAGGGCCGAGGCGGGCCTGGGAGGCGGCCCGGAG
GTGGGGCGCCGCTGGGGCCGGCCCGCACGGGCTTCATCTGAGGGCGCACGGCCCGCGACC
GAGCGTGCGGACTGGCCTCCCAAGCGTGGGGCGACAAGCTGCCGGAGCTGCAATGGGCCG
CGGCTGGGGATTCTTGTGTTGGCCTCCTGGGCGCCGTGTGGCTGCTCAGCTCGGGCCACGG
AGAGGAGCAGCCCCCGGAGACAGCGGCACAGAGGTGCTTCTGCCAGGTTAGTGGTTACTT
GGATGATTGTACCTGTGATGTTGAAACCATTGATAGATTTAATAACTACAGGCTTTTCCC
AAGACTACAAAACTTCTTGAAAGTGACTACTTTAGGTATTACAAGGTAAACCTGAAGAG
GCCGTGTCCTTTCTGGAATGACATCAGCCAGTGTGGAAGAAGGGACTGTGCTGTCAAACC
ATGTCAATCTGATGAAGTTCCTGATGGAATTAAATCTGCGAGCTACAAGTATTCTGAAGA
AGCCAATAATCTCATTGAAGAATGTGAACAAGCTGAACGACTTGGAGCAGTGGATGAATC
TCTGAGTGAGGAAACACAGAAGGCTGTTCTTCAGTGGACCAAGCATGATGATTCTTCAGA
TAACCTTCTGTGAAGCTGATGACATTGAGTCCCCTGAAGCTGAATATGTAGATTTGCTTCT
TAATCCTGAGCGCTACACTGGTTACAAGGGACCAGATGCTTGGAAAATATGGAATGTCAT
CTACGAAGAAAACGTGTTTTAAGCCACAGACAATTAAGAACCTTTAAATCCTTTGGCTTC
TGGTCAAGGGACAAGTGAAGAGAACACTTTTTACAGTTGGCTAGAAGGTCTCTGTGTAGA
AAAAAGAGCATTCTACAGACTTATATCTGGCCTACATGCAAGCATTAATGTGCATTTGAG
TGCAAGATATCTTTTACAAGAGACCTGGTTAGAAAAGAAATGGGGACACAACATTACAGA
ATTTCAACAGCGATTTGATGGAATTTTGAAGTGAAGGAGAAGGTCCAAGAAGGCTTAAGAA
CTTGATATTTTCTCTACTTAATAGAACTAAGGGCTTTATCCAAAGTGTTACCATTCTTCGA
GCGCCAGATTTTCAACTCTTTACTGGAAATAAAATTCAGGATGAGGAAAACAAAATGTT
ACTTCTGGAAATACTTCATGAAATCAAGTCATTTCTTTGCATTTTGATGAGAATTCATT
TTTTGCTGGGGATAAAAAAGAAGCACACAACTAAAGGAGGACTTTCGACTGCATTTTAG
AAATATTTCAAGAATTATGGATTGTGTTGGTTGTTTTAAATGTCGTCTGTGGGGAAAGCT
TCAGACTCAGGGTTTGGGCACCTGCTCTGAAGATCTTATTTTCTGAGAAATTGATAGCAA
TATGCCAGAAAGTGGACCTAGTTATGAATTCCATCTAACCAGACAAGAAATAGTATCATT
ATTCAACGCATTTGGAAGAATTTCTACAAGTGTGAAAGAATTAGAAAACCTTCAGGAACTT
GTTACAGAATATTCATTAAAGAAAACAAGCTGATATGTGCCTGTTTCTGGACAATGGAGG
CGAAAGAGTGGAATTTTCATTCAAAGGCATAATAGCAATGACAGTCTTAAGCCAAACATTT
TATATAAAGTTGCTTTTGTAAAGGAGAATTATATTGTTTAAAGTAAACACATTTTAAAA
ATTGTGTTAAGTCTATGTATAATACTACTGTGAGTAAAAGTAATACTTTAATAATGTGGT
ACAAATTTTAAAGTTTAAATATTGAATAAAAGGAGGATTATCAAATTAATAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAA

105/392

FIGURE 100

MGRGWGFLFGLLGAVWLLSSGHGEEQPPETAARCFQVSGYLDDCTCDVETIDRFNNYR
LFPRLQKLLSDYFRYYKVNLRPCPFWNDISQCGRRDCAVKPCQSDEVPDGIKSASYKY
SEEANNLIEECEQAERLGAVDESLSEETQKAVLQWTKHDDSSDNFCEADDIQSPEAEYVD
LLNPERYTGKGPDAWKIWNVIYEENCFKPQTIKRPLNPLASGQGTSEENTFYSWLEGL
CVEKRAFYRLISGLHASINVHLSARYLLQETWLEKKWGHNITEFQQRFDGILTEGEGPRR
LKNLYFLYLIELRALSKVLPFFERPDLFTGNKIQDEENKMLLLEILHEIKSFPLHFDE
NSFFAGDKKEAHKLKEDFRLHFRNISRIMDCVGCFCRLWGKLQTQGLGTALKILFSEKL
IANMPESGPSYEFHLTRQEIIVSLFNAFGRISTSVKELENFRNLLQNIH

Important features:**Signal peptide:**

amino acids 1-23

N-glycosylation site:

amino acids 280-283 and 384-387

Amidation site:

amino acids 94-97

Glycosaminoglycan attachment site:

amino acids 20-23 and 223-226

Aminotransferases class-V pyridoxal-phosphate:

amino acids 216-222

Interleukin-7 proteins:

amino acids 338-343

106/392

FIGURE 101

GCCTAGCCAGGCCAAGAATGCAATTGCCCCGGTGGTGGGAGCTGGGAGACCCCTGTGCTT
GGACGGGACAGGGTCGGGGGACACGCAGGATGAGCCCCGCGACCACTGGCACATTCTTG
TGACAGTGTACAGTATTTTCTCCAAGGTACACTCCGATCGGAATGTATACCCATCAGCAG
GTGTCTCTTTGTTTCATGTTTTGGAAAGAGAATATTTTAAGGGGGAATTTCCACCTTACC
CAAAACCTGGCGAGATTAGTAATGATCCCATAACATTTAATACAAATTTAATGGGTTACC
CAGACCGACCTGGATGGCTTCGATATATCCAAAGGACACCATATAGTGATGGAGTCCTAT
ATGGGTCCCCAACAGCTGAAAATGTGGGGAAGCCAACAATCATTGAGATAACTGCCTACA
ACAGGCGCACCTTTGAGACTGCAAGGCATAATTTGATAATTAATATAATGTCTGCAGAAG
ACTTCCCGTTGCCATATCAAGCAGAATTCCTTCATTAAAGAATATGAATGTAGAAGAAATGT
TGGCCAGTGAGGTTCTTGGGAGACTTTCTTGGCGCAGTGAAAAATGTGTGGCAGCCAGAGC
GCCTGAACGCCATAAACATCACATCGGCCCTAGACAGGGGTGGCAGGGTGGCACTTCCCA
TTAATGACCTGAAGGAGGGCGTTTATGTCATGGTTGGTGCAGATGTCCCGTTTTCTTCTT
GTTTACGAGAAGTTGAAAATCCACAGAATCAATTGAGATGTAGTCAAGAAATGGAGCCTG
TAATAACATGTGATAAAAAATTTTCGTACTCAATTTTACATTGACTGGTGCAAAATTTTCAT
TGGTTGATAAAACAAAGCAAGTGTCCACCTATCAGGAAGTGATTCGTGGAGAGGGGATTT
TACCTGATGGTGGAGAATACAAACCCCCCTTCTGATTCTTTGAAAAGCAGAGACTATTACA
CGGATTTCTTAATTACACTGGCTGTGCCCTCGGCAGTGGCACTGGTCCTTTTTCTAATAC
TTGCTTATATCATGTGCTGCCGACGGGAAGGCGTGGAAAAGAGAAACATGCAAAACACCAG
ACATCCAACCTGGTCCATCACAGTGCTATTTCAGAAATCTACCAAGGAGCTTCGAGACATGT
CCAAGAATAGAGAGATAGCATGGCCCCGTGCAACGCTTCCTGTGTTCCACCCTGTGACTG
GGGAAATCATACCTCCTTTACACACAGACAACTATGATAGCACAAACATGCCATTGATGC
AAACGCAGCAGAACTTGCCACATCAGACTCAGATCCCCAACAGCAGACTACAGGTAAAT
GGTATCCCTGAAGAAAGAAAACCTGACTGAAGCAATGAATTTATAATCAGACAATATAGCA
GTTACATCACATTTCTTTTCTTTCCAATAATGCATGAGCTTTTCTGGCATATGTTATGC
ATGTTGGCAGTATTAAGTGTATACCAAATAATACAACATAACTTTCATTTTACTAATGTA
TTTTTTTGTACTTAAAGCATTTTTTGACAATTTGTAAAACATTGATGACTTTATATTTGTT
ACAATAAAAGTTGATCTTTAAAATAAATATTATTAATGAAGCCTAAAAA

107/392

FIGURE 102

MQLPRWWELGDPCAWTGQGRGTRRMSPATTGTFLTVYSIFSKVHSDRNVYPSAGVLFVH
VLEREYFKGEFPPYPKPGEISNDPITFNTNLMGYPDPRPGWLRYIQRTPYSDGVLYGSPTA
ENVGKPTIIETAYNRRTFETARHNLIIINIMSAEDFPLPYQAEFFIKNMNVEEMLASEVL
GDFLGAVKNVWQPERLNAINITSALDRGGRVPLPINDLKEGVYVMVGADVPFSSCLREVE
NPQNQLRCSQEMEPVITCDKKFRTQFYIDWCKISLVDKTKQVSTYQEVIRGEGILPDGGE
YKPPSDSLKSRDYYTDFLITLAVPSAVALVFLILAYIMCCRREGVEKRNMQTPDIQLVH
HSAIQKSTKELRDMSKNREIAWPLSTLPVFHPVTGEIIPPLHTDNYDSTNMPLMQTQQNL
PHQTQIPQQOTTGKWYP

signal sequence:

Amino acids 1-46

transmembrane domain:

Amino acids 319-338

N-glycosylation site:

Amino acids 200-204

**cAMP- and cGMP-dependent protein kinase phosphorylation
site:**

Amino acids 23-27

Tyrosine kinase phosphorylation site:

Amino acids 43-52

N-myristoylation sites:

Amino acids 17-23;112-118;116-122;185-191

108/392

FIGURE 103

CAGAAGAGGGGGCTAGCTAGCTGTCTCTGCGGACCAGGGAGACCCCCGCGCCCCCCCCGGT
GTGAGGCGGCCTCACAGGGCCGGGTGGGCTGGCGAGCCGACGCGGCGGCGGAGGAGGCTG
TGAGGAGTGTGTGGAACAGGACCCGGGACAGAGGAACCAATGGCTCCGCAGAACCTGAGCA
CCTTTTGCCTGTTGCTGCTATACCTCATCGGGGCGGTGATTGCCGGACGAGATTTCTATA
AGATCTTGGGGGTGCCTCGAAGTGCCTCTATAAAGGATATTAAAAAGGCCTATAGGAAAC
TAGCCCTGCAGCTTCATCCCGACCGGAACCCCTGATGATCCACAAGCCCAGGAGAAATTCC
AGGATCTGGGTGCTGCTTATGAGGTTCTGTGAGATAGTGAGAAACGGAAACAGTACGATA
CTTATGGTGAAGAAGGATTAAAAGATGGTCATCAGAGCTCCCATGGAGACATTTTTTTCAC
ACTTCTTTGGGGATTTTTGGTTTTCATGTTTTGGAGGAACCCCTCGTCAGCAAGACAGAAATA
TTCCAAGAGGAAGTGATATTATTGTAGATCTAGAAGTCACTTTGGAAGAAGTATATGCAG
GAAATTTTGTGGAAGTAGTTAGAAACAAACCTGTGGCAAGGCAGGCTCCTGGCAAACGGA
AGTGCAATTGTGCGCAAGAGATGCGGACCACCCAGCTGGGGCCCTGGGCGCTTCCAAATGA
CCCAGGAGGTGGTCTGCGACGAATGCCCTAATGTCAAACCTAGTGAATGAAGAACGAACGC
TGGAAGTAGAAATAGAGCCTGGGGTGAGAGACGGCATGGAGTACCCCTTTATTGGAGAAG
GTGAGCCTCACGTGGATGGGGAGCCTGGAGATTTACGGTTCCGAATCAAAGTTGTCAAGC
ACCCAATATTTGAAAGGAGAGGAGATGATTTGTACACAAATGTGACAATCTCATTAGTTG
AGTCACTGGTTGGCTTTGAGATGGATATTACTCACTTGGATGGTCACAAGGTACATATTT
CCCGGGATAAGATCACCAGGCCAGGAGCGAAGCTATGGAAGAAAGGGGAAGGGCTCCCCA
ACTTTGACAACAACAATATCAAGGGCTCTTTGATAATCACTTTTGATGTGGATTTTCCAA
AAGAACAGTTAACAGAGGAAGCGAGAGAAGGTATCAAACAGCTACTGAAACAAGGGTCAG
TGCAGAAGGTATACAATGGACTGCAAGGATATTAGAGAGTGAATAAAAATTGGACTTTGTTT
AAAATAAGTGAATAAGCGATATTTATTATCTGCAAGGTTTTTTTTTGTGTGTGTTTTTGT
TTATTTTCAATATGCAGTGTAGGCTTAATTTTTTTTATCTAATGATCATCATGAAATGAAT
AAGAGGGCTTAAGAATTTGTCCATTTGCATTTCGGAAGAAAGAAATGACCAGCAAAAGGTTTAC
TAATACCTCTCCCTTTGGGGATTTAATGTCTGGTGCTGCCGCCTGAGTTTCAAGAATTAA
AGCTGCAAGAGGACTCCAGGAGCAAAAGAAACACAATATAGAGGGTTGGAGTTGTTAGCA
ATTTCAATTCAAAATGCCAACTGGAGAAGTCTGTTTTTAAATACATTTTGTGTATTATTTTAA

109/392

FIGURE 104

MAPQNLSTFCLLLLYLIGAVIAGRDFYKILGVPRASIKDIKKAYRKLALQLHPDRNPDD
PQAQEKFDLGAAYEVLSDSEKQYDQTYGEEGLKDGHQSSHGDI FSHFFGDFGFMFGGT
PRQQDRNIPRGSDIIVDLEVTLEEVYAGNFVEVVRNKPVARQAPGKRKCNCRQEMRTTQL
GPGRFQMTQEVVCDECPNVKLVNEERTLEVEIEPGVRDGMETPFGEGEPHVDGEPGDLR
FRIKVVKHPIFERRGDDLYTNVTISLVESLVGFEMDITHLDGHKVHISRDKITRPGAKLW
KKGEGLPNFDNNNIKGSLIITFDVDFPKEQLTEEAREGIKQLLKQGSVQKVYNGLQGY

Important features:**Signal peptide:**

amino acids 1-22

Cell attachment sequence:

amino acids 254-257

Nt-dnaJ domain signature:

amino acids 67-87

Homologous region to Nt-dnaJ domain proteins:

amino acids 26-58

N-glycosylation site:

amino acids 5-9, 261-265

Tyrosine kinase phosphorylation site:

amino acids 253-260

N-myristoylation site:

amino acids 18-24, 31-37, 93-99, 215-221

Amidation site:

amino acids 164-168

110/392

FIGURE 105

GGCACGAGGCGGCGGGGCAGTCGCGGGATGCGCCCGGAGCCACAGCCTGAGGCCCTCAG
GTCTCTGCAGGTGTCGTGGAGGAACCTAGCACCTGCCATCCTCTTCCCCAATTTGCCACT
TCCAGCAGCTTTAGCCCATGAGGAGGATGTGACCGGGACTGAGTCAGGAGCCCTCTGGAA
GCATGGAGACTGTGGTGATTGTTGCCATAGGTGTGCTGGCCACCATCTTTCTGGCTTCGT
TTGCAGCCTTGGTGCTGGTTTGCAGGCAGCGCTACTGCCGCGCGGAGACCTGCTGCAGC
GCTATGATTCTAAGCCCATTTGTGGACCTCATTGGTGCCATGGAGACCCAGTCTGAGCCCT
CTGAGTTAGAACTGGACGATGTCGTTATCACCAACCCCCACATTGAGGCCATTCTGGAGA
ATGAAGACTGGATCGAAGATGCCTCGGGTCTCATGTCCCACTGCATTGCCATCTTGAAGA
TTTGTCACTCTGACAGAGAAGCTTGTTGCCATGACAATGGGCTCTGGGGCCAAGATGA
AGACTTCAGCCAGTGTGACGACATCATTTGTGGTGGCCAAGCGGATCAGCCCCAGGGTGG
ATGATGTTGTGAAGTCGATGTACCCTCCGTTGGACCCCCAACTCCTGGACGCACGGACGA
CTGCCCTGCTCCTGTCTGTGTCAGTCACCTGGTGCTGGTGACAAGGAATGCCTGCCATCTGA
CGGGAGGCCTGGACTGGATTGACCAGTCTCTGTGCGCTGCTGAGGAGCATTTGGAAGTCC
TTCGAGAAGCAGCCCTAGCTTCTGAGCCAGATAAAGGCCTCCCAGGCCCTGAAGGCTTCC
TGCAGGAGCAGTCTGCAATTTAGTGCCTACAGGCCAGCAGCTAGCCATGAAGGCCCTGC
CGCCATCCCTGGATGGCTCAGCTTAGCCTTCTACTTTTTTCTATAGAGTTAGTTGTTCTC
CACGGCTGGAGAGTTTACGCTGTGTGTGCATAGTAAAGCAGGAGATCCCCGTCAGTTTATG
CCTCTTTTGCAGTTGCAAACTGTGGCTGGTGAGTGGCAGTCTAATACTACAGTTAGGGGA
GATGCCATTCACCTCTCTGCAAGAGGAGTATTGAAAACCTGGTGGACTGTCAGCTTTATTTA
GCTCACCTAGTGTTTTTCAAGAAAATTGAGCCACCGTCTAAGAAATCAAGAGGTTTACAT
TAAATTAGAATTTCTGGCCTCTCTCGATCGGTCAGAATGTGTGGCAATTCTGATCTGCA
TTTTTCAAGAGGACAATCAATTGAACTAAGTAGGGGTTTCTTCTTTTGGCAAGACTTG
TACTCTCTCACCTGGCCTGTTTTCATTTATTTGTATTATCTGCCTGGTCCCTGAGGCGTCT
GGGTCTCTCCTCTCCCTTGCAGGTTTGGGTTTGAAGCTGAGGAACTACAAAGTTGATGAT
TTCTTTTTTATCTTTATGCCTGCAATTTTACCTAGCTACCACTAGGTGGATAGTAAATTT
ATACTTATGTTTCCCTCAAAAAAAAAAAAAAA

111/392

FIGURE 106

METVVIVAIGVLATIFLASFAALVLVCRQRYCRPRDLLQRYDSKPIVDLIGAMETQSEPS
ELELDDVVITNPHIEAILENEDWIEDASGLMSHCIAILKICHTLTEKLVAMTMGSGAKMK
TSASVSDIIVVAKRISPRVDDVVKSMYPPLDPKLLDARTTALLSVSHLVLVTRNACHLT
GGLDWIDQSLSAAEHLEVLREAALASEPDKGLPGPEGFLQEQSAI

112/392

FIGURE 107

GCTTCATTTCTCCCGACTCAGCTTCCCACCCCTGGGCTTTCCGAGGTGCTTTGCGCCGCTGT
CCCCACCACTGCAGCCATGATCTCCTTAACGGACACGCAGAAAATTGGAATGGGATTAAC
AGGATTTGGAGTGTTTTCTGTTCTTTGGAATGATTCTCTTTTTTGACAAAGCACTACT
GGCTATTGGAATGTTTTATTTGTAGCCGGCTTGGCTTTTGTAATTGGTTTAGAAAGAAC
ATTCAGATTCTTCTTCCAAAAACATAAAATGAAAGCTACAGGTTTTTTCTGGGTGGTGT
ATTTGTAGTCCTTATTGGTTGGCCTTTGATAGGCATGATCTTCGAAATTTATGGATTTTT
TCTCTTGTTCAAGGGCTTCTTTCCTGTCGTTGTTGGCTTTATTAGAAGAGTGCCAGTCCT
TGGATCCCTCCTAAATTTACCTGGAATTAGATCATTTGTAGATAAAGTTGGAGAAAGCAA
CAATATGGTATAAACAACAAGTGAATTTGAAGACTCATTTAAAATATTGTGTTATTTATAA
AGTCATTTGAAGAATATTCAGCACAAAATTAAATTACATGAAATAGCTTGTAATGTTCTT
TACAGGAGTTTAAAACGTATAGCCTACAAAGTACCAGCAGCAAATTAGCAAAGAAGCAGT
GAAAACAGGCTTCTACTCAAGTGAACCTAAGAAGAAGTCAGCAAGCAAAGTGAGAGAGGTG
AAATCCATGTTAATGATGCTTAAGAACTCTTGAAGGCTATTTGTGTTGTTTTCCACAA
TGTGCGAACTCAGCCATCCTTAGAGAACTGTGGTGCCTGTTTCTTTTCTTTTATTTTG
AAGGCTCAGGAGCATCCATAGGCATTTGCTTTTTTAGAAGTGTCCACTGCAATGGCAAAA
TATTTCCAGTTGCACTGTATCTCTGGAAGTGATGCATGAATTCGATTGGATTGTGTCATT
TTAAAGTATTAAAACCAAGGAAACCCCAATTTTGATGTATGGATTACTTTTTTTTGNGCN
CAGGGCC

113/392

FIGURE 108

MISLTDQTQKIGMGLTGFGVFVFLFFGMILFFDKALLAIGNVLFVAGLAFVIGLERTFRFFF
QKHKMKATGFFLGGVFVVLIGWPLIGMIFEIYGFFLLFRGFFPVVVGFIRRVFVLGSLLN
LPGIRSFVDKVGESNNMV

Important features:**Transmembrane domains:**

amino acids 12-30 (typeII), 33-52, 69-89 and 93-109

N-myristoylation sites:

amino acids 11-16, 51-56 and 116-121

Aminoacyl-transfer RNA synthetases class-II protein:

amino acids 49-59

114/392

FIGURE 109

CCAGTCTGTCGCCACCTCACTTGGTGTCTGCTGTCCCCGCCAGGCAAGCCTGGGGTGAGA
GCACAGAGGAGTGGGCCGGGACCATGCGGGGGACGCGGCTGGCGCTCCTGGCGCTGGTGC
TGGCTGCCTGCGGAGAGCTGGCGCCGGCCCTGCGCTGCTACGTCTGTCCGGAGCCCACAG
GAGTGTGCGACTGTGTCACCATCGCCACCTGCACCACCAACGAAACCATGTGCAAGACCA
CACTCTACTCCCGGGAGATAGTGTACCCCTTCCAGGGGGACTCCACGGTGACCAAGTCCT
GTGCCAGCAAGTGTAAGCCCTCGGATGTGGATGGCATCGGCCAGACCCTGCCCCTGTCCT
GCTGCAATACTGAGCTGTGCAATGTAGACGGGGCGCCCGCTCTGAACAGCCTCCACTGCG
GGGCCCTCACGCTCCTCCCACTCTTGAGCCTCCGACTGTAGAGTCCCCGCCACCCCCAT
GGCCCTATGCGGCCAGCCCCGAATGCCTTGAAGAAGTGCCCCCTGCACCAGGAAAAAA
AAAAA

115/392

FIGURE 110

MRGTRLALLALVLAACGELAPALRCYVCPEPTGVSDCVTIATCTTNETMCKTTLYSREIV
YPFQGDSTVTKSCASKCKPSDVDGIGQTLFVSCCNTELCNVDGAPALNSLHCGALTLLPL
LSLRL

Important features:

Signal peptide:

amino acids 1-17

N-glycosylation site:

amino acids 46-49

116/392

FIGURE 111

GCGCCGCCAGGCGTAGGCGGGGTGGCCCTTGCGTCTCCCGCTTCCTTGAAAAACCCGGCG
GGCGAGCGAGGCTGCGGGCCGGCCGCTGCCCTTCCCCACACTCCCCGCCGAGAAGCCTCG
CTCGGCGCCCAACATGGCGGGTGGGCGCTGCGGCCCGCAGCTAACGGCGCTCCTGGCCGC
CTGGATCGCGGCTGTGGCGGCGACGGCAGGCCCCGAGGAGGCCGCGCTGCCGCCGGAGCA
GAGCCGGGTCCAGCCCATGACCGCCTCCAACCTGGACGCTGGTGATGGAGGGCGAGTGGAT
GCTGAAATTTTACGCCCCATGGTGTCCATCCTGCCAGCAGACTGATTGAGAATGGGAGGC
TTTTGCAAAGAATGGTGAAATACCTCAGATCAGTGTGGGGAAGGTAGATGTCATTCAAGA
ACCAGGTTTGAGTGGCCGCTTCTTTGTCACCACTCTCCCAGCATTTTTTCATGCAAAGGA
TGGGATATTCCGCCGTTATCGTGGCCAGGAATCTTCGAAGACCTGCAGAATTATATCTT
AGAGAAGAAATGGCAATCAGTCGAGCCTCTGACTGGCTGGAAATCCCCAGCTTCTCTAAC
GATGCTCTGGAATGGCTGGTCTTTTTAGCATCTCTGGCAAGATATGGCATCTTCACAACTA
TTTCACAGTGACTCTTGGAATTCCTGCTTGGTGTCTTATGTGTTTTTCGTCATAGCCAC
CTTGGTTTTTGGCCTTTTTATGGGTCTGGTCTTGGTGGTAATATCAGAATGTTTCTATGT
GCCACTTCCAAGGCATTTATCTGAGCGTCTCTGAGCAGAATCGGAGATCAGAGGAGGCTCA
TAGAGCTGAACAGTTGCAGGATGCGGAGGAGGAAAAAGATGATTCAAATGAAGAAGAAAA
CAAAGACAGCCTTGTAGATGATGAAGAAGAGAAAGAAGATCTTGGCGATGAGGATGAAGC
AGAGGAAGAAGAGGAGGAGGACAACCTGGCTGCTGGTGTGGATGAGGAGAGAAGTGAGGC
CAATGATCAGGGGGCCCCCAGGAGAGGACGGTGTGACCCGGGAGGAAGTAGAGCCTGAGGA
GGCTGAAGAAGGCATCTCTGAGCAACCCTGCCAGCTGACACAGAGGTGGTGGAAAGACTC
CTTGAGGCAGCGTAAAAGTCAGCATGCTGACAAGGGACTGTAGATTTAATGATGCGTTTTT
CAAGAATACACACCAAAACAATATGTCAGCTTCCCTTTGGCCTGCAGTTTGTACCAAATC
CTTAATTTTTTCTGAATGAGCAAGCTTCTCTTAAAAGATGCTCTCTAGTCATTTGGTCTC
ATGGCAGTAAGCCTCATGTATACTAAGGAGAGTCTTCCAGGTGTGACAATCAGGATATAG
AAAAACAAACGTAGTGTTGGGATCTGTTTGGAGACTGGGATGGGAACAAGTTCATTTACT
TAGGGGTGAGAGAGTCTCGACCAGAGGAGGCCATTTCCAGTCCTAATCAGCACCTTCCAG
AGACAAGGCTGCAGGCCCTGTGAAATGAAAGCCAAGCAGGAGCCTTGGCTCCTGAGCATC
CCCAAAGTGTAACGTAGAAGCCTTGCATCCTTTTCTTGTGTAAAGTATTTATTTTTGTCA
AATTGCAGGAAACATCAGGCACCACAGTGCATGAAAAATCTTTCACAGCTAGAAATTGAA
AGGGCCTTGGGTATAGAGAGCAGCTCAGAAGTCATCCAGCCCTCTGAATCTCCTGTGCT
ATGTTTTATTTCTTACCTTTAATTTTTCCAGCATTTCCACCATGGGCATTCAGGCTCTCC
ACACTCTTCACTATTATCTCTTGGTCAGAGGACTCCAATAACAGCCAGGTTTACATGAAC
TGTGTTTGTTTCATTCTGACCTAAGGGGTTTAGATAATCAGTAACCATAACCCCTGAAGCT
GTGACTGCCAAACATCTCAAATGAAATGTTGTGGCCATCAGAGACTCAAAGGAAGTAAG
GATTTTACAAGACAGATTAAAAAAAATTGTTTTGTCCAAAATATAGTTGTTGTTGATTT
TTTTTTAAGTTTTCTAAGCAATATTTTTCAAGCCAGAAGTCCTCTAAGTCTTGCCAGTAC
AAGGTAGTCTTGTGAAGAAAAGTTGAATACTGTTTTGTTTTCATCTCAAGGGGTTCCCTG
GGTCTTGAACACTTTAATAATAACTAAAAAACCACTTCTGATTTTCCTTCAGTGATGTG
CTTTTGGTGAAAGAATTAATGAACTCCAGTACCTGAAAGTGAAAGATTTGATTTTGTTTC
CATCTTCTGTAATCTTCCAAAGAATTATCTTTGTAAATCTCTCAATACTCAATCTACT
GTAAGTACCCAGGGAGGCTAATTCTTT

117/392

FIGURE 112

MAGGRCGPQLTALLAAWIAAVAATAGPEEAALPPEQSRVQPMASNWTLMMEGEWMLKFY
APWCPSCQQTDSWEAFAKNGEILQISVGKVDVIQEPGLSGRFFVTTLPAFFHAKDGI FR
RYRGP G IFEDLQNYILEKKWQSVEPLTGWKSPASLTMSGMAGLFSISGKIWHLHNYFTVT
LGIPAWCSYVFFVIATLVFGLFMGLVLVVISSECFYVPLPRHLSESEQNRRSEEAHRAEQ
LQDAEEEEKDDSNEEENKDSLVDDEEEKEDLGDEDEAEEDNLAAGVDEERSEANDQG
PPGEDGVTREEVEPEEAEEGISEQPCPADTEVVEDSLRQRKSQHADKGL

Important features:**Signal peptide:**

amino acids 1-22

Transmembrane domain:

amino acids 191-211

N-glycosylation site:

amino acids 46-49

Thioredoxin family proteins: (homologous region to disulfide
isomerase)

amino acids 56-72

Flavodoxin proteins:

amino acids 173-187

118/392

FIGURE 113

GAGGAACCTACCGGTACCGGCCGCGCGCTGGTAGTCGCCGGTGTGGCTGCACCTCACCAA
TCCCGTGCGCCGCGGCTGGGCCGTCGGAGAGTGCCTGTGCTTCTCTCCTGCACGCGGTGC
TTGGGCTCGGCCAGGCGGGTCCGCCGCCAGGGTTTGAGGATGGGGAGTAGCTACAGGA
AGCGACCCCGCGATGGCAAGGTATATTTTTGTGGAATGAAAAGGAAGTATTAGAAATGAG
CTGAAGACCATTACAGATTAATATTTTTGGGGACAGATTTGTGATGCTTGATTCACCCCT
TGAAGTAATGTAGACAGAAGTTCTCAAATTTGCATATTACATCAACTGGAACCAGCAGTG
AATCTTAATGTTCACTTAAATCAGAACTTGCATAAGAAAGAGAATGGGAGTCTGGTTAAA
TAAAGATGACTATATCAGAGACTTGAAAAGGATCATTTCTCTGTTTTCTGATAGTGTATAT
GGCCATTTTAGTGGGCACAGATCAGGATTTTTACAGTTTACTTGGAGTGTCCAAAACCTGC
AAGCAGTAGAGAAATAAGACAAGCTTTCAAGAAATTGGCATTGAAGTTACATCCTGATAA
AAACCCGAATAACCCAAATGCACATGGCGATTTTTTAAAAATAAATAGAGCATATGAAGT
ACTCAAAGATGAAGATCTACGGAAAAAGTATGACAAATATGGAGAAAAGGGACTTGAGGA
TAATCAAGGTGGCCAGTATGAAAGCTGGAACATTATCGTTATGATTTTGGTATTTATGA
TGATGATCCTGAAATCATAACATTGGAAAAGAGAATTTGATGCTGCTGTTAATTCTGG
AGAACTGTGGTTTGTAATTTTTACTCCCCAGGCTGTTCACTGCCATGATTTAGCTCC
CACATGGAGAGACTTTGCTAAAGAAGTGGATGGGTACTTCGAATTGGAGCTGTAACTG
TGGTGAATGATAGAATGCTTTGCCGAATGAAAGGAGTCAACAGCTATCCAGTCTCTTCAT
TTTTCGGTCTGGAATGGCCCCAGTGAAATATCATGGAGACAGATCAAAGGAGAGTTTAGT
GAGTTTTGCAATGCAGCATGTTAGAAGTACAGTGACAGAACTTTGGACAGGAAATTTTGT
CAACTCCATACAACTGCTTTTGCTGCTGGTATTGGCTGGCTGATCACTTTTTGTTCAAA
AGGAGGAGATTGTTTGACTTCACAGACACGACTCAGGCTTAGTGGCATGTTGTTTCTCAA
CTCATTGGATGCTAAAGAAATATATTTGGAAGTAATACATAATCTTCCAGATTTTGAAC
ACTTTTCGGCAAACACACTAGAGGATCGTTTGGCTCATCATCGGTGGCTGTTATTTTTTCA
TTTTGGAAAAAATGAAAATTCAAATGATCCTGAGCTGAAAAAACTAAAACTCTACTTAA
AAATGATCATATTCAAGTTGGCAGGTTTGACTGTTTCTCTGCACCAGACATCTGTAGTAA
TCTGTATGTTTTTTCAGCCGTCTCTAGCAGTATTTAAAGGACAAGGAACCAAAGAATATGA
AATTCATCATGGAAAGAAGATTCTATATGATATACTTGCCTTTGCCAAAGAAAGTGTGAA
TTCTCATGTTACCACGCTTGGACCTCAAAATTTTCTGCCAATGACAAAGAACCATGGCT
TGTTGATTTCTTTGCCCCCTGGTGTCCACCATGTGAGCTTTACTACCAGAGTTACGAAG
AGCATCAAATCTTCTTTATGGTCAGCTTAAAGTTTGGTACACTAGATTGTACAGTTCATGA
GGGACTCTGTAAACATGTATAACATTCAAGGCTTATCCAACAACAGTGGTATTCAACCAGTC
CAACATTCATGAGTATGAAGGACATCACTCTGCTGAACAAATCTTGGAGTTCATAGAGGA
TCTTATGAATCCTTCAGTGGTCTCCCTTACACCCACCACCTTCAACGAACCTAGTTACACA
AAGAAAACACAACGAAGTCTGGATGGTTGATTCTATTCTCCGTGGTGTATCCTTGCCA
AGTCTTAATGCCAGAATGGAAAAGAATGGCCCGGACATTAAGTGGACTGATCAACGTGGG
CAGTATAGATTGCCAACAGTATCATTTCTTTTGTGCCAGGAAAACGTTCAAAGATACCC
TGAGATAAGATTTTTTCCCCCAAATCAAATAAAGCTTATCAGTATCACAGTTACAATGG
TTGGAATAGGGATGCTTATTCCTGAGAATCTGGGGTCTAGGATTTTTTACCTCAAGTATC
CACAGATCTAACACCTCAGACTTTTCAAGTGAAGAAAGTTCTACAAGGGAAAAATCATTGGGT
GATTGATTTCTATGCTCCTTGGTGTGGACCTTGCCAGAATTTTGCTCCAGAATTTGAGCT
CTTGGCTAGGATGATTAAAGGAAAAGTGAAGCTGGAAAAGTAGACTGTCAGGCTTATGC
TCAGACATGCCAGAAAGCTGGGATCAGGGCTATCCAAGTGAAGTTTATTTTCTACGA
AAGAGCAAAGAGAAATTTTCAAGAAGAGCAGATAAATACCAGAGATGCAAAGCAATCGC
TGCCTTAATAAGTGAAAATTTGAAAATCTCCGAAATCAAGGCAAGAGGAATAAGGATGA
ACTTTGATAATGTTGAAGATGAAGAAAAAGTTTAAAAAGAAATCTGACAGATGACATCAG
AAGACACCTATTTAGAATGTTACATTTATGATGGGAATGAATGAACATTATCTTAGACTT

119/392

GCAGTTGTTACTGCCAGAATTATCTACAGCACTGGTGTAAAAGAAGGGTCTGCAAACCTTTT
TCTGTAAAGGGCCGGTTTATAAATATTTTAGACTTTGCAGGCTATAATATATGGTTCACA
CATGAGAACAAGAATAGAGTCATCATGTATTCTTTGTTATTTGCTTTTAAACAACCTTTAA
AAAATATTTAAACGATTCTTAGCTCAGAGCCATACAAAAGTAGGCTGGATTCAGTCCATG
GACCATAGATTGCTGTCCCCCTCGACGGACTTATAATGTTTCAGGTGGCTGGCTTGAACA
TGAGTCTGCTGTGCTATCTACATAAATGTCTAAGTTGTATAAAGTCCACTTTCCCTTCAC
GTTTTTTGGCTGACCTGAAAAGAGGTAACTTAGTTTTTGGTCACTTGTTCTCCTAAAAAT
GCTATCCCTAACCATATATTTATATTTTCGTTTTAAAAACACCCATGATGTGGCACAGTAA
ACAAACCCTGTTATGCTGTATTATTATGAGGAGATTCTTCATTGTTTTCTTCCCTTCTCA
AAGGTTGAAAAAATGCTTTTAAATTTTTTACAGCCGAGAAACAGTGCAGCAGTATATGTGC
ACACAGTAAGTACACAAATTTGAGCAACAGTAAGTGCACAAATTCTGTAGTTTGCTGTAT
CATCCAGGAAAACCTGAGGGAAAAAAATTATAGCAATTAAGTGGGCATTGTAGAGTATCC
TAAATATGTTATCAAGTATTTAGAGTTCTATATTTTAAAGATATATGTGTTTCATGTATTT
TCTGAAATTGCTTTCATAGAAAATTTTCCCACTGATAGTTGATTTTTGAGGCATCTAATAT
TTACATATTTGCCTTCTGAACTTTGTTTTGACCTGTATCCTTTATTTACATTGGGTTTTT
CTTTCATAGTTTTTGGTTTTTCACTCCTGTCCAGTCTATTTATTATTCAAATAGGAAAAAT
TACTTTACAGGTTGTTTTACTGTAGCTTATAATGATACTGTAGTTATTCCAGTTACTAGT
TTACTGTCAGAGGGCTGCCTTTTTTTCAGATAAATATTGACATAATAACTGAAGTTATTTTT
ATAAGAAAATCAAGTATATAAATCTAGGAAAGGGATCTTCTAGTTTCTGTGTTGTTTAGA
CTCAAAGAATCACAAATTTGTCAGTAACATGTAGTTGTTTAGTTATAATTTCAGAGTGTAC
AGAATGGTAAAAATTCCAATCAGTCAAAAGAGGTCAATGAATTAAAAGGCTTGCAACTTT
TTCAAAAAAAAAAAAAAAAAA

120/392

FIGURE 114

MGVWLNKDDYIRD LKRIILCFLIVYMAILVGT DQDFYSL LGVSKTASSREIRQAFKKLAL
KLHPDKNPNNPNAHGDF LKINRAYEVLKDEDLRKKYDKYGEK GLEDNQGGQYESWNYRY
DFGIYDDDPEIITLERREFDA AVNSGELW FVNFYSPGCSHCHDLAPTWRDFAKEVDGLLR
IGAVNCGDDRMLCRMKG VNSYPSLFIFRSGMAPVKYHGDRSKESLV SFAMQHVRSTVTEL
WTGNFVNSIQTAFAAGIGW LITFCSKGGDCLTSQTRLRLSGMLF LNSLDAKEIYLEVIHN
LPDFELLSANTLEDRLA HHRWLLFFHFGKNENSNDPELKKLKTLLKNDHIQVGRFDCSSA
PDICSNLYVFQPSLAVFKG QGTKEYEIHGKKILYDILAFAKESVNSHVTTLGPQNFPAN
DKEPWLVDFFAPWCPPCR ALLPELRRASNLLYGQLKFGTLDCTVHEGLCNMYNIQAYPTT
VVFNQSNIEHEYEGHSAEQ ILEFIEDLMNPSVVSLTPTTFNELVTQRKHNEVWMVDFYSP
WCHPCQVLMPEWKRMARTLTGLIN VGSIDCQQYHSFCAQENVQRYPEIRFFPPKSNKAYQ
YHSYNGWNRDAYSLRIWGLG FL PQVSTD LTPQTFSEKVLQGKNHWVIDFYAPWCGPCQNF
APEFELLARMIKGVKAGKVDCQAYA QTCQKAGIRAYPTVKFYFYERAKRNFQEEQINTR
DAKAIAALISEKLETLRNQ GKRNKDEL

Important features:

Endoplasmic reticulum targeting sequence:

amino acids 744-747

Cytochrome c family heme-binding site signature:

amino acids 158-163

Nt-dnaJ domain signature:

amino acids 77-96

N-glycosylation site:

amino acids 484-487

121/392

FIGURE 115

GCGGGCTGTTGACGGCGCTGCGATGGCTGCCTGCGAGGGCAGGAGAAGCGGAGCTCTCGG
TTCCTCTCAGTCGGACTTCCTGACGCCGCCAGTGGGCGGGGCCCCCTTGGGCCGTCGCCAC
CACTGTAGTCATGTACCCACCGCCGCCGCCGCCCTCATCGGGACTTCATCTCGGTGAC
GCTGAGCTTTGGCGAGAGCTATGACAACAGCAAGAGTTGGCGGGCGGCGCTCGTGCTGGAG
GAAATGGAAGCAACTGTCGAGATTGCAGCGGAATATGATTCTCTTCCTCCTTGCCCTTTCT
GCTTTTCTGTGGACTCCTCTTCTACATCAACTTGGCTGACCATTGGAAAGCTCTGGCTTT
CAGGCTAGAGGAAGAGCAGAAGATGAGGCCAGAAATTGCTGGGTTAAAACCAGCAAATCC
ACCCGTCTTACCAGCTCCTCAGAAGGCGGACACCGACCCTGAGAACTTACCTGAGATTTCT
GTCACAGAAGACACAAAGACACATCCAGCGGGGACCACCTCACCTGCAGATTAGACCCCC
AAGCCAAGACCTGAAGGATGGGACCCAGGAGGAGGCCACAAAAGGCAAGAAGCCCCCTGT
GGATCCCCGCCCGGAAGGAGATCCGCAGAGGACAGTCATCAGCTGGAGGGGAGCGGTGAT
CGAGCCTGAGCAGGGCACCGAGCTCCCTTCAAGAAGAGCAGAAGTGCCACCAAGCCTCC
CCTGCCACCGGCCAGGACACAGGGCACACCAAGTGCATCTGAACATATCGCCAGAAGGGCGT
GATTGACGTCTTCCTGCATGCATGGAAGGATAACCGCAAGTTTGCATGGGGCCATGACGA
GCTGAAGCCTGTGTCCAGGTCTTCAGTGAGTGGTTTGGCCTCGGTCTCACACTGATCGA
CGCGCTGGACACCATGTGGATCTTGGGTCTGAGGAAAGAATTTGAGGAAGCCAGGAAGTG
GGTGTGAAGAAGTTACACTTTGAAAAGGACGTGGACGTCAACCTGTTTGAGAGCACGAT
CCGCATCCTGGGGGGGCTCCTGAGTGCCTACCACCTGTCTGGGGACAGCCTCTTCCTGAG
GAAAGCTGAGGATTTTGGAAATCGGCTAATGCCTGCCTTCAGAACACCATCCAAGATTCC
TTACTCGGATGTGAACATCGGTACTGGAGTTGCCCCACCGCCACGGTGGACCTCCGACAG
CACTGTGGCCGAGGTGACCAGCATTAGCTGGAGTTCCGGGAGCTCTCCCGTCTCACAGG
GGATAAGAAGTTTCAGGAGGCAGTGGAGAAGGTGACACAGCACATCCACGGCCTGTCTGG
GAAGAAGGATGGGCTGGTGCCCATGTTTATCAATACCCACAGTGGCCTCTTACCCACCT
GGGCGTATTACGCTGGGCGCCAGGGCCGACAGCTACTATGAGTACCTGCTGAAGCAGTG
GATCCAGGGCGGGAAGCAGGAGACACAGCTGCTGGAAGACTACGTGGAAGCCATCGAGGG
TGTCAGAACGCACCTGCTGCGGCACTCCGAGCCCAGTAAGCTCACCTTTGTGGGGGAGCT
TGCCACAGGCCGCTTCAGTGCCAAGATGGACCACCTGGTGTGCTTCTGCCAGGGACGCT
GGCTCTGGGCGTCTACCACGGCCTGCCCGCCAGCCACATGGAGCTGGCCAGGAGCTCAT
GGAGACTTGTACCAGATGAACCGGCAGATGGAGACGGGGCTGAGTCCCGAGATCGTGCA
CTTCAACCTTTACCCCCAGCCGGGCCGTCGGGACGTGGAGGTCAAGCCAGCAGACAGGCA
CAACCTGCTGCGGCCAGAGACCGTGGAGAGCCTGTTCTACCTGTACCGCGTCACAGGGGA
CCGCAAATACCAGGACTGGGGCTGGGAGATTCTGCAGAGCTTCAGCCGATTACACGGGT
CCCCCTCGGGTGGCTATTCTTCCATCAACAATGTCCAGGATCCTCAGAAGCCCGAGCCTAG
GGACAAGATGGAGAGCTTCTTCTGGGGGAGACGCTCAAGTATCTGTTCTTGCTCTTCTC
CGATGACCCAAACCTGCTCAGCCTGGACGCCTACGTGTTCAACACCGAAGCCCACCCTCT
GCCTATCTGGACCCCTGCCTAGGGTGGATGGCTGCTGGTGTGGGGACTTCGGGTGGGCAG
AGGCACCTTGCTGGGTCTGTGGCATTTTCCAAGGGCCACGTCAGCACCGGCAACCGCCAA
GTGCCCAGGCTCTGAACCTGGCTCTGGGCTCCTCCTCGTCTCTGCTTTAATCAGGACACC
GTGAGGACAAGTGAGGCCGTGAGTCTTGGTGTGATGCGGGGTGGGCTGGGCCGCTGGAGC
CTCCGCTGCTTCTCCAGAAGACACGAATCATGACTCACGATTGCTGAAGCCTGAGCAG
GTCTCTGTGGGCCGACCAGAGGGGGGCTTCGAGGTGGTCCCTGGTACTGGGGTGACCGAG
TGGACAGCCCAGGGTGCAGCTCTGCCCGGGCTCGTGAAGCCTCAGATGTCCCAATCCAA
GGGTCTGGAGGGGCTGCCGTGACTCCAGAGGCCTGAGGCTCCAGGGCTGGCTCTGGTGT
TACAAGCTGGACTCAGGGATCCTCCTGGCCGCCCCGAGGGGGCTTGGAGGGCTGGACGG
CAAGTCCGTCTAGCTCACGGGCCCTCCAGTGGAAATGGGTCTTTTCGGTGGAGATAAAAG
TTGATTTGCTCTAACCGCAA

122/392

FIGURE 116

MAACEGRRSGALGSSQSDFLTPPVGGAPWAVATTVMYPPPPPPPHRDFISVTLSFGESY
DNSKSWRRRSCWRKWKQLSRLQRNMILFLLAFLLLFCGLLFYINLADHWKALAFRLEEEQK
MRPEIAGLKPANPPVLPAPQKADTDPENLPEISSQKTQRHIQRGPPHLQIRPPSQDLKDG
TQEEATKRQEAPVDPRPEGDPQRTVISWRGAVIEPEQGTELPSSRAEVPTKPPLPPARTQ
GTPVHLNRYRQKGVIDVFLHAWKGYRKFAWGHDELKPVSRSFSEWFGLGLTLIDALDTMWI
LGLRKEFEEARKWVSKKLHFEDVDVNLFESTITIRILGGLLSAYHLSGDSLFRLKAEDFGN
RLMPAFRTPSKIPYSDVNIGTGVAHPPRWTS DSTVAEVT SIQLEFRELSRLTGDKKFQEA
VEKVTQHIHGLSGKKDGLVPMFINTHSGLFTHLGVFTLGARADSYEYLLKQWIQGGKQE
TQLLEDYVEAIEGVRTHLLRHSEPSKLT FVGELAHGRFSAKMDHLVCFLPGTLALGVYHG
LPASHMELAQELMETCYQMNROMETGLSPEIVHFNLYPQGRRDVEVKPADRHNLRLPET
VESLFYLYRVTGDRKYQDWGWEILQSFSRFRTRVPSGGYSSINNVQDPQKPEPRDKMESFF
LGETLKYLFLLFSDDPNLLSLDAYVFNTTEAHPLPIWTPA

Important features of the protein:

Transmembrane domain:

amino acids 21-40 and 84-105 (type II)

123/392

FIGURE 117

GTGGGATTTATTTGAGTGCAAGATCGTTTTCTCAGTGGTGGTGGAAAGTTGCCTCATCGCA
GGCAGATGTTGGGGCTTTGTCCGAACAGCTCCCCCTCTGCCAGCTTCTGTAGATAAGGGTT
AAAACTAATATTTATATGACAGAAGAAAAAGATGTCATTCCGTAAAGTAAACATCATCA
TCTTGGTCCTGGCTGTTGCTCTCTTCTTACTGGTTTTTGCACCATAACTTCCTCAGCTTGA
GCAGTTTGTAAAGGAATGAGGTTACAGATTCAGGAATTGTAGGGCCTCAACCTATAGACT
TTGTCCCAAATGCTCTCCGACATGCAGTAGATGGGAGACAAGAGGAGATTCTGTGGTCA
TCGCTGCATCTGAAGACAGGCTTGGGGGGGCCATTGCAGCTATAAACAGCATTCAGCACA
ACACTCGCTCCAATGTGATTTTCTACATTGTTACTCTCAACAATACAGCAGACCATCTCC
GGTCCTGGCTCAACAGTGATTCCCTGAAAAGCATCAGATACAAAATTGTCAATTTTGACC
CTAAACTTTTGGAAGGAAAAGTAAAGGAGGATCTGACCAGGGGGAATCCATGAAACCTT
TAACCTTTGCAAGGTTCTACTTGCCAATTCTGGTTCCCAGCGCAAAGAAGGCCATATACA
TGGATGATGATGTAATTGTGCAAGGTGATATCTTGCCCTTTACAATACAGCACTGAAGC
CAGGACATGCAGCTGCATTTTCAGAAGATTGTGATTGAGCCTCTACTAAAGTTGTCATCC
GTGGAGCAGGAAACCAGTACAATTACATTGGCTATCTTGACTATAAAAAGGAAAGAATTC
GTAAGCTTTCCATGAAAGCCAGCACTTGCTCATTTAATCCTGGAGTTTTTGTGCAAACC
TGACGGAATGGAAACGACAGAATATAACTAACCAACTGGAAAAATGGATGAAACTCAATG
TAGAAGAGGGACTGTATAGCAGAACCTGGCTGGTAGCATCACAAACACCTCCTCTGCTTA
TCGTATTTTATCAACAGCACTCTACCATCGATCCTATGTGGAATGTCCGCCACCTTGTT
CCAGTGCTGGAAAACGATATTCACCTCAGTTTGTAAAGGCTGCCAAGTTACTCCATTGGA
ATGGACATTTGAAGCCATGGGGAAGGACTGCTTCATATACTGATGTTTGGGAAAAATGGT
ATATTCCAGACCCAACAGGCAAATTC AACCTAATCCGAAGATATAACCGAGATCTCAAACA
TAAAGTGAAACAGAATTTGAACTGTAAGCAAGCATTTCTCAGGAAGTCCTGGAAGATAGC
ATGCATGGGAAGTAACAGTTGCTAGGCTTCAATGCCTATCGGTAGCAAGCCATGGAAAAA
GATGTGTCAGCTAGGTAAAGATGACAAACTGCCCTGTCTGGCAGTCAGCTTCCCAGACAG
ACTATAGACTATAAATATGTCTCCATCTGCCTTACCAAGTGTTTTCTTACTACAATGCTG
AATGACTGGAAAGAAGAACTGATATGGCTAGTTGAGCTAGCTGGTACAGATAATTCAAAA
CTGCTGTTGGTTTTAATTTTGTAACCTGTGGCCTGATCTGTAAATAAACTTACATTTTTC

124/392

FIGURE 118

MSFRKVNIIILVLAVALFLLVLHHNFLSLSSLLRNEVTD SGIVGPQPIDFVPNALRHAVD
GRQEEIPVVIAASEDRLGGAIAAINSIQHNTRSNVIFYIVTLNNTADHLRSWLNSDSLKS
IRYKIVNFDPKLLEGKVKEPDQGESMKPLTFARFYLPILVPSAKKAIYMDDDVIVQGDI
LALYNTALKPGHAAAFSEDCDSASTKV VIRGAGNQYNYIGYLDYKKERIRKLSMKASTCS
FNPGVFVANLTEWKRQNI TNQLEKWMKLNVEEGLYSRTL AGSITTPPLLIVFYQQHSTID
PMWNVRLGSSAGKRYSPQFVKA AKLLHWNGHLKPWGRTASYTDVWEKWIIPDPTGKFNL
IRRYTEISNIK

125/392

FIGURE 119

CCATCCCTGAGATCTTTTTATAAAAAACCCAGTCTTTGCTGACCAGACAAAGCATACCAG
ATCTCACCAGAGAGTCGCAGACACTATGCTGCCTCCCATGGCCCTGCCCAGTGTGTCCTG
GATGCTGCTTTCCTGCCTCATTCTCCTGTGTCAGGTTCAAGGTGAAGAAACCCAGAAGGA
ACTGCCCTCTCCACGGATCAGCTGTCCCAAAGGCTCCAAGGCCTATGGCTCCCCCTGCTA
TGCCTTGTTTTTGTACCAAAAATCCTGGATGGATGCAGATCTGGCTTGCCAGAAGCGGCC
CTCTGGAAACTGGTGTCTGTGCTCAGTGGGGCTGAGGGATCCTTCGTGTCCTCCCTGGT
GAGGAGCATTAGTAACAGCTACTCATACATCTGGATTGGGCTCCATGACCCACACAGGG
CTCTGAGCCTGATGGAGATGGATGGGAGTGGAGTAGCACTGATGTGATGAATTACTTTGC
ATGGGAGAAAAATCCCTCCACCATCTTAAACCCTGGCCACTGTGGGAGCCTGTCAAGAAG
CACAGGATTTCTGAAGTGGAAGATTATAACTGTGATGCAAAGTTACCCTATGTCTGCAA
GTTCAAGGACTAGGGCAGGTGGGAAGTCAGCAGCCTCAGCTTGGCGTGCAGCTCATCATG
GACATGAGACCAGTGTGAAGACTCACCTGGAAGAGAATATTCTCCCCAACTGCCCTAC
CTGACTACCTTGTCATGATCCTCCTTCTTTTCTTTTCTTCACCTTCATTTCAGGCTT
TTCTCTGTCTTCCATGTCTTGAGATCTCAGAGAATAATAATAAAATGTTACTTTATAAA
AAAAAAAAAAAAAAAAAAAA

126/392

FIGURE 120

MLPPMALPSVSWMLLSCLILLCQVQGEETQKELPSPRISCPKGSKAYGSPCYALFLSPKS
WMDADLACQKRPSGKLVSVLGAEGSFVSSLVRSISNSYSYIWIGLHDPTQGSEPDGDGW
EWSSTDVMNYFAWEKNPSTILNPGHCGSLSRSTGFLKWKDYNCDAKLPYVCKFKD

Important features:

Signal peptide:

amino acids 1-26

C-type lectin domain signature:

amino acids 146-171

127/392

FIGURE 121

AAAGTTACATTTTCTCTGGAACCTCTCCTAGGCCACTCCCTGCTGATGCAACATCTGGGTT
TGGGCAGAAAGGAGGGTGCTTCGGAGCCCCGCCCTTTCTGAGCTTCCTGGGCCGGCTCTAG
ACAATTCAGGCTTCGCTGCGACTCAGACCTCAGCTCCAACATATGCATTCTGAAGAAAG
ATGGCTGAGATGGACAGAATGCTTTATTTTGGAAAGAAACAATGTTCTAGGTCAAACCTGA
GTCTACCAAATGCAGACTTTTACAATGGTTCTAGAAGAAATCTGGACAAGTCTTTTCATG
TGGTTTTTCTACGCATTGATTCCATGTTTGCTCACAGATGAAGTGGCCATTCTGCCTGCC
CCTCAGAACCTCTCTGTACTCTCAACCAACATGAAGCATCTCTTGATGTGGAGCCCAGTG
ATCGCGCCTGGAGAAACAGTGTACTATTCTGTCTGAATACCAGGGGGAGTACGAGAGCCTG
TACACGAGCCACATCTGGATCCCCAGCAGCTGGTGCTCACTCACTGAAGGTCTGAGTGT
GATGTCACTGATGACATCACGGCCACTGTGCCATACAACCTTCGTGTCAGGGCCACATTG
GGCTCACAGACCTCAGCCTGGAGCATCCTGAAGCATCCCTTTAATAGAAACTCAACCATC
CTTACCCGACCTGGGATGGAGATCACCAAAGATGGCTTCCACCTGGTTATTGAGCTGGAG
GACCTGGGGCCCCAGTTTGAGTTCCTTGTGGCCTACTGGAGGAGGGAGCCTGGTGCCGAG
GAACATGTCAAAATGGTGAGGAGTGGGGGTATTCCAGTGCACCTAGAAACCATGGAGCCA
GGGGCTGCATACTGTGTGAAGGCCAGACATTCGTGAAGGCCATTGGGAGGTACAGCGCC
TTCAGCCAGACAGAATGTGTGGAGGTGCAAGGAGAGGCCATTCCCCTGGTACTGGCCCTG
TTTGCCCTTTGTTGGCTTCATGCTGATCCTTGTGGTTCGTGCCACTGTTTCGTCTGAAAAATG
GGCCGGCTGCTCCAGTACTCCTGTTGCCCGTGGTGGTCTCCAGACACCTTGAAAATA
ACCAATTCACCCAGAAAGTTAATCAGCTGCAGAAGGGAGGAGGTGGATGCCTGTGCCACG
GCTGTGATGTCTCCTGAGGAACCTCCTCAGGGCCTGGATCTCATAGGTTTGCGGAAGGGCC
CAGGTGAAGCCGAGAACCTGGTCTGCATGACATGGAAACCATGAGGGGACAAGTTGTGTT
TCTGTTTTCCGCCACGGACAAGGGATGAGAGAAGTAGGAAGAGCCTGTTGTCTACAAGTC
TAGAAGCAACCATCAGAGGCAGGGTGGTTTGTCTAACAGAACTGACTGAGGCTTAGGG
GATGTGACCTCTAGACTGGGGGCTGCCACTTGCTGGCTGAGCAACCCTGGGAAAAGTGAC
TTCATCCCTTCGGTCCTAAGTTTTCTCATCTGTAATGGGGGAATTACCTACACACCTGCT
AAACACACACACACAGAGTCTCTCTATATATACACACGTACACATAAATACACCCAGC
ACTTGCAAGGCTAGAGGGAACTGGTGACACTCTACAGTCTGACTGATTCAGTGTTTTCTG
GAGAGCAGGACATAAATGTATGATGAGAATGATCAAGGACTCTACACACTGGGTGGCTTG
GAGAGCCCACTTTCCAGAAATAATCCTTGAGAGAAAAGGAATCATGGGAGCAATGGTGT
GAGTTCACCTCAAGCCCAATGCCGGTGCAGAGGGGAATGGCTTAGCGAGCTCTACAGTAG
GTGACCTGGAGGAAGGTACAGCCCACTGAAAATGGGATGTGCATGAACACGGAGGATC
CATGAACTACTGTAAAGTGTGACAGTGTGTGCACACTGCAGACAGCAGGTGAAATGTAT
GTGTGCAATGCGACGAGAATGCAGAAGTCAGTAACATGTGCATGTTTGTGTGCTCCTTT
TTTCTGTTGGTAAAGTACAGAATTCAGCAAATAAAAAGGGCCACCCTGGCCAAAAGCGGT
AAAAAAAAAAAAAAAA

128/392

FIGURE 122

MQTFTMVLEEIWTSLFMWFFYALIPCLLTDEVAILPAPQNLSVLSTNMKHLMLWSPVIAP
GETVYYSVEYQGEYESLYTSHIWIPSSWCSLTEGPECDVTDDITATVPYNLRVRATLGSQ
TSAWSILKHFPNRRNSTILTRPGMEITKDG FHLVIELEDLGPQFEFLVAYWRREPGEAEHV
KMVRSGGIPVHLETMEPGAAYCVKAQTFVKAIGRYSAFSQTECVEVQGEAIPLVLALFAF
VGFMLILVVVPLFVWKMGRLQLYSCCPVVVLPDTLKITNSPQKLISCRREEVDACATAVM
SPEELLRAWIS

Important features:**Signal peptide:**

amino acids 1-29

Transmembrane domain:

amino acids 230-255

N-glycosylation sites:

amino acids 40-43 and 134-137

Tissue factor proteins homology:

amino acids 92-119

Integrins alpha chain protein homology:

amino acids 232-262

129/392

FIGURE 123

CGGACGCGTGGGCGGCCACCTCCGGAACAAGCCATGGTGGCGGCGACGGTGGCAGCGGCG
TGGCTGCTCCTGTGGGCTGCGGCCTGCGCGCAGCAGGAGCAGGACTTCTACGACTTCAAG
GCGGTCAACATCCGGGGCAAACCTGGTGTGCTGGAGAAGTACCGCGGATCGGTGTCCCTG
GTGGTGAATGTGGCCAGCGAGTGGGCTTACAGACCAGCACTACCGAGCCCTGCAGCAG
CTGCAGCGAGACCTGGGCCCCCACCCTTTAACGTGCTCGCCTTCCCCTGCAACCAGTTT
GGCCAACAGGAGCCTGACAGCAACAAGGAGATTGAGAGCTTTGCCCCCGCACCTACAGT
GTCTCATTCCCCATGTTTAGCAAGATTGCAGTCACCGGTACTGGTGCCCATCCTGCCTTC
AAGTACCTGGCCCAGACTTCTGGGAAGGAGCCACCTGGAACCTTCTGGAAGTACCTAGTA
GCCCCAGATGGAAAGGTGGTAGGGGCTTGGGACCCAACCTGTGTCAGTGGAGGAGGTCAGA
CCCCAGATCACAGCGCTCGTGAGGAAGCTCATCCTACTGAAGCGAGAAGACTTATAACCA
CCGCGTCTCCTCCTCCACCACCTCATCCCGCCACCTGTGTGGGGCTGACCAATGCAAAC
TCAAATGGTGTCTTCAAAGGGAGAGACCCACTGACTCTCCTTCCTTTACTCTTATGCCATT
GGTCCCATCATTCTTGTGGGGGAAAAATTCTAGTATTTTGATTATTTGAATCTTACAGCA
ACAAATAGGAACCTCTGGCCAATGAGAGCTCTTGACCAGTGAATCACCAGCCGATACGAA
CGTCTTGCCAACAAAAATGTGTGGCAAATAGAAGTATATCAAGCAATAATCTCCACCCA
AGGCTTCTGTAAACTGGGACCAATGATTACCTCATAGGGCTGTTGTGAGGATTAGGATGA
AATACCTGTGAAAGTGCCTAGGCAGTGCCAGCCAAATAGGAGGCATTCAATGAACATTTT
TTGCATATAAACCAAAAAATAACTTGTTATCAATAAAAACTTGCATCCAACATGAATTTCT
CAGCCGATGATAATCCAGGCCAAAGGTTTAGTTGTTGTTATTTTCTCTGTATTATTTTCT
TCATTACAAAAGAAATGCAAGTTCATTGTAACAATCCAAACAATACCTCACGATATAAAA
TAAAAATGAAAGTATCCTCCTCAAAAA

130/392

FIGURE 124

MVAATVAAAWLLLWAAACAQQEQDFYDFKAVNIRGKLVSLVKYRGSVSLVNVASECGFT
DQHYRALQQLQORDLGPHHFNVLAFFPCNQFGQQEPDSNKEIESFARRTYSVSFPMFSKIAV
TGTGAHPAFKYLAQTSGKEPTWNFWKYLVA PDGKVVGAWDPTVSVEEVRPQITALVRKLI
LLKREDL

131/392

FIGURE 125

CGGACGCGTGGGCGGACGCGTGGGCGGACGCGTGGGTTGGGAGGGGGCAGGATGGGAGGG
AAAGTGAAGAAAACAGAAAAGGAGAGGGACAGAGGCCAGAGGACTTCTCATACTGGACAG
AAACCGATCAGGCATGGAACTCCCCCTTCGTCACTCACCTGTTCTTGCCCCCTGGTGTTCTT
GACAGGTCTCTGCTCCCCCTTTAACCTGGATGAACATCACCCACGCCTATTCCCAGGGCC
ACCAGAAAGCTGAATTTGGATACAGTGTCTTACAACATGTTGGGGGTGGACAGCGATGGAT
GCTGGTGGGCGCCCCCTGGGATGGGCCTTCAGGCGACCGGAGGGGGGACGTTTATCGCTG
CCCTGTAGGGGGGGGCCACAATGCCCCATGTGCCAAGGGCCACTTAGGTGACTACCAACT
GGGAAATTCATCTCATCCTGCTGTGAATATGCACCTGGGGATGTCTCTGTTAGAGACAGA
TGGTGATGGGGGATTTCATGGTGAGCTTAAGGAGAGGGTGGTGGCAGTGTCTCTGAAGGTCC
ATAAAAGAAAAAAGAGAAGTGTGGTAAAGGAAAATGGTCTGTGTGGAGGGGTCAAGGAGT
TAAAAACCCTAGAAAGCAAAAGGTAGGTAATGTCAGGGAGTAGTCTTCATGCCTCCTTCA
ACTGGGAGCATGTTCTGAGGGTGCCCTCCCAAGCCTGGGAGTAACATATTTCCCCCATCCC
CAGGCCTGTGCCCCCTCTCTGGTCTCGTGCTTGTGGCAGCTCTGTCTTCAGTTCTGGGATA
TGTGCCCCGTGTGGATGCTTCATTCCAGCCTCAGGGAAGCCTGGCACCCACTGCCCAACGT
GAGCCAGAGGAAGGCTGAGTACTTGGTTCCAGAAGGAGATACTGGGTGGGAAAAAGATG
GGCAAAGCGGTATGATGCCTGGCAAAGGGCCTGCATGGCTATCCTCATTGCTACCTAAT
GTGCTTGCAAAGCTCCATGTTTCCTAACAGATTGAGACTCCTGGCCAGGTGTGGTGGCC
CACACCTGTAATTCTAGCACTTTGGGAGGCCAAGGTGGGCAGATCACTTGAGGTCAGGAG
TTCAAGACCAGCCTGGCCAACATGGTGAACTCCATCTCTACTAAAAAATAATAACA
AAAATTAGCTGGGTGCGCTAGTGCATGCCTGTAATCTCATCTACTCGGGAGGCTAAGACA
GGAGACTCTCACTTCAACCCAGGAGGTGGAGGTTGCGGTGAGCCAAGATTGTGCCTCTGC
ACTCTAGCGTGGGTGACAGAGTAAGCGAGACTCCATCTCAAAAATAATAATAATAAT
TCAGACTCCTTATCAGGAGTCCATGATCTGGCCTGGCACAGTAACTCATGCCTGTAATCC
CAACATTTTGGGAGGCCAACGCAGGAGGATTGCTTGAGGTCTGGAGGTTTGAGACCAGCC
TGGGCAACATAGAAAGACCCCATCTCTAAATAAATGTTTAAAAAT

132/392

FIGURE 126

MELPFVTHLFLPLVFLTGLCSPFNLDEHHPRLFPGPPEAEFGYSVLQHVGGGQRWMLVGA
PWDGPSGDRRGDVYRCPVGGAHNAPCAKGHLGDYQLGNSSHPAVNMHLGMSLLETGDG
FMVS

Important features:

Signal peptide:

amino acids 1-22

Cell attachment sequence:

amino acids 70-73

N-glycosylation site:

amino acids 98-101

Integrins alpha chain proteins:

amino acids 67-81

133/392

FIGURE 127

GAGAGGACGAGGTGCCGCTGCCTGGAGAATCCTCCGCTGCCGTCGGCTCCCGGAGCCCAG
CCCTTTCCTAACCCAACCCAACCTAGCCCAGTCCCAGCCGCCAGCGCCTGTCCCTGTAC
GGACCCCAGCGTTACCAATGCATCCTGCCGTCTTCCTATCCTTACCCGACCTCAGATGCTC
CCTTCTGCTCCTGGTAACTTGGGTTTTTACTCCTGTAACAACCTGAAATAACAAGTCTTGC
TACAGAGAATATAGATGAAATTTTAAACAATGCTGATGTTGCTTTAGTAAATTTTATGC
TGACTGGTGTTCGTTTCAGTCAGATGTTGCATCCAATTTTGTAGGAAGCTTCCGATGTCAT
TAAGGAAGAATTTCCAAATGAAAATCAAGTAGTGTGTTGCCAGAGTTGATTGTGATCAGCA
CTCTGACATAGCCCAGAGATACAGGATAAGCAAATACCCAACCCCTCAAATTGTTTCGTAA
TGGGATGATGATGAAGAGAGAATACAGGGGTCAGCGATCAGTGAAAGCATTGGCAGATTA
CATCAGGCAACAAAAAAGTGACCCCATTCAGAAATTCGGGACTTAGCAGAAATCACCAC
TCTTGATCGCAGCAAAAGAAATATCATTGGATATTTTGAGCAAAAGGACTCGGACAACCTA
TAGAGTTTTTGAACGAGTAGCGAATATTTTGCATGATGACTGTGCCTTTCTTTCTGCATT
TGGGGATGTTTCAAAACCGGAAAGATATAGTGGCGACAACATAATCTACAAACCACCAGG
GCATTCTGCTCCGGATATGGTGTACTTGGGAGCTATGACAAATTTTGATGTGACTTACAA
TTGGATTCAAGATAAATGTGTTCTCTTGTCCGAGAAATAACATTTGAAAATGGAGAGGA
ATTGACAGAAGAAGGACTGCCTTTTCTCATACTCTTTCACATGAAAGAAGATACAGAAAG
TTTAGAAATATTCCAGAATGAAGTAGCTCGGCAATTAATAAGTGAAAAAGGTACAATAAA
CTTTTTACATGCCGATTGTGACAAATTTAGACATCCTCTTCTGCACATACAGAAAACCTCC
AGCAGATTGTCTGTAAATCGCTATTGACAGCTTTAGGCATATGTATGTGTTTGGAGACTT
CAAAGATGTATTAATTCCTGGAAAACTCAAGCAATTCGTATTTGACTTACATTCTGGAAA
ACTGCACAGAGAATTCCATCATGGACCTGACCCAACTGATACAGCCCCAGGAGAGCAAGC
CCAAGATGTAGCAAGCAGTCCACCTGAGAGCTCCTTCCAGAACTAGCACCCAGTGAATA
TAGGTATACTCTATTGAGGGATCGAGATGAGCTTTTAAAAACTTGAAAAACAGTTTGTAAG
CCTTTCAACAGCAGCATCAACCTACGTGGTGGAATAGTAAACCTATATTTTTCATAATTC
TATGTGTATTTTTTATTTTGAATAAACAGAAAGAAATTTAAAAA
AAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAA

134/392

FIGURE 128

MHPAVFLSLPDLRCSLLLLVTWVFTPVTTEITSLATENIDEILNNADVALVNFYADWCRF
SQMLHPIFEEASDVIKEEFPNENQVVFARVDCDQHS DIAQRYRISKYPTLKLFRNGMMMK
REYRGQRSVKALADYIRQQKSDPIQEIRDLAEITTLDRSKRNIIGYFEQKDS DN YRVFER
VANILHDDCAFLSAFGDVSKPERYSGDNIIYKPPGHSAPDMVYLGAMTNFDVTYNWIQDK
CVPLVREITFENGEEELTEEGLPFLILFHMKEDTESLEIFQNEVARQLISEKGTINFLHAD
CDKFRHPLLHIQKTPADCPVIAIDSFRHMYVFGDFKDVLIPGKLKQFVFDLHSGKLHREF
HHGPDPTDTAPGEQAQDVASSPPESSFQKLAPSEYRYTLLRDRDEL

Important features:

Signal peptide:

amino acids 1-29

Endoplasmic reticulum targeting sequence:

amino acids 403-406

Tyrosine kinase phosphorylation site:

amino acids 203-211

Thioredoxin family proteins:

amino acids 50-66

135/392

FIGURE 129

GAGCAGGACGGAGCCATGGACCCCGCCAGGAAAGCAGGTGCCCAGGCCATGATCTGGACT
GCAGGCTGGCTGCTGCTGCTGCTGCTTTCGCGGAGGAGCGCAGGCCCTGGAGTGCTACAGC
TGCGTGAGAAAGCAGATGACGGATGCTCCCCGAACAAGATGAAGACAGTGAAGTGCGCG
CCGGGCGTGGACGTCTGCACCGAGGCCGTGGGGGCGGTGGAGACCATCCACGGACAATTC
TCGCTGGCAGTGCGGGGTTGCGGTTCTGGGACTCCCCGGCAAGAATGACCGCGGCCTGGAT
CTTCACGGGCTTCTGGCGTTCATCCAGCTGCAGCAATGCGCTCAGGATCGCTGCAACGCC
AAGCTCAACCTCACCTCGCGGGCGCTCGACCCGGCAGGTAATGAGAGTGATACCCGCC
AACGGCGTGGAGTGCTACAGCTGTGTGGGCCTGAGCCGGGAGGCGTGCCAGGGTACATCG
CCGCCGGTCTGAGCTGCTACAACGCCAGCGATCATGTCTACAAGGGCTGCTTCGACGGC
AACGTCACCTTGACGGCAGCTAATGTGACTGTGTCTTGCCTGTCCGGGGCTGTGTCCAG
GATGAATTCTGCACTCGGGATGGAGTAACAGGCCAGGGTTCACGCTCAGTGGCTCCTGT
TGCCAGGGGTCCCGCTGTAACCTTGACCTCCGCAACAAGACCTACTTCTCCCCTCGAATC
CCACCCCTTGTCGGGCTGCCCCCTCCAGAGCCACGACTGTGGCCTCAACCACATCTGTC
ACCACTTCTACCTCGGCCCCAGTGAGACCCACATCCACCACCAAACCCATGCCAGCGCCA
ACCAGTCAGACTCCGAGACAGGGAGTAGAACACGAGGCCTCCCGGGATGAGGAGCCAGG
TTGACTGGAGGCGCCGCTGGCCACCAGGACCGCAGCAATTCAGGGCAGTATCCTGCAAAA
GGGGGGCCCCAGCAGCCCCATAATAAAGGCTGTGTGGCTCCACAGCTGGATTGGCAGCC
CTTCTGTTGGCCGTGGCTGCTGGTGTCTTACTGTGAGCTTCTCCACCTGGAAATTTCCCT
CTCACCTACTTCTCTGGCCCTGGGTACCCCTCTTCTCATCACTTCCTGTTCCCACTG
GACTGGGCTGGCCAGCCCCCTGTTTTTCCAACATTCGCCAGTATCCCAGCTTCTGCTGC
GCTGGTTTGCGGCTTTGGGAAATAAAATACCGTTGTATATATTCTGCCAGGGGTGTTCTA
GCTTTTGGAGACAGCTCCTGTATCCTTCTCATCCTTGTCTCTCCGCTTGTCTCTTGTG
ATGTTAGGACAGAGTGAGAGAAGTCAGCTGTCACGGGGAAGGTGAGAGAGAGGATGCTAA
GCTTCCTACTCACTTTCTCCTAGCCAGCCTGGACTTTGGAGCGTGGGGTGGGTGGGACAA
TGGCTCCCCACTCTAAGCACTGCCTCCCCTACTCCCCGCATCTTTGGGGAATCGGTTCCC
CATATGTCTTCTTACTAGACTGTGAGCTCCTCGAGGGGGGGCCCGGTACCCAATTGCCC
CTATAGTGAGTCGTA

136/392

FIGURE 130

MDPARKAGAQAMIWTAGWLLLLLLLRGGAQALECYSCVQKADDGCS PNKMKT VKCAPGV DV
CTEAVGAVETIHGQFSLAVRGCGSLPGKNDRGLDLHGLLAFIQLQQCAQDRCN AKLNLT
SRALDPAGNESAYPPNGVECYSCVGLSREACQGTSPFVVS CYNASDHVYKGC FDNVTLT
AANVTVSLPVRGCVQDEFCTRDGVTGPGFTLSGSCCQGSRCNSDLRNKTYFSPRI PPLVR
LPPPEPTTVASTTSVTTST SAPVRPTSTTKPMPAPTSQTPRQGEHEASRDEE PRLTGGA
AGHQDRSNSGQYPAKG GPQQPHNKGC VAPTAGLAALLLAVAAGVLL

137/392

FIGURE 131

AAACTTGACGCCATGAAGATCCCGGTCCTTCCTGCCGTGGTGCTCCTCTCCCTCCTGGTG
CTCCACTCTGCCCAGGGAGCCACCCTGGGTGGTCCTGAGGAAGAAAGCACCATTGAGAAT
TATGCGTCACGACCCGAGGCCTTTAACACCCCGTTCCTGAACATCGACAAATTGCGATCT
GCGTTTAAGGCTGATGAGTTCCTGAACTGGCACGCCCTCTTGAGTCTATCAAAAGGAAA
CTTCCTTTCCTCAACTGGGATGCCTTTCCTAAGCTGAAAGGACTGAGGAGCGCAACTCCT
GATGCCCAGTGACCATGACCTCCACTGGAAGAGGGGGCTAGCGTGAGCGCTGATTCTCAA
CCTACCATAACTCTTTCCTGCCTCAGGAACTCCAATAAAACATTTTCCATCCAAA

138/392

FIGURE 132

MKIPVLPVAVLLSLLVLHSAQGATLGGPEEESTIENYASRPEAFNTPFLNIDKLRSAFKA
DEFLNWHALFESIKRKLPFLNWDAFPKLKGLRSATPDAQ

139/392

FIGURE 133

CAGTTCCTGAAATCAATGGAGTTAATTTAGGGAATACAAACCAGCCATGGGGGGTGGAGATT
GCCTTTGCCTCAGTGATTCTCACCTGCCTCTCCCTTCTGGCAGCAGGAGTCTCCAGGTT
GTTCTTCTCCAGCCAGTTCCAACCTCAGGAGACAGGTCCCAAGGCCATGGGAGATCTCTCC
TGTGGCTTTGCCGGCCACTCATGAGAGTGTTTTTGTGTAAAGTATTTTTTTAGAATACTGT
TGACTTCTTCATGATTTAATAACCATCCTTTGCGAAGTTTTATGAGGCTTTAGGGGAATG
TCAACCCTCAAATTTTTGTTATACTAGATGGCTTCCATTTACCCACCACTATTTTAAGGT
CCCTTTATTTTTAGGTTCAAGGTTCAATTTGACTTGAGAAAGTGCCCTTCTGCAGCTTCAT
TGATTTTTGTTTATCTTCACTATTAATTGTAACGATTAAAAAAGAATAAGAGCACGCAGAC
CTCTAGGAGAATATTTTTATCCCTGGGTGCCCCCTGACACATTTATGTAGTGATCCCAAAA
TGTGATTGTTAATTTAAATGTTATTCTAATATTAGTACATTTCAGTTGTGATGTAATATGA
ATAACCAGAATCTATTTCTTAAAAGTTTTGAGTATATTTTTTCACTAGATATTTGTATAG
AAAGACTGAATAGTGATG

140/392

FIGURE 134

MGVEIAFASVILTCLSLAAGVSQVLLQFVPTQETGPKAMGDLSCGFAGHS

141/392

FIGURE 135

GGGGAATCTGCAGTAGGTCTGCCGGCGATGGAGTGGTGGGCTAGCTCGCCGCTTCGGCTC
TGGCTGCTGTTGTTCCCTCCTGCCCTCAGCGCAGGGCCGCCAGAAGGAGTCAGGTTCAAAA
TGGAAAGTATTTATTGACCAAATTAACAGGTCTTTGGAGAATTACGAACCATGTTCAAGT
CAAACTGCAGCTGCTACCATGGTGT CATAGAAGAGGATCTAACTCCTTTCCGAGGAGGC
ATCTCCAGGAAGATGATGGCAGAGGTAGTCAGACGGAAGCTAGGGACCCACTATCAGATC
ACTAAGAACAGACTGTACCGGGAAAATGACTGCATGTTCCCCTCAAGGTGTAGTGGTGT
GAGCACTTTATTTTGGAAAGTGATCGGGCGTCTCCCTGACATGGAGATGGTGATCAATGTA
CGAGATTATCCTCAGGTTCCCTAAATGGATGGAGCCTGCCATCCCAGTCTTCTCCTTCAGT
AAGACATCAGAGTACCATGATATCATGTATCCTGCTTGGACATTTTGGGAAGGGGGACCT
GCTGTTTGGCCAATTTATCCTACAGGTCTTGGACGGTGGGACCTCTTCAGAGAAGATCTG
GTAAGGTGAGCAGCACAGTGGCCATGGAAAAAGAAAACTCTACAGCATATTTCCGAGGA
TCAAGGACAAGTCCAGAACGAGATCCTCTCATTCTTCTGTCTCGGAAAAACCCAAAACTT
GTTGATGCAGAATACACCAAAAACCAGGCCTGGAAATCTATGAAAGATACCTTAGGAAAG
CCAGCTGCTAAGGATGTCCATCTTGTGGATCACTGCAAATACAAGTATCTGTTTAATTTT
CGAGGCGTAGCTGCAAGTTTCCGGTTTAAACACCTCTTCTGTGTGGCTCACTTGTTTTTC
CATGTTGGTGATGAGTGGCTAGAATTCTTCTATCCACAGCTGAAGCCATGGGTTCATAT
ATCCCAGTCAAAACAGATCTCTCCAATGTCCAAGAGCTGTTACAATTTGTAAAAGCAAAT
GATGATGTAGCTCAAGAGATTGCTGAAAGGGGAAGCCAGTTTATTAGGAACCATTTGCAG
ATGGATGACATCACCTGTTACTGGGAGAACCTCTTGAGTGAATACTCTAAATTCCTGTCT
TATAATGTAACGAGAAGGAAAGGTTATGATCAAATTATTCCCAAAATGTTGAAAACCTGAA
CTATAGTAGTCATCATAGGACCATAGTCCTCTTTGTGGCAACAGATCTCAGATATCCTAC
GGTGAGAAGCTTACCATAAGCTTGGCTCCTATACCTTGAATATCTGCTATCAAGCCAAAT
ACCTGGTTTTTCCTTATCATGCTGCACCCAGAGCAACTCTTGAGAAAGATTTAAAATGTGT
CTAATACACTGATATGAAGCAGTTCAACTTTTTTGGATGAATAAGGACCAGAAATCGTGAG
ATGTGGATTTTTGAACCCAACTCTACCTTTCATTTTCTTAAGACCAATCACAGCTTGTGCC
TCAGATCATCCACCTGTGTGAGTCCATCACTGTGAAATTGACTGTGTCCATGTGATGATG
CCCTTTGTCCCATTTATTTGGAGCAGAAAATTCGTCAATTTGGAAGTAGTACAACCTATTGC
TGGAATTGTGAAATTATTCAAGGCGTGATCTCTGTCACTTTATTTTAATGTAGGAAACCC
TATGGGGTTTATGAAAAATACTTGGGGATCATTCTCTGAATGGTCTAAGGAAGCGGTAGC
CATGCCATGCAATGATGTAGGAGTTCTCTTTTGTAAAACCATAAACTCTGTTACTCAGGA
GGTTTCTATAATGCCACATAGAAAGAGGCCAATTGCATGAGTAATTATTGCAATTGGATT
TCAGGTTCCCTTTTTTGTGCCTTCATGCCCTACTTCTTAATGCCTCTCTAAAGCCAAA

142/392

FIGURE 136

MEWWASSPLRLWLLLFLLPSAQGRQKESGSKWKVFIDQINRSLENYEPCSSQNCSCYHGV
IEEDLTPFRGGISRKMAEVVRRKLGTHYQITKNRLYREND CMFP SRC SGVEHFILEVIG
RLPD MEMVINVRDYPQVPKWMEPAIPVFSFSKTSEYHDIMYPAWTFWEGGPAVWP IYPTG
LGRWDLFREDLVRSA AQWPWKKNSTAYFRGSRTSPERDPLILLSRKNP KLVD AEYTKNQ
AWKSMKDTLGKPAAKDVHLVDHCKYKYLNFNFRGVAASFRFKHLFLCGSLVFHVGDEWLEF
FYPQLKPWVHYIPVKTDLSNVQELLQFVKANDDVAQEIAERGSQFIRNHLQMD DITCYWE
NLLSEYSKFLSYNVTRRKG YDQIIPKMLKTEL

143/392

FIGURE 137

ATTCTCCTAGAGCATCTTTGGAAGCATGAGGCCACGATGCTGCATCTTGGCTCTTGTCTG
CTGGATAACAGTCTTCCTCCTCCAGTGTTCAAAGGAACTACAGACGCTCCTGTTGGCTC
AGGACTGTGGCTGTGCCAGCCGACACCCAGGTGTGGGAACAAGATCTACAACCCTTCAGA
GCAGTGCTGTTATGATGATGCCATCTTATCCTTAAAGGAGACCCGCCGCTGTGGCTCCAC
CTGCACCTTCTGGCCCTGCTTTGAGCTCTGCTGTCCCGAGTCTTTTGGCCCCCAGCAGAA
GTTTCTTGTGAAGTTGAGGGTTCTGGGTATGAAGTCTCAGTGTCACCTTATCTCCCATCTC
CCGGAGCTGTACCAGGAACAGGAGGCACGTCCTGTACCCATAAAAAACCCAGGCTCCACT
GGCAGACGGCAGACAAGGGGAGAAGAGACGAAGCAGCTGGACATCGGAGACTACAGTTGA
ACTTCGGAGAGAAGCAACTTGACTTCAGAGGGATGGCTCAATGACATAGCTTTGGAGAGG
AGCCCAGCTGGGGATGGCCAGACTTCAGGGGAAGAATGCCTTCCTGCTTCATCCCCTTTC
CAGCTCCCCCTTCCCCTGAGAGCCACTTTCATCGGCAATAAAATCCCCACATTTACCATCT

144/392

FIGURE 138

MRPRCCILALVCWITVFLQCSKGTTDAPVGSLWLCQPTPRCGNKIYNPSEQCCYDDAI
LSLKETRRCGSTCTFWPCFELCCPESFGPQQKFLVKLRVLGMKSQCHLSPIRSCTRNR
HVLYP

Important features:

Signal sequence:

amino acids 1-21

N-myristoylation sites:

amino acids 33-39, 70-76

145/392

FIGURE 139

CCTCTGTCCACTGCTTTTCGTGAAGACAAGATGAAGTTCACAATTGTCTTTGCTGGACTTC
TTGGAGTCTTTCTAGCTCCTGCCCTAGCTAACTATAATATCAACGTCAATGATGACAACA
ACAATGCTGGAAGTGGGCAGCAGTCAGTGAGTGTCAACAATGAACACAATGTGGCCAATG
TTGACAATAACAACGGATGGGACTCCTGGAATTCCATCTGGGATTATGGAAATGGCTTTG
CTGCAACCAGACTCTTTCAAAGAAGACATGCATTGTGCACAAAATGAACAAGGAAGTCA
TGCCCTCCATTCAATCCCTTGATGCACTGGTCAAGGAAAAGAAGCTTCAGGGTAAGGGAC
CAGGAGGACCACCTCCCAAGGGCCTGATGTACTCAGTCAACCCAAACAAAGTCGATGACC
TGAGCAAGTTCGGAAAAAACATTGCAAACATGTGTCTGGGATTCCAACATACATGGCTG
AGGAGATGCAAGAGGCAAGCCTGTTTTTTTACTCAGGAACGTGCTACACGACCAGTGTAC
TATGGATTGTGGACATTTCTTCTGTGGAGACACGGTGGAGAACTAAACAATTTTTTAAA
GCCACTATGGATTTAGTCATCTGAATATGCTGTGCAGAAAAAATATGGGCTCCAGTGGTT
TTTACCATGTCAATCTGAAATTTTCTCTACTAGTTATGTTTGATTTCTTTAAGTTTCAA
TAAATCATTTAGCATTGAAAAAA

146/392

FIGURE 140

MKFTIVFAGLLGVFLAPALANYNINVNDDNNNAGSGQQSVSVNNEHNVANVDNNGWDSW
NSIWDYGNNGFAATRLFQKKTCIVHKMNKEVMPSIQSLDALVKEKKLQKGPGGPPPKGLM
YSVNPKNKVDLDSKFGKNIANMCRGIPTYMAEEMQEASLFFYSGTCYTTSVLWIVDISFCG
DTVEN

Signal Peptide:
amino acids 1-20

N-myristoylation Sites:
amino acids 67-72, 118-123, 163-168

Flavodoxin protein homology:
amino acids 156-174

147/392

FIGURE 141

GGTCCTTAATGGCAGCAGCCGCCGCTACCAAGATCCTTCTGTGCCTCCCGCTTCTGCTCC
TGCTGTCCGGCTGGTCCCGGGCTGGGCGAGCCGACCCTCACTCTCTTTGCTATGACATCA
CCGTCATCCCTAAGTTTCAAGACCTGGACCACGGTGGTGTGCGGTTCAAGGCCAGGTGGATG
AAAAGACTTTTCTTCACTATGACTGTGGCAACAAGACAGTCACACCTGTCAGTCCCCTGG
GGAAGAACTAAATGTCACAACGGCCTGGAAAGCACAGAACCCAGTACTGAGAGAGGTGG
TGGACATACTTACAGAGCAACTGCGTGACATTGAGCTGGAGAATTACACACCCAAGGAAC
CCCTCACCTGTCAGGCAAGGATGTCTTGTGAGCAGAAAGCTGAAGGACACAGCAGTGGAT
CTTGGCAGTTCAGTTTCGATGGGCAGATCTTCCTCCTCTTTGACTCAGAGAAGAGAATGT
GGACAACGGTTCATCCTGGAGCCAGAAAGATGAAAGAAAAGTGGGAGAATGACAAGGTG
TGGCCATGTCCTTCCATTACTTCTCAATGGGAGACTGTATAGGATGGCTTGAGGACTTCT
TGATGGGCATGGACAGCACCTGGAGCCAAGTGCAGGAGCACCCTCGCCATGTCCTCAG
GCACAACCCAACCTCAGGGCCACAGCCACCACCCTCATCCTTTGCTGCCTCCTCATCATCC
TCCCCTGCTTCATCCTCCCTGGCATCTGAGGAGAGTCCTTTAGAGTGACAGGTTAAAGCT
GATACCAAAGGCTCCTGTGAGCACGGTCTTGATCAAACCTCGCCCTTCTGTCTGGCCAGC
TGCCACGACCTACGGTGTATGTCCAGTGGCCTCCAGCAGATCATGATGACATCATGGAC
CCAATAGCTCATTCAGTGCCTTGATTCTTTTGCCAACAATTTTACCAGCAGTTATACCT
AACATATTATGCAATTTTCTCTTGGTGCTACCTGATGGAATTCCTGCACTTAAAGTTCTG
GCTGACTAAACAAGATATATCATTTTTCTTTCTTCTTTTTGTTTGGAAAATCAAGTACT
TCTTTGAATGATGATCTCTTTCTTGCAAATGATATTGTCAGTAAAATAATCACGTTAGAC
TTCAGACCTCTGGGGATTCTTTCCGTGTCCTGAAAGAGAATTTTTAAATTATTTAATAAG
AAAAAATTTATATTAATGATTGTTTCCTTTAGTAATTTATTGTTCTGTACTGATATTTAA
ATAAAGAGTTCTATTTCCCAAAAAAAAAAAAAAAAAAAAA

148/392

FIGURE 142

MAAAAATKILLCLPLLLLLSGWSRAGRADPHSLCYDITVIPKFRPGPRWCAVQGQVDEKT
FLHYDCGNKTVTPVSPLGKKLNVTTAWKAQNPVLREVVDILTEQLRDIQLENYTPKEPLT
LQARMSCEQKAEGHSSGSWQFSFDGQIFLLFDSEKRMWTTVHPGARKMKEKWENDKVVAM
SFHYFSMGDCIGWLEDFLMGMDSTLEPSAGAPLAMSSGTTQLRATATTLILCCLLIILPC
FILPGI

Important features:**Signal peptide:**

amino acids 1-25

Transmembrane domain:

amino acids 224-246

N-glycosylation site:

amino acids 68-72, 82-86

N-myristoylation site:

amino acids 200-206, 210-216

Amidation site:

amino acids 77-81

149/392

FIGURE 143

AATGTGAGAGGGGCTGATGGAAGCTGATAGGCAGGACTGGAGTGTTAGCACCAGTACTGG
ATGTGACAGCAGGCAGAGGAGCACCTTAGCAGCTTATTCAGTGTCCGATTCTGATTCCGGC
AAGGATCCAAGCATGGAATGCTGCCGTCGGGCAACTCCTGGCACACTGCTCCTCTTTCTG
GCTTTCTGCTCCTGAGTTCCAGGACCGCACGCTCCGAGGAGGACCGGGACGGCCTATGG
GATGCCTGGGGCCCATGGAGTGAATGCTCACGCACCTGCGGGGGAGGGGCTCCTACTCT
CTGAGGCGCTGCCTGAGCAGCAAGAGCTGTGAAGGAAGAAATATCCGATACAGAACATGC
AGTAATGTGGACTGCCACCAGAAGCAGGTGATTTCCGAGCTCAGCAATGCTCAGCTCAT
AATGATGTCAAGCACCATGGCCAGTTTTATGAATGGCTTCCTGTGTCTAATGACCCTGAC
AACCCATGTTCACTCAAGTGCCAAGCCAAAGGAACAACCCTGGTTGTTGAAC TAGCACCT
AAGGTCTTAGATGGTACGCGTTGCTATACAGAATCTTTGGATATGTGCATCAGTGGTTTA
TGCCAAATTGTTGGCTGCGATCACCAGCTGGGAAGCACCGTCAAGGAAGATAACTGTGGG
GTCTGCAACGGAGATGGGTCCACCTGCCGGCTGGTCCGAGGGCAGTATAAATCCCAGCTC
TCCGCAACCAAATCGGATGATACTGTGGTTGCACTTCCCTATGGAAGTAGACATATTCGC
CTTGTCTTAAAGGTCTGATCACTTATATCTGGAAACCAAACCCTCCAGGGGACTAAA
GGTGAAAACAGTCTCAGCTCCACAGGAACCTTTCTTGTGGACAATTCTAGTGTGGACTTC
CAGAAATTTCCAGACAAAGAGATACTGAGAATGGCTGGACCACTCACAGCAGATTTTATT
GTCAAGATTTCGTAACCTCGGGCTCCGCTGACAGTACAGTCCAGTTCATCTTCTATCAACCC
ATCATCCACCGATGGAGGGAGACGGATTTCTTTCTTGCTCAGCAACCTGTGGAGGAGGT
TATCAGCTGACATCGGCTGAGTGCTACGATCTGAGGAGCAACCGTGTGGTTGCTGACCAA
TACTGTCACTATTACCCAGAGAACATCAAACCCAAACCCAAGCTTCAGGAGTGCAACTTG
GATCCTTGTCCAGCCAGTGACGGATACAAGCAGATCATGCCTTATGACCTCTACCATCCC
CTTCCTCGGTGGGAGGCCACCCCATGGACCGCGTGCTCCTCCTCGTGTGGGGGGGGCATC
CAGAGCCGGGCAGTTTCTGTGTGGAGGAGGACATCCAGGGGCATGTCACTTCAGTGGAA
GAGTGGAAATGCATGTACACCCCTAAGATGCCCATCGCGCAGCCCTGCAACATTTTTTGAC
TGCCCTAAATGGCTGGCACAGGAGTGGTCTCCGTGCACAGTGACATGTGGCCAGGGCCTC
AGATAACCGTGTGGTCTCTGCATCGACCATCGAGGAATGCACACAGGAGGCTGTAGCCCA
AAAACAAAGCCCCACATAAAAGAGGAATGCATCGTACCCACTCCCTGCTATAAACCCAAA
GAGAACTTCCAGTCGAGGCCAAGTTGCCATGGTTCAAACAAGCTCAAGAGCTAGAAGAA
GGAGCTGCTGTGTGAGAGGAGCCCTCGTAAGTTGTAAAGCACAGACTGTTCTATATTG
AACTGTTTTGTTTAAAGAAAGCAGTGTCTCACTGGTTGTAGCTTTTCATGGGTCTCTGAAC
TAAGTGTAATCATCTCACCAAAGCTTTTTGGCTCTCAAATTAAAGATTGATTAGTTTTCAA
AAAAAAAAA

150/392

FIGURE 144

MECCRATPGTLLLFLAFLLLSSRTARSEEDRDGLWDAWGPWSECSRTC GGGASYSLRRC
LSSKSCEGRNIRYRTCSNVDCPPEAGDFRAQQCSAHNDVKHHGQFYEWLPVSNDPDNPCS
LKCQAKGTTLVVELAPKVLDGTRCYTESLDMCISGLCQIVGCDHQLGSTVKEDNCGVCNG
DGSTCRLVRGQYKSQLSATKSDDTVVALPYGSRHIRLVKGPDHLYLETKTLQGTKGENS
LSSTGTFLVDNSSVDFQKFPDKEILRMAGPLTADFIKIRNSGSADSTVQFIFYQPIIHR
WRETDFFPCSATCGGGYQLTSAECYDLRSNRVVADQYCHYYPENIKPKPKLQECNLDPCP
ASDGYKQIMPYDLYHPLPRWEATPWTACSSSCGGGIQSRVSCVEEDIQGHVTSVEEWKC
MYTPKMPIAQPCNIFDCPKWLAQEWSPCTVTCGQGLRYRVVLCIDHRGMHTGGCSPKTKP
HIKEECIVPTPCYKPKEKLPVEAKLPWFKQAQELEEGA AVSEEPS

Important features:**Signal peptide:**

amino acids 1-25

N-glycosylation site:

amino acids 251-254

Thrombospondin 1:

amino acids 385-399

von Willebrand factor type C domain proteins:

amino acids 385-399, 445-459 and 42-56

151/392

FIGURE 145

GGAGGAGGGAGGGCGGGCAGGCGCCAGCCCAGAGCAGCCCCGGGCACCAGCACGGACTCT
CTCTTCCAGCCCAGGTGCCCCCACTCTCGCTCCATTTCGGCGGGAGCACCCAGTCCTGTA
CGCCAAGGAACTGGTCTCTGGGGGCACCAATGGTTTTCGGCGGCAGCCCCCAGCCTCCTCATC
CTTCTGTTGCTGCTGCTGGGGTCTGTGCCTGCTACCGACGCCCCGCTCTGTGCCCTGAAG
GCCACGTTCTCTGGAGGATGTGGCGGGTAGTGGGGAGGCCGAGGGCTCGTCGGCCTCCTCC
CCGAGCCTCCCGCCACCCTGGACCCCGGCCCTCAGCCCCACATCGATGGGGCCCCAGCCC
ACAACCTTGGGGGGCCCATCACCCCCACCAACTTCCTGGATGGGATAGTGGACTTCTTC
CGCCAGTACGTGATGCTGATTGCTGTGGTGGGCTCCCTGGCCTTTCTGCTGATGTTTCATC
GTCTGTGCCGCGGTTCATCACCCGGCAGAAGCAGAAGGCCTCGGCCTATTACCCATCGTCC
TTCCCCAAGAAGAAGTACGTGGACCAGAGTGACCGGGCCGGGGCCCCCGGGCCTTCAGT
GAGGTCCCCGACAGAGCCCCCGACAGCAGGCCCGAGGAAGCCCTGGATTCTCTCCCGGCAG
CTCCAGGCCGACATCTTGCCCGCCACCCAGAACCTCAAGTCCCCACCAGGGCTGCACTG
GGCGGTGGGGACGGAGCCAGGATGGTGGAGGGCAGGGGCGCAGAGGAAGAGGAGAAGGGC
AGCCAGGAGGGGGACAGGAAGTCCAGGGACATGGGGTCCCAGTGGAGACACCAGAGGCG
CAGGAGGAGCCGTGCTCAGGGGTCTTGAGGGGGCTGTGGTGGCCGGTGAGGGCCAAGGG
GAGCTGGAAGGGTCTCTCTTGTTAGCCAGGAAGCCAGGGACCAGTGGGTCCCCCGAA
AGCCCCCTGTGCTTGCAGCAGTGTCCACCCAGTGTCTAACAGTCCTCCCGGGCTGCCAGC
CCTGACTGTTCGGGCCCCCAAGTGGTCACCTCCCCGTGTATGAAAAGGCCTTCAGCCCTGA
CTGCTTCCTGACACTCCCTCCTTGGCCTCCCTGTGGTGCCAATCCCAGCATGTGCTGATT
CTACAGCAGGCAGAAATGCTGGTCCCCGGTGCCCCGGAGGAATCTTACCAAGTGCCATCA
TCCTTCACCTCAGCAGCCCCAAAGGGCTACATCCTACAGCACAGCTCCCTTGACAAAGTG
AGGGAGGGCACGTGTCCCTGTGACAGCCAGGATAAAACATCCCCCAAAGTGCTGGGATTA
CAGGCGTGAGCCACCGTGCCCGGCCAACTACTTTTAAAACAGCTACAGGGTAAATC
CTGCAGCACCCACTCTGGAAAATACTGCTCTTAATTTTCTGAAGGTGGCCCCCTGTTTC
TAGTTGGTCCAGGATTAGGGATGTGGGGTATAGGGCATTTAAATCCTCTCAAGCGCTCTC
CAAGCACCCCCGGCCTGGGGGTGAGTTTCTCATCCCGCTACTGCTGCTGGGATCAGGTTG
AATGAATGGAACCTTTCCTGTCTGGCCTCCAAAGCAGCCTAGAAGCTGAGGGGCTGTGTT
TGAGGGGACCTCCACCTGGGGGAAGTCCGAGGGGCTGGGGAAGGGTTTCTGACGCCAGC
CTGGAGCAGGGGGGCCCTGGCCACCCCTGTTGCTCACACATTGTCTGGCAGCCTGTGTC
CACAATATTCTGTCAGTCTCGACAGGGAGCCTGGGCTCCGTCTGCTTTAGGGAGGCTCT
GGCAGGAGGTCTCTCCCCATCCCTCCATCTGGGGCTCCCCAACCTCTGCACAGCTCT
CCAGGTGCTGAGATATAATGCACCAGCACATAAACCTTTATTCCGGCCTGAAAAAAAAA
AAGA

152/392

FIGURE 146

MVSAAAPSLILLLLLLLGSVPATDARSVPLKATFLEDVAGSGEAEAGSSASSPSLPPPWTP
ALSPTSMGPQPTTLGGPSPPTNFLDGIVDFFRQYVMLIAVVGSLAFLLMFIVCAAVITRQ
KQKASAYYPSSFPKKKYVDQSDRAGGPRAFSEVPDRAPDSRPEEALDSSRQLQADILAAT
QNLKSPTRAALGGGDGARMVEGRGAEEEEKGSQEGDQEVQGHGVPVETPEAQEEPCSGVL
EGAVVAGEGQGELEGSLLLAQEAQGPVGPPESPACSSVHPSV

Signal peptide:
amino acids 1-25

Transmembrane domain:
amino acids 94-118

N-myristoylation site:
amino acids 18-24, 40-46, 46-52, 145-151, 192-198, 193-199,
211-217, 238-244, 242-248

153/392

FIGURE 147

GAAAGACGTGGTCCTGACAGACAGACAATCCTATTCCCTACCAAATGAAGATGCTGCTG
CTGCTGTGTTTGGGACTGACCCTAGTCTGTGTCCATGCAGAAGAAGCTAGTTCTACGGGA
AGGAACTTTAATGTAGAAAAGATTAATGGGGAATGGCATACTATTATCCTGGCCTCTGAC
AAAAGAGAAAAGATAGAAGAACATGGCAACTTTAGACTTTTTCTGGAGCAAATCCATGTC
TTGGAGAATTCTTAGTTCTTAAAGTCCATACTGTAAGAGATGAAGAGTGCTCCGAATTA
TCTATGGTTGCTGACAAAACAGAAAAGGCTGGTGAATATTCTGTGACGTATGATGGATTCT
AATACATTTACTATACCTAAGACAGACTATGATAACTTTCTTATGGCTCACCTCATTAACT
GAAAAGGATGGGGAAACCTTCCAGCTGATGGGGCTCTATGGCCGAGAACCAGATTTGAGT
TCAGACATCAAGGAAAGGTTTGCACAACCTATGTGAGGAGCATGGAATCCTTAGAGAAAAT
ATCATTGACCTATCCAATGCCAATCGCTGCCTCCAGGCCCGAGAATGAAGAATGGCCTGA
GCCTCCAGTGTTGAGTGGACACTTCTCACCAGGACTCCACCATCATCCCTTCCTATCCAT
ACAGCATCCCCAGTATAAATTCTGTGATCTGCATTCCATCCTGTCTCACTGAGAAGTCCA
ATTCCAGTCTATCAACATGTTACCTAGGATACCTCATCAAGAATCAAAGACTTCTTTAAA
TTTCTCTTTGATACACCCTTGACAATTTTTCATGAAATTATTCCTCTTCCTGTTCAATAA
ATGATTACCCTTGCACTTAA

154/392

FIGURE 148

MKMLLLLCLGLTLVCVHAEESSTGRNFNVEKINGEWHTIILASDKREKIEEHGNFRLFL
EQIHVLENSLVKLVHTVRDEECSELSMVADKTEKAGEYSVTYDGFNTFTIPKTDYDNFLM
AHLINEKDGETFQLMGLYGREPDLSSDIKERFAQLCEEHGILRENIIDLSNANRCLQARE

155/392

FIGURE 149

GTGGACTCTGAGAAGCCCAGGCAGTTGAGGACAGGAGAGAGAAGGCTGCAGACCCAGAGG
GAGGGAGGACAGGGAGTCGGAAGGAGGAGGACAGAGGAGGGCACAGAGACGCAGAGCAAG
GGCGGCAAGGAGGAGACCTTGGTGGGAGGAAGACACTCTGGAGAGAGAGGGGGCTGGGCA
GAGATGAAGTTCCAGGGGCCCCCTGGCCTGCCTCCTGCTGGCCCTCTGCCTGGGCAGTGGG
GAGGCTGGCCCCCTGCAGAGCGGAGAGGAAAGCACTGGGACAAATATTGGGGAGGCCCTT
GGACATGGCCTGGGAGACGCCCTGAGCGAAGGGGTGGGAAAGGCCATTGGCAAAGAGGCC
GGAGGGGCAGCTGGCTCTAAAGTCAGTGAGGCCCTTGGCCAAGGGACCAGAGAAGCAGTT
GGCACTGGAGTCAGGCAGGTTCCAGGCTTTGGCGCAGCAGATGCTTTGGGCAACAGGGTC
GGGGAAGCAGCCCATGCTCTGGGAAACACTGGGCACGAGATTGGCAGACAGGCAGAAGAT
GTCATTCGACACGGAGCAGATGCTGTCCGCGGCTCCTGGCAGGGGGTGCCTGGCCACAGT
GGTGCTTGGGAAACTTCTGGAGGCCATGGCATCTTTGGCTCTCAAGGTGGCCTTGGAGGC
CAGGGCCAGGGCAATCCTGGAGGTCTGGGGACTCCGTGGGTCCACGGATACCCCGGAAAC
TCAGCAGGCAGCTTTGGAATGAATCCTCAGGGAGCTCCCTGGGGTCAAGGAGGCAATGGA
GGGCCACCAAACCTTTGGGACCAACACTCAGGGAGCTGTGGCCCAGCCTGGCTATGGTTCA
GTGAGAGCCAGCAACCAGAATGAAGGGTGACGAATCCCCCACCATTCTGGCTCAGGTGGA
GGCTCCAGCAACTCTGGGGGAGGCAGCGGCTCACAGTCGGGCAGCAGTGGCAGTGGCAGC
AATGGTGACAACAACAATGGCAGCAGCAGTGGTGGCAGCAGCAGTGGCAGCAGCAGTGGC
AGCAGCAGTGGCGGCAGCAGTGGCGGCAGCAGTGGTGGCAGCAGTGGCAACAGTGGTGGC
AGCAGAGGTGACAGCGGCAGTGAGTCCTCCTGGGGATCCAGCACCGGCTCCTCCTCCGGC
AACCACGGTGGGAGCGGCGGAGGAAATGGACATAAACCCGGGTGTGAAAAGCCAGGGGAAT
GAAGCCCGCGGGAGCGGGGAATCTGGGATTTCAGGGCTTCAGAGGACAGGGAGTTTCCAGC
AACATGAGGGAAATAAGCAAGAGGGCAATCGCCTCCTTGGAGGCTCTCGAGACAAATTAT

156/392

FIGURE 150

MKFQGPLACLLLLALCLGSGEAGPLQSGEESTGTNIGEALGHGLGDALSEGVGKAIGKEAG
 GAAGSKVSEALQGQTREAVGTGVRQVPGFGAADALGNRVGEAAHALGNTGHEIGRQAEVD
 IRHGADAVRGSWQGVPGHSGAWETSGGHGIFGSQGGLGGQQQGNPGGLGTPWVHGYPGNS
 AGSFGMNPQGAPWGQGGNGGPPNFGTNTQGAVAQPGYGSVRASNQNEGCTNPPPSGSGGG
 SSNSGGGSGSQSGSSGSGSNGDNNNGSSSGSSSGSSSGSSSGSSSGSSSGSSSGSSSGS
 RGDSGSESSWGSSTGSSSGNHGGSGGGNGHKPGCEKPGNEARGSGESGIQGFRGQGVSSN
 MREISKEGNRLLGSGDNYRGQSSSWGSGGGDAVGGVNTVNSETSPGMFNFDTFWKNFKS
 KLGFINWDAINKDORSSRIP

Signal peptide:

amino acids 1-21

N-glycosylation site:

amino acids 265-269

Glycosaminoglycan attachment site:

amino acids 235-239, 237-241, 244-248, 255-259, 324-328,
388-392

Casein kinase II phosphorylation site:

amino acids 26-30, 109-113, 259-263, 300-304, 304-308

N-myristoylation site:

amino acids 17-23, 32-38, 42-48, 50-56, 60-66, 61-67, 64-70,
74-80, 90-96, 96-102, 130-136, 140-146, 149-155, 152-158,
155-161, 159-165, 163-169, 178-184, 190-196, 194-200,
199-205, 218-224, 236-242, 238-244, 239-245, 240-246,
245-251, 246-252, 249-252, 253-259, 256-262, 266-272,
270-276, 271-277, 275-281, 279-285, 283-289, 284-290,
287-293, 288-294, 291-297, 292-298, 295-301, 298-304,
305-311, 311-317, 315-321, 319-325, 322-328, 323-329,
325-331, 343-349, 354-360, 356-362, 374-380, 381-387,
383-389, 387-393, 389-395, 395-401

Cell attachment sequence:

amino acids 301-304

157/392

FIGURE 151

CGGCCACAGCTGGCATGCTCTGCCTGATCGCCATCCTGCTGTATGTCCTCGTCCAGTACC
TCGTGAACCCCGGGGTGCTCCGCACGGACCCAGATGTCAAGAATATGAACACGTGGCTG
CTGTTCCCTCCCCCTGTTCCCGGTGCAGGTGCAGACCCTGATAGTCGTGATCATCGGGATG
CTCGTGCTCCTGCTGGACTTTCTTGGCTTGGTGCACCTGGGCCAGCTGCTCATCTTCCAC
ATCTACCTGAGTATGTCCCCACCCCTAAGCCCCGATCCCCCAAGGCTGGGTGGTCAGA
GCTGCTCATCTTACACCTCTACTTGAGTATGTCCCTAACCCTGAGCCCCCACGCCTGGG
GCCAGAGTCTTTGTCCCCGTGTGCGCATGTGTTACAGGTGAGCCTCTCCAGAAGTGAG
ATCATGGACAAAAAGGGCAAATCACAGGAAGAAATTAAATCCATGAGGACCCAGCAGGCC
CAGCAAGAAGCTGAACTCACGCCGAGACCTGCAGGAGTGGTGCCAGGTGCTTGAAGTAAC
AAGTTTAAATGTTTCAAGACAATGGAATGGAATCTATTAGGCAAGAACAGGACATTATG
AAATAAGGACAGGTGGACTTCCAAAAACACAAGTAGAAATTCTAACAAATGAAATATATTA
CAGGCAGGTCACCCACTAACCAAACTGAAGCGAGAGCTGTGGTCTTGCTTGGTCTCA
CAGTGGGCACAGCGGTAGGCGGTGAGTCATGTTGCTGAACGACGGAGGGTAAACTCCCCA
GCCCCAAGAAACCTGTGTTGGAAGTAACAACAACCTCCCTGCTCCTGGCACCAGCCGTT
TTGGTCATGGTGGGCCAGCTGCAAAGCGTCTTCCATTCTCTGGGCAGTGGTGGCCCCGAG
GCTGTGGCCTCTCAGGGGGTTTCTGTGGACACGGGCAGCAGAGTGTGTCCAGGCCAGCCC
CCAAGAATGCCCTGCTCCTGACAGCTTGGCCAACCCCTGGTCAGGGCAGAGGGAGTTGGG
TGGGTCAGGCTCTGGGCTCACCTCCATCTCCAGAGCATCCCCTGCCTGCAGTTGTGGCAA
GAACGCCCAGCTCAGAAATGAACACACCCCAAGAGCCTCCTTGTTTCATAACCACAGGT
TACCCTACAAACCACTGTCCCCACACAACCTGGGGATGTTTTTAAACACACACCTCTAA
CGCATATCTTACAGTCACTGTTGTCTTGCCTGAGGGTTGAATTTTTTTTAAATGAAAGTGC
AATGAAAATCACTGGATTAAATCCTACGGACACAGAGCTGAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAA

158/392

FIGURE 152

MNTWLLFLPLFPVQVQTLIVVIIGMLVLLLDLGLVHLGQLLI FHIYLSMSPTLSPRSPQ
GWVVRAAHLTPLLEYVPNPEPPTPGARVFVPRVRMCSGSASPRSEIMDKKGKSQEEIKSM
RTQQAQQEAEELTPRPAGVVPGA

159/392

FIGURE 153

AACTGGAAGGAAAGAAAGAAAGGTCAGCTTTGGCCCAGATGTGGTTACCCCTTGGTCTCC
TGTCTTTATGTCCTTCTCCTCTTCCTATTCTGTCATCTCCCTCACTTAAGTCTCAGGCCT
GTCAGCAGCTCCTGTGGACATTGCCATCCCCTCTGGTAGCCTTCAGAGCAAACAGGACAA
CCTATGTTATGGATGTTTCCACCAACCAGGGTAGTGGCATGGAGCACCGTAACCATCTGT
GCTTCTGTGATCTCTATGACAGAGCCACTTCTCCACCTCTGAAATGTTCCCTGCTCTGAA
ATCTGGCATGAGATGGCACAGGTGACCACGCAGAAGCCACCAGAATCTTGCCTGCCCTAT
TCCTCCTCCCAAGTCTGTTCTCTTATTGTCAACCTCAGCACAAACAGGCTGGCGCCAATGG
CATTACAGAGAAAGCAATCTGTGTGGCTAGTGGGCAGATTACCATGCAAGCCCCAGGAGA
AATGGAGGAGCTTTGTAGCCACCTCCCTGTGAGCCAGTATTAACATGTCCCCTTCCCCCT
GCCCCGCCGTAGATTGAGGACATTGCCCCCTGTGTGCCACCAAACAGGACTTCCCCCTT
GGCTTGGCATCCCTGGCTCTCTCCTGGTACCCAGCAAGACGTCTGTTCCAGGGCAGTGTA
GCATCTTTCAAGCTCCGTTACTATGGCGATGGCCATGATGTTACAATCCCACTTGCCTGA
ATAATCAAGTGGAAGGGGAAGCAGAGGGAAATGGGGCCATGTGAATGCAGCTGCTCTGT
TCTCCCTACCCTGAGGAAAAACCAAAGGGAAGCAACAGGAACTTCTGCAACTGGTTTTTA
TCGGAAGATCATCCTGCCTGCAGATGCTGTTGAAGGGGCACAAGAAATGTAGCTGGAGA
AGATTGATGAAAGTGCAGGTGTGTAAGGAAATAGAACAGTCTGCTGGGAGTCAGACCTGG
AATTCTGATTCCAACTCTTTATTACTTTGGGAAGTCACTCAGCCTCCCCGTAGCCATCT
CCAGGGTGACGGAACCCAGTGTATTACCTGCTGGAACCAAGGAACTAACAATGTAGGTT
ACTAGTGAATACCCCAATGGTTTCTCCAATTATGCCCATGCCACCAAAACAATAAAACAA
AATTCTCTAACACTGAAA

160/392

FIGURE 154

MWLPLGLLSLCLSPLPILSSPSLKSQACQQLLWTLPSPLVAFRANRTTYVMDVSTNQGS
MEHRNHLCFCDLYDRATSPPLKCSLL

162/392

FIGURE 156

MAGIKALISLSFGGAIGLMFLMLGCALPIYNKYWPLFVLFFYILSPIPYCIARRLVDDTD
AMSNACKELAIFLTGTGIVVSAFGLPIVFARAHLEWGACALVLTGNTVIFATILGFFLVF
GSNDDFSQWQW

163/392

FIGURE 157

GTTCCTCATAGTTGGCGTCTTCTAAAGGAAAAACACTAAAATGAGGAACTCAGCGGACCG
GGAGCGACGCAGCTTGAGGGAAGCATCCCTAGCTGTTGGCGCAGAGGGGCGAGGCTGAAG
CCGAGTGGCCCGAGGTGTCTGAGGGGCTGGGGCAAAGGTGAAAGAGTTTCAGAACAAAGCT
TCCTGGAACCCATGACCCATGAAGTCTTGTCGACATTTATACCGTCTGAGGGTAGCAGCT
CGAAACTAGAAGAAGTGGAGTGTGTCAGGGACGGCAGTATCTCTTTGTGTGACCCTGGC
GGCCTATGGGACGTTGGCTTCAGACCTTTGTGATACACCATGCTGCGTGGGACGATGACG
GCGTGGAGAGGAATGAGGCCTGAGGTCACTGGCTTGCTCCTCCTAGCCACAGCAGGC
TGCTTTGCTGACTTGAACGAGGTCCCTCAGGTACCGTCCAGCCTGCGTCCACCGTCCAG
AAGCCCGGAGGCACCTGTGATCTTGGGCTGCGTGGTGGAACTCCAAGGATGAATGTAACC
TGGCGCCTGAATGAAAGGAGCTGAATGGCTCGGATGATGCTCTGGGTGTCTCATCACC
CACGGGACCCTCGTCATCACTGCCCTTAACAACCACACTGTGGGACGGTACCAGTGTGTG
GCCCCGATGCCTGCGGGGGCTGTGGCCAGCGTGCCAGCCACTGTGACACTAGCCAATCTC
CAGGACTTCAAGTTAGATGTGCAGCACGTGATTGAAGTGGATGAGGGAAACACAGCAGTC
ATTGCCTGCCACCTGCCTGAGAGCCACCCCAAAGCCAGGTCCGGTACAGCGTCAAACAA
GAGTGGCTGGAGGCCTCCAGAGGTAACCTGATCATGCCCTCAGGGAACCTCCAGATT
GTGAATGCCAGCCAGGAGGACGAGGGCATGTACAAGTGTGCAGCCTACAACCCAGTGACC
CAGGAAGTGAAAACCTCCGGCTCCAGCGACAGGCTACGTGTGCGCCGCTCCACCGCTGAG
GCTGCCCCGCATCATCTACCCCCCAGAGGCCCAAACCATCATCGTCACCAAAGGCCAGAGT
CTCATTCTGGAGTGTGTGGCCAGTGGAATCCACCCCCACGGGTCACTGGGCCAAGGAT
GGGTCCAGTGTACCGGCTACAACAAGACGCGCTTCTGTGAGCAACCTCCTCATCGAC
ACCACCAGCGAGGAGGACTCAGGCACCTACCGCTGCATGGCCGACAATGGGGTTGGGCAG
CCCGGGGCAGCGGTCACTCTTACAATGTCCAGGTGTTTGAACCCCCTGAGGTACCATG
GAGCTATCCCAGCTGGTCATCCCCCTGGGGCCAGAGTGCCAAGCTTACCTGTGAGGTGCGT
GGGAACCCCCCGCCCTCCGTGCTGTGGCTGAGGAATGCTGTGCCCTCATCTCCAGCCAG
CGCCTCCGGCTCTCCCGCAGGGCCCTGCGCGTGCTCAGCATGGGGCCTGAGGACGAAGGC
GTCTACCAGTGCATGGCCGAGAACGAGGTTGGGAGCGCCCATGCCGTAGTCCAGCTGCCG
ACCTCCAGGCCAAGCATAACCCCAAGGCTATGGCAGGATGCTGAGCTGGCTACTGGCACA
CCTCCTGTATCACCCCTCCAAACTCGGCAACCCCTGAGCAGATGCTGAGGGGGCAACCGGCG
CTCCCCAGACCCCCAACGTCAGTGGGGCCTGCTTCCCCGAAGTGTCCAGGAGAGAAGGGG
CAGGAGGGCTCCCGCCGAGGCTCCCATCATCTCAGCTCGCCCCGCACCTCCAAGACAGAC
TCATATGAACCTGGTGTGGCGGCTCGGCATGAGGGCAGTGGCCGGGCGCAATCCTCTAC
TATGTGGTGAAACACCGCAAGCAGGTACAAATTCCTCTGACGATTGGACCATCTCTGGC
ATTCCAGCCAACCAGCACCGCCTGACCCCTACCAGACTTGACCCCGGGAGCTTGTATGAA
GTGGAGATGGCAGCTTACAACCTGTGCGGGAGAGGGCCAGACAGCCATGGTCACCTTCCGA
ACTGGACGGCGGCCCAAACCCGAGATCATGGCCAGCAAAGAGCAGCAGATCCAGAGAGAC
GACCCTGGAGCCAGTCCCCAGAGCAGCAGCCAGCCAGACCACGGCCGCCTCTCCCCCCCA
GAAGCTCCCGACAGGCCACCATCTCCACGGCCTCCGAGACCTCAGTGTACGTGACCTGG
ATTCCCCGTGGGAATGGTGGGTTCCCAATCCAGTCCTTCCGTGTGGAGTACAAGAAGCTA
AAGAAAGTGGGAGACTGGATTCTGGCCACCAGCGCCATCCCCCATCGCGGCTGTCCGTG
GAGATCACGGGCCTAGAGAAAGGCACCTCCTACAAGTTTCGAGTCCGGGCTCTGAACATG
CTGGGGGAGAGCGAGCCAGCGCCCCCTCTCGGCCCTACGTGGTGTGCGGCTACAGCGGT
CGCGTGTACGAGAGGCCCCGTGGCAGGTCTTATATCACCTTACGGATGCGGTCAATGAG
ACCACCATCATGAGTCAAGTGGATGTACATCCCAGCAAGTAACAACAACACCCCAATCCAT
GGCTTTTATATCTATTATCGACCCACAGACAGTGAACAATGATAGTGACTACAAGAAGGAT
ATGGTGAAGGGGACAAGTACTGGCACTCCATCAGCCACCTGCAGCCAGAGACCTCCTAC
GACATTAAGATGCAGTGCTTCAATGAAGGAGGGGAGAGCGAGTTCAGCAACGTGATGATC
TGTGAGACCAAAGCTCGGAAGTCTTCTGGCCAGCCTGGTCGACTGCCACCCCCAACTCTG
GCCCCACCACAGCCGCCCTTCTGAAACCATAGAGCGGCCGGTGGGCACTGGGGCCATG
GTGGCTCGCTCCAGCGACCTGCCCTATCTGATTGTGCGGGTCTGCTGGGCTCCATCGTT
CTCATCATCGTCACCTTCATCCCCTTCTGCTTGTGGAGGGCCTGGTCTAAGCAAAAACAT
ACAACAGACCTGGGTTTTCTCGAAGTGCCCTTCCACCTCCTGCCCCGTATACTATGGTG
CCATTGGGAGGACTCCAGGCCACCAGGCCAGTGGACAGCCCTACCTCAGTGGCATCAGT

164/392

GGACGGGCCTGTGCTAATGGGATCCACATGAATAGGGGCTGCCCCTCGGCTGCAGTGGGC
TACCCGGGCATGAAGCCCCAGCAGCACTGCCCAGGCGAGCTTCAGCAGCAGAGTGACACC
AGCAGCCTGCTGAGGCAGACCCATCTTGGCAATGGATATGACCCCCAAAGTCACCAGATC
ACGAGGGGTCCCAAGTCTAGCCCGGACGAGGGCTCTTTCTTATACACACTGCCCGACGAC
TCCACTCACCAGCTGCTGCAGCCCCATCACGACTGCTGCCAACGCCAGGAGCAGCCTGCT
GCTGTGGGCCAGTCAGGGGTGAGGAGAGCCCCGACAGTCCTGTCTGGAAGCAGTGTGG
GACCTCCATTTCACTCAGGGCCCCCATGCTGCTTGGGCCTTGTGCCAGTTGAAGAGGTG
GACAGTCCTGACTCCTGCCAAGTGAGTGGAGGAGACTGGTGTCCCCAGCACCCCGTAGGG
GCCTACGTAGGACAGGAACCTGGAATGCAGCTCTCCCCGGGGGCCACTGGTGCGTGTGTCT
TTTGAACACACCCTCTCACAATTTAGGCAGAAGCTGATATCCCAGAAAGACTATATATT
GTTTTTTTTTTTAAAAAAAAAAGAAGAAAAAAGAGACAGAGAAAATTGGTATTTATTTTC
TATTATAGCCATATTTATATATTTATGCACTTGTAATAAATGTATATGTTTTATAATTC
TGGAGAGACATAAGGAGTCCTACCCGTTGAGGTTGGAGAGGGAAAATAAAGAAGCTGCCA
CCTAACAGGAGTCACCCAGGAAAGCACCGCACAGGCTGGCGCGGGACAGACTCCTAACCT
GGGGCCTCTGCAGTGGCAGGCGAGGCTGCAGGAGGCCACAGATAAGCTGGCAAGAGGAA
GGATCCCAGGCACATGGTTCATCACGAGCATGAGGGAACAGCAAGGGGCACGGTATCACA
GCCTGGAGACACCCACACAGATGGCTGGATCCGGTGCTACGGGAAACATTTTCCTAAGAT
GCCCATGAGAACAGACCAAGATGTGTACAGCACTATGAGCATTAACCACTTCCAGAAT
CAATAATCCGTGGCAACATATCTCTGTAAAAACAAACACTGTAACCTCTAAATAAATGTT
TAGTCTTCCCTGTAAAA

165/392

FIGURE 158

MLRGTMTAWRGMRPEVTLACLLLLATAGCFADLNEVPQVTVQPASTVQKPGGTIVILGCVVE
PPRMNVTWRLNGKELNGSDDALGVLI THGTLVITALLNNHTVGRYQCVARMPAGAVASVPA
TVTLANLQDFKLDVQHVIEVDEGNTAVIACHLPESHKPAQVRYSVKQEWLEASRGNYLIM
PSGNLQIVNASQEDEGMYKCAAYNPVTQEVKTSGSSDRLRVRRSTAEAAARI IYPPEAQTI
IVTKGQSLILECVASGIPPPRVTWAKDGSSVTGYNKTRFLLSNLLIDTTSEEDSGTYRCM
ADNGVGQPGA AVILYNVQVFEPPEVTMELS QLVIPWGQSAKLTCEVRGNPPPSVLWLRNA
VPLISSQRLRLSRRALRVLSMGPEDEGVYQCAENEVGSAAHVVQLRTSRPSITPRLWQD
AELATGTPPVSPSKLGNPEQMLRGQPALPRPPTSVGPASPKCPGEGKGQAGAPAEAPIILSS
PRTSKTDSYELVWRPRHEGSGRAPILYYVVKHRKQVTNSSDDWTISGIPANQHRLTLTRL
DPGSLYEVEMAAYNCAGEGQTAMVTFRTGRRPKPEIMASKEQQIQRDDPGAS PQSSSQPD
HGRLSPPEAPDRPTISTASETSVYVTWIPRGNGGFPIQSFRVEYKKLKKVGDWILATSAI
PPSRLSVEITGLEKGT SYKFRVRALNMLGESEPSAPSRPYVVSGYSGRVYERP VAGPYIT
FTDAVNETTIMLKWMIIPASNNNTPIHGFYIYYRPTSDSDNDSYKKDMVEGD KYWHSISH
LQPETSYDIKMQCFNEGGESEFSNVMICETKARKSSGQPGRLPPPTLAPPQPPLPETIER
PVG TGAMVARSSDLPYLIVGVVLGSI VLIIVTFIPFCLWRAW SKQKHTTDLGFPR SALPP
SCPYTMVPLGGLPGHQASGQPYLSGISGRACANGIHMNRGCPSAAVGYPGMKPQQHCPGE
LQQQSDTSSLLRQTHLGNGYDPQSHQITRGPKSSPDEGSFLYTL PDDSTHQLLQPHHDCC
QRQE QPAAVGQSGVRRAPDSPVLEAVWDPPFHSGPPCCLGLVPVEEVDSPDSCQVSGGDW
CPQHVPVGAYVQGEPGMQLSPGPLVRVSFETPPLTI

Signal peptide:
amino acids 1-30

Transmembrane domain:
amino acids 16-30 (type II), 854-879

166/392

FIGURE 159

CCCACGCGTCCGCCCACGCGTCCGCCCACGCGTCCGCCCACGCGTCCGCGTCCG
CCCACGCGTCCGCCCACGCGTCCGGTGCAAGCTCGCGCCGCACACTGCCTGGTGGAGGGA
AGGAGCCCCGGGCGCCTCTCGCCGCTCCCCGCGCCGCGTCCGCACCTCCCCACCGCCCGC
CGCCCCGCGCCCGCCGCCCCGCAAAGCATGAGTGAGCCCGCTCTCTGCAGCTGCCCGGGGC
GCGAATGGCAGGCTGTTTCCGCGGAGTAAAGGTGGCGCCGGTCAGTGGTCGTTTCCAAT
GACGGACATTAACCAGACTGTCAGATCCTGGGGAGTCGCGAGCCCCGAGTTTGGAGTTTT
TTCCCCCCCACAACGTACAGTCCGAAGTGCAGAGGGAAAGGAAGGCGGCAGGAAGGCGAA
GCTCGGGCTCCGGCACGTAGTTGGGAAACTTGCGGGTCTAGAAAGTCGCCTCCCCGCCTT
GCCGGCCGCCCTTGACAGCCCCGAGCCGAGCAGCAAAGTGAGACATTGTGCGCTGCCAGA
TCCGCCGGCCCGCGGACCGGGGCTGCCTCGGAAACACAGAGGGGTCTTCTCTCGCCCTGCA
TATAATTAGCCTGCACACAAAGGGAGCAGCTGAATGGAGGTTGTCACTCTCTGGAAGG
ATTTCTGACCGAGCGCTTCCAATGGACATTCTCCAGTCTCTCTGGAAGATTCTCGCTAA
TGGATTTCTCTGCTGCTCGGTCTCTGTCTATACTGGCTGCTGAGGAGGCCCTCGGGGGTGG
TCTTGTGTCTGCTGGGGGCTGCTTTTCAAGTGTGCTGCGCCCGCCCGCCAGCGGTGCCCGC
AGCTGTGCCGGTGCGAGGGGCGGTGCTGTACTGCGAGGCGCTCAACCTCACCGAGGCGC
CCCACAACCTGTCCGGCCTGCTGGGCTTGTCCCTGCGCTACAACAGCCTCTCGGAGCTGC
GCGCCGGCCAGTTCACGGGGTTAATGCAGCTCACGTGGCTCTATCTGGATCACAATCACA
TCTGCTCCGTGCAGGGGGACGCCTTTAGAACTGCGCCGAGTTAAGGAACCTCACGTGA
GTTCCAACCAGATCACCCAACCTGCCAACACCACTTCCGGCCCATGCCAACCTGCGCA
GCGTGACCTCTCGTACAACAAGCTGCAGGCGCTCGCGCCCGACCTCTTCCACGGGCTGC
GGAAGCTCACACGCTGCATATGCGGGCCAACGCCATCCAGTTTGTGCCCCGTGCGCATCT
TCCAGGACTGCCGAGCCTCAAGTTTCTCGACATCGGATACAATCAGCTCAAGAGTCTGG
CGCGCAACTCTTTCGCCGGCTTGTTTAAAGCTCACCGAGCTGCACCTCGAGCACAACGACT
TGGTCAAGGTGAACCTCGCCCACTTCCCGCGCCTCATCTCCCTGCACTCGCTCTGCCTGC
GGAGGAACAAGGTGGCCATTGTGGTCAGCTCGCTGGACTGGGTTTGGAACTGGAGAAAA
TGGACTTGTCGGGCAACGAGATCGAGTACATGGAGCCCCATGTGTTGAGACCGTGCCGC
ACCTGCAGTCCCTGCAGCTGGACTCCAACCGCCTCACCTACATCGAGCCCCGGATCCTCA
ACTCTTGGAAGTCCCTGACAAGCATCACCTGGCCGGGAACCTGTGGGATTGCGGGCGCA
ACGTGTGTGCCCTAGCCTCGTGGCTCAGCAACTTCCAGGGGCGCTACGATGGCAACTTGC
AGTGCGCCAGCCCGGAGTACGCAACAGGGCGAGGACGTCTTGACGCGGTGTACGCCTTCC
ACCTGTGCGAGGATGGGGCCGAGCCCAACAGCGGCCACCTGCTCTCGGCCGTACCAACC
GCAGTGATCTGGGGCCCCCTGCCAGCTCGGCCACCAGCTCGCGGACGGCGGGGAGGGGC
AGCACGACGGCACATTTCGAGCCTGCCACCGTGGCTCTTCCAGGCGGCGAGCACGCCGAGA
ACGCCGTGCAGATCCACAAGGTGGTCACGGGCACCATGGCCCTCATCTTCTCCTTCCTCA
TCGTGGTCTTGGTGCTCTACGTGTCTTGGAAAGTGTTCCTCAGCCAGCCTCAGGCAGCTCA
GACAGTGCTTTGTACGCGAGCGCAGGAAGCAAAAGCAGAAACAGACCATGCATCAGATGG
CTGCCATGTCTGCCAGGAATACTACGTTGATTACAAACCGAACCACATTGAGGGAGCCC
TGGTGATCATCAACGAGTATGGCTCGTGTACCTGCCACCAGCAGCCCGCGAGGGAATGCG
AGGTGTGAATTGTCCAGTGGCTCTCAACCCATGCGCTACCAAATACGCCTGGGCAGCCGG
GACGGGCCGGCGGGCACCAGGCTGGGGTCTCCTTGTCTGTGCTCTGATATGCTCCTTGAC
TGAAACTTTAAGGGGATCTCTCCAGAGACTTGACATTTTAGCTTTATTGTGTCTTAAAA
ACAAAAGCGAATTAAAAACACAACAAAAAACCCACCCCAACCTTCAGGACAGTCTATC
TTAAATTTTCATATGAGAACTCCTTCCTCCCTTTGAAGATCTGTCCATATTAGGAATCTG
AGAGTGTAAGAAAGGTGGCCATAAGACAGAGAGAGAATAATCGTGCTTTGTTTTATGCTA
CTCCTCCACCCCTGCCCATGATTAAACATCATGTATGTAGAAGATCTTAAGTCCATACGC
ATTTTCATGAAGAACCATTGGAAAGAGGAATCTGCAATCTGGGAGCTTAAGAGCAAATGAT

167/392

GACCATAGAAAGCTATGTTCTTACTTTGTGTGTGTGTCTGTATGTTTCTGCGTTGTGTGT
CTTTGTAGGCAAGCAAACGTTGTCTACACAAACGGGAATTTAGCTCACATCATTTCATGC
CCCTGTGCCTCTAGCTCTGGAGATTGGTGGGGGGAGGTGGGGGGAAACGGCAGGAATAAG
GGAAAGTGGTAGTTTTTAAGGTTTGTAACTTGAATCTTTTCTTCTCAAATTA
ATTATCTTTAAGCTTCAAGAACTTGCTCTGACCCCTCTAAGCAAACCTACTAAGCATTTA
AAAGAGAATCTAATTTTTTAAAGGTGTAGCACCTTTTTTTTTTATTCTTCCACAGAGGGTG
CTAATCTCATTATGCTGTGCTATCTGAAAAGAACTTAAGGCCACAATTCACGTCTCGTCC
TGGGCATTGTGATGGATTGACCCTCCATTTGCAGTACCTTCCCAGCTGATTAAAGTTCAG
CAGTGGTATTGAGGTTTTTCGAATATTTATATAGAAAAAAGTCTTTTCACATGACAAAT
GACACTCTCACACCAGTCTTAGCCCTAGTAGTTTTTTTAGGTTGGACCAGAGGAAGCAGGT
TAAATGAGACCTGTCCTCTGCTGCACTCAGAAAAAATAGGCAGTCCCTGATGCTCAGATC
TTAGCCTTGATATTAATAGTTGAGACCACCTACCCACAATGCAGCCTATACTCCCAAGAC
TACAAAGTTACCATCGCAAAGGAAAGGTTATTCCAGTAAAAGGAAATAGTTTTCTCAACC
ATTTAAAAATATTCTTCTGAACTCATCAAAGTAGAAGAGCCCCCAACCTTTTCTCTCTGC
CTTCAAGAAGGCAGACATTTGGTATGATTTAGCATCAACAACACATTTATGAGTATATGT
AAGTAATCAGAGGGGCAAATGCCACTTGTTATTCTTCCCAAGTTTTCCAAGCAAGTACAC
ACAGATCTCTGGTAGGATTAGGGGCCACTTGTTGTTTCCGGCTTATTTTAGTCGACTTGTC
AGCAAGTTTGATGCCTAGTCTATCTGACATGGCCCAGTAGAACAGGGCATTGATGGATCA
CATGAGATGGTAGAAGGAACATCATCACATACCCCTCTCACAGAGAAAATTATCAAAGAA
CCAGAAATTATATCTGTTTTGGAGCAAGAGTGTATAATGTTTCAGGGTAGTCAAATAA
ACATAAATTATCTCCTCTAGATGAGTGGCGATGTTGGCTGATTTGGGTCTGCCATTGACA
GAATGTCAAATAAAAAGGAATTAGCTAGAATATGACCATTAAATGTGCTTCTGAAATATA
TTTTGAGATAGTTTTAGAATGTCA

168/392

FIGURE 160

MDFLLLGLCLYWLLRRPSGVVLCLLGACFQMLPAAPSGCPQLCRCEGRLLYCEALNLTEA
PHNLSGLLGLSLRYNSLSELRAGQFTGLMQLTWLYLDHNHICSVQGDADFQKLRRVKELTL
SSNQITQLPNTTFRPMPNLRSDLSYNKLQALAPDLFHGLRKLTTLHMRANAIQFVPVRI
FQDCRSKFLDIGYNQLKSLARNSFAGLFLKTELHLEHNDLVKVNFAHFPRILSLHSLCL
RRNKVAIVVSSLDWVWNLEKMDLSGNEIEYMEPHVFETVPHLQSLQLDSNRLTYIEPRIL
NSWKSLSITLAGNLWDCGRNVCALASWLSNFQGRYDGNLQCASPEYAQGEDVLDVYAF
HLCEDGAEP TSGHLLSAVTNRSDLGPPASSATT LADGGEGQHDGTFEPATVALPGGEHAE
NAVQIHKVVTGTMALIFSF L I V V L V L Y V S W K C F P A S L R Q L R Q C F V T Q R R K Q K Q K T M H Q M
AAMSAQEY Y V D Y K P N H I E G A L V I I N E Y G S C T C H Q Q P A R E C E V

170/392

FIGURE 162

MAPQSLPSSRMAPLGMLLGLLMAACFTFCLSHQNLKEFALTNPEKSSTKETERKETKAEE
ELDAEVLEV FHP THEWQALQPGQAVPAGSHVRLNLQTGEREAKLOYEDKFRNNLKGKRLD
INTNTYTSQDLKSALAKFKEGAEMESSKEDKARQAEVKRLFRPIEELKKDFDELNVVIET
DMQIMVRLINKFNSSSSSLEEKIAALFDLEYVYVHQMDNAQDLLSFGGLQVVINGLNSTEP
LVKEYAAFVLGAAFSSNPKVQVEAIEGGALQKLLVILATEQPLTAKKKVLFALCSLLRHF
PYAQRQFLKLGGGLQVLR TLVQEKGTEVLAVRVVTL LYDLVTEKMF AEEEEAE LTQEMSPEK
LQQYRQVHLLPGLWEQGWCEITAHLLALPEHDAREKVLQTLGVLLTTCDRYRQDPQLGR
TLASLQAEYQVLASLELQDGEDEGYFQELLGSVNSLLKELR

Important features:

Signal peptide:

amino acids 1-29

Hypothetical YJL126w/YLR351c/yhcX family protein:

amino acids 364-373

N-glycosylation site:

amino acids 193-197, 236-240

N-myristoylation site:

amino acids 15-21, 19-25, 234-240, 251-257, 402-408, 451-457

Homologous region SLS1 protein:

amino acids 68-340

171/392

FIGURE 163

CAGAGAGGAGGCTTTGGGAATTGTCCAGCAGAAACAGAGAAGTCTGAGGTGGTGTCAAGA
CAAAAGATGCTTCAGCTTTGGAACTTGTTCTCCTGTGCGGCGTGCTCACTGGGACCTCA
GAGTCTCTTCTTGACAATCTTGGCAATGACCTAAGCAATGTCGTGGATAAGCTGGAACCT
GTTCTTCACGAGGGACTTGAGACAGTTGACAATACTCTTAAAGGCATCCTTGAGAACTG
AAGGTCGACCTAGGAGTGCTTCAGAAATCCAGTGCTTGGCAACTGGCCAAGCAGAAGGCC
CAGGAAGCTGAGAAATTGCTGAACAATGTCATTTCTAAGCTGCTTCCAACCTAACACGGAC
ATTTTTGGGTTGAAAATCAGCAACTCCCTCATCCTGGATGTCAAAGCTGAACCGATCGAT
GATGGCAAAGGCCTTAACCTGAGCTTCCCTGTCAACGCGAATGTCACTGTGGCCGGGCCC
ATCATTGGCCAGATTATCAACCTGAAAGCCTCCTTGGACCTCCTGACCGCAGTCACAATT
GAAACTGATCCCCAGACACACCAGCCTGTTGCCGTCTGGGAGAATGCGCCAGTGACCCA
ACCAGCATCTCACTTTCCTTGCTGGACAAACACAGCCAAATCATCAACAAGTTCGTGAAT
AGCGTGATCAACACGCTGAAAAGCACTGTATCCTCCCTGCTGCAGAAGGAGATATGTCCA
CTGATCCGCATCTTCATCCACTCCCTGGATGTGAATGTCATTTCAGCAGGTTCGTGATAAT
CCTCAGCACAAAACCCAGCTGCAAACCCTCATCTGAAGAGGACGAATGAGGAGGACCACT
GTGGTGATGCTGATTGGTTCCAGTGCTTGCCCCACCCCCTTATAGCATCTCCCTCCA
GGAAGCTGCTGCCACCACCTAACCAGCGTGAAAGCCTGAGTCCCACCAGAAGGACCTTCC
CAGATACCCCTTCTCCTCACAGTCAGAACAGCAGCCTCTACACATGTTGTCTGCCCCTG
GCAATAAAGGCCCATTTCTGCACCCTTAA

172/392

FIGURE 164

MLQLWKLVLLCGVLTGTSESLLDNLGNDLSNVVDKLEPVLHEGLETVDNTLKGILEKLKV
DLGVLQKSSAWQLAKQKAQEA EKLLNNVISKLLPTNTDIFGLKISNSLI LDVKAEPIDDG
KGLNLSFPVTANVTVAGPIIGQIINLKASDLLTAVTIETDPQTHQPVAVLGECASDPTS
ISLSLLDKHSQIINKFVNSVINTLKSTVSSLLQKEICPLIRIFIHSLDVNVIQQVVDNPQ
HKTQLQTLI

Important features:**Signal peptide:**

1-15

Transmembrane domain:

none

N-glycosylation site:

124-128, 132-136

N-myristoylation site:

12-18, 16-22, 26-32, 101-107, 122-128, 141-147

Leucine zipper pattern:

44-66

173/392

FIGURE 165

GCAGTCAGAGACTTCCCCTGCCCCCTCGCTGGGAAAGAACATTAGGAATGCCTTTTGTAGTGC
CTTGCTTTCCTGAACTAGCTCACAGTAGCCCGGCGGCCAGGGCAATCCGACCACATTTCA
CTCTCACCGCTGTAGGAATCCAGATGTCAGGCCAAGTACAGCAGCACGAGGGACATGCTGG
ATGATGATGGGGACACCACCATGAGCCTGCATTCTCAAGCCTCTGCCACAACCTCGGCATC
CAGAGCCCCGGCGCACAGAGCACAGGGCTCCCTCTTCAACGTGGCGACCAGTGGCCCTGA
CCCTGCTGACTTTGTGCTTGGTGCTGCTGATAGGGCTGGCAGCCCTGGGGCTTTTGTTTT
TTCAGTACTACCAGCTCTCCAATACTGGTCAAGACACCATTTCTCAAATGGAAGAAAGAT
TAGGAAATACGTCCCAAGAGTTGCAATCTCTTCAAGTCCAGAATATAAAGCTTGCAGGAA
GTCTGCAGCATGTGGCTGAAAACTCTGTCTGAGCTGTATAACAAAGCTGGAGCACACA
GGTGCAGCCCTTGTACAGAACAATGGAAATGGCATGGAGACAATTGCTACCAGTTCTATA
AAGACAGCAAAAGTTGGGAGGACTGTAAATATTTCTGCCTTAGTGAAAACCTTACCATGC
TGAAGATAAACAAACAAGAAGACCTGGAATTTGCCGCGTCTCAGAGCTACTCTGAGTTTT
TCTACTCTTATTGGACAGGGCTTTTGCGCCCTGACAGTGGCAAGGCCTGGCTGTGGATGG
ATGGAACCCCTTTCACTTCTGAACTGTTCCATATTATAATAGATGTCACCAGCCCAAGAA
GCAGAGACTGTGTGGCCATCCTCAATGGGATGATCTTCTCAAAGGACTGCAAAGAATTGA
AGCGTTGTGTCTGTGAGAGAAGGGCAGGAATGGTGAAGCCAGAGAGCCTCCATGTCCCCC
CTGAAACATTAGGCGAAGGTGACTGATTTCGCCCTCTGCAACTACAAATAGCAGAGTGAGC
CAGGCGGTGCCAAAGCAAGGGCTAGTTGAGACATTGGGAAATGGAACATAATCAGGAAAG
ACTATCTCTCTGACTAGTACAAAATGGGTCTCTCGTGTTTCCTGTTTCAGGATCACCAGCAT
TTCTGAGCTTGGGTTTTATGCACGTATTTAACAGTCAACAAGAAGTCTTATTTACATGCCAC
CAACCAACCTCAGAAACCCATAATGTCATCTGCCTTCTTGGCTTAGAGATAACTTTTAGC
TCTCTTTCTTCTCAATGTCTAATATCACCTCCCTGTTTTTCATGTCTTCCTTACACTTGGT
GGAATAAGAACTTTTTTGAAGTAGAGGAAATACATTGAGGTAACATCCTTTTCTCTGACA
GTCAAGTAGTCCATCAGAAATTGGCAGTCACCTCCAGATTGTACCAGCAAATACACAAG
GAATTCTTTTTTGTGTTTTCAGTTCATACTAGTCCCTTCCCAATCCATCAGTAAAGACCC
CATCTGCCTTGTCCATGCCGTTTCCCAACAGGGATGTCACTTGATATGAGAATCTCAAAT
CTCAATGCCTTATAAGCATTCCTTCTGTGTCCATTAAGACTCTGATAATTGTCTCCCCCT
CCATAGGAATTTCTCCAGGAAAGAAATATATCCCATCTCCGTTTCATATCAGAACTAC
CGTCCCCGATATTCCCTTCAGAGAGATTAAAGACCAGAAAAAAGTGAGCCTCTTCATCTG
CACCTGTAATAGTTTCAGTTCCTATTTTCTTCCATTGACCCATATTTATACCTTTCAGGT
ACTGAAGATTTAATAATAATAAATGTAAATACTGTGAAAAA

174/392

FIGURE 166

MQAKYSSTRDMLDDDGDTTMSLHSQASATTRHPEPRRTEHRAPSSTWRPVALTLLTLCLV
LLIGLAALGLLFFQYYQLSNTGQDTISQMEERLGNTSQELQSLQVQNIKLAGSLQHVAEK
LCRELYNKAGAHRCSPCTEQWKWHGDNCYQFYKDSKSWEDCKYFCLSENSTMLKINKQED
LEFAASQSYSEFFYSYWTGLLRPD SGKAWLWMDGTPFTSELFHIIIDVTSPRSRDCVAIL
NGMIFSKDCKELKRCVCERRAGMVKPESLHVPPETLGED

175/392

FIGURE 167

CGGACGGGCAGGACGCCCCGTTTCGCCTAGCGCGTGCTCAGGAGTTGGTGTCTGCCTGCG
CTCAGGATGAGGGGGAATCTGGCCCTGGTGGGCGTTCTAATCAGCCTGGCCTTCCTGTCA
CTGCTGCCATCTGGACATCCTCAGCCGGCTGGCGATGACGCCTGCTCTGTGCAGATCCTC
GTCCCTGGCCTCAAAGGGGATGCGGGAGAGAAGGGAGACAAAGGCGCCCCCGGACGGCCT
GGAAGAGTCGGCCCCACGGGAGAAAAAGGAGACATGGGGGACAAAGGACAGAAAGGCAGT
GTGGGTCGTTCATGGAAAAATTGGTCCCATTGGCTCTAAAGGTGAGAAAGGAGATTCCGGT
GACATAGGACCCCCTGGTCCTAATGGAGAACCAGGCCTCCCATGTGAGTGCAGCCAGCTG
CGCAAGGCCATCGGGGAGATGGACAACCAGGTCTCTCAGCTGACCAGCGAGCTCAAGTTC
ATCAAGAATGCTGTGCGCCGGTGTGCGCGAGACGGAGAGCAAGATCTACCTGCTGGTGAAG
GAGGAGAAGCGCTACGCGGACGCCCAGCTGTCCTGCCAGGGCCGCGGGGGCACGCTGAGC
ATGCCCAAGGACGAGGCTGCCAATGGCCTGATGGCCGCATACCTGGCGCAAGCCGGCCTG
GCCCCTGTCTTCATCGGCATCAACGACCTGGAGAAGGAGGGCGCCTTCGTGTACTCTGAC
CACTCCCCCATGCGGACCTTCAACAAGTGGCGCAGCGGTGAGCCCAACAATGCCTACGAC
GAGGAGGACTGCGTGGAGATGGTGGCCTCGGGCGGCTGGAACGACGTGGCCTGCCACACC
ACCATGTACTTCATGTGTGAGTTTGACAAGGAGAACATGTGAGCCTCAGGCTGGGGCTGC
CCATTGGGGGCCCCACATGTCCCTGCAGGGTTGGCAGGGACAGAGCCCAGACCATGGTGC
CAGCCAGGGAGCTGTCCCTCTGTGAAGGGTGGAGGCTCACTGAGTAGAGGGCTGTTGTCT
AAACTGAGAAAATGGCCTATGCTTAAGAGGAAAATGAAAGTGTTCCCTGGGGTGCTGTCTC
TGAAGAAGCAGAGTTTCATTACCTGTATTGTAGCCCCAATGTCATTATGTAATTATTACC
CAGAATTGCTCTTCCATAAAGCTTGTGCCTTTGTCCAAGCTATACAATAAAATCTTTAAG
TAGTGCAGTAGTTAAGTCCAAAAAAAAAAAAAAAAAAAAA

176/392

FIGURE 168

MRGNLALVGVLISLAFLSLLPSGHPQPAGDDACSVQILVPGLKGDAGEKGDKGAPGRPGR
VGPTGEKGDMDKGQKGSVGRHGKIGPIGSKGEKGDSDIGPPGPNGEPGLPCECSQLRK
AIGEMDNQVSQLTSELKFIKNAVAGVRETESKIYLLVKEEKRYADAQLSCQGRGGTLSMP
KDEAANGLMAAYLAQAGLARVFIGINDLEKEGAFVYSDHSPMRTFNKWRSGEPNNAYDEE
DCVEMVASGGWNDVACHTTMYFMCEFDKENM

177/392

FIGURE 169

AGTGACTGCAGCCTTCCTAGATCCCCTCCACTCGGTTTCTCTCTTTGCAGGAGCACCGGC
AGCACCAGTGTGTGAGGGGAGCAGGCAGCGTCTAGCCAGTTCCTTGATCCTGCCAGAC
CAGCCAGCCCCCGGCACAGAGCTGCTCCACAGGCACCAATGAGGATCATGCTGCTATTAC
AGCCATCCTGGCCTTCAGCCTAGCTCAGAGCTTTGGGGCTGTCTGTAAGGAGCCACAGGA
GGAGGTGGTTCTTGGCGGGGGCCGCAGCAAGAGGGATCCAGATCTCTACCAGCTGCTCCA
GAGACTCTTCAAAAGCCACTCATCTCTGGAGGGATTGCTCAAAGCCCTGAGCCAGGCTAG
CACAGATCCTAAGGAATCAACATCTCCCGAGAAACGTGACATGCATGACTTCTTTGTGGG
ACTTATGGGCAAGAGGAGCGTCCAGCCAGAGGGAAAGACAGGACCTTTCTTACCTTCAGT
GAGGGTTCCTCGGCCCCCTTCATCCCAATCAGCTTGGATCCACAGGAAAGTCTTCCCTGGG
AACAGAGGAGCAGAGACCTTTATAAGACTCTCCTACGGATGTGAATCAAGAGAACGTCCC
CAGCTTTGGCATCCTCAAGTATCCCCCGAGAGCAGAATAGGTACTCCACTTCCGGAATCC
TGGACTGCATTAGGAAGACCTCTTCCCTGTCCCAATCCCCAGGTGCGCACGCTCCTGTT
ACCCTTTCTCTTCCCTGTTCTTGTAACATTCTTGTGCTTTGACTCCTTCTCCATCTTTTC
TACCTGACCCTGGTGTGGAACTGCATAGTGAATATCCCCAACCCCAATGGGCATTGACT
GTAGAATACCCTAGAGTTCCTGTAGTGTCTACATTAAAAATATAATGTCTCTCTCTATT
CCTCAACAATAAAGGATTTTTGCATATGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAA

178/392

FIGURE 170

MRIMLLFTAILAFSLAQSFQAVCKEPQEEVVPGGGRSKRDPDLYQLLQRLFKSHSSLEGL
LKALSQASTDPKESTSPEKRDMDFFVGLMGKRSVQPEGKTGPFLPSVRVPRPLHPNQLG
STGKSSLGTEEQRPL

Important features:

Signal peptide:

amino acids 1-18

Tyrosine kinase phosphorylation site:

amino acids 36-45

N-myristoylation site:

amino acids 33-39, 59-65

Amidation site:

amino acids 90-94

Leucine zipper pattern:

amino acids 43-65

Tachykinin family signature:

amino acids 86-92

179/392

FIGURE 171

TGGCCTCCCCAGCTTGCCAGGCACAAGGCTGAGCGGGAGGAAGCGAGAGGCATCTAAGCA
GGCAGTGTTTTGCCTTCACCCCAAGTGACCAATGAGAGGTGCCACGCGAGTCTCAATCATG
CTCCTCCTAGTAACTGTGTCTGACTGTGCTGTGATCACAGGGGCCTGTGAGCGGGATGTC
CAGTGTGGGGCAGGCACCTGCTGTGCCATCAGCCTGTGGCTTCGAGGGCTGCGGATGTGC
ACCCCGCTGGGGCGGGAAGGCGAGGAGTGCCACCCCGGCAGCCACAAGGTCCCCTTCTTC
AGGAAACGCAAGCACACACCTGTCCTTGCTTGCCCAACCTGCTGTGCTCCAGGTTCCCG
GACGGCAGGTACCGCTGCTCCATGGACTTGAAGAACATCAATTTTTTAGGCGCTTGCCTGG
TCTCAGGATACCCACCATCCTTTTCCTGAGCACAGCCTGGATTTTTATTTCTGCCATGAA
ACCCAGCTCCCATGACTCTCCAGTCCCTACACTGACTACCCTGATCTCTCTTGTCTAGT
ACGCACATATGCACACAGGCAGACATACTCCCATCATGACATGGTCCCCAGGCTGGCCT
GAGGATGTACAGCTTGAGGCTGTGGTGTGAAAGGTGGCCAGCCTGGTTCTCTTCCCTGC
TCAGGCTGCCAGAGAGGTGGTAAATGGCAGAAAGGACATTCCCCCTCCCCTCCCCAGGTG
ACCTGCTCTCTTTCTGGGCCCTGCCCTCTCCCCACATGTATCCCTCGGTCTGAATTAG
ACATTCTGGGCACAGGCTCTTGGGTGCATTGCTCAGAGTCCCAGGTCCTGGCCTGACCC
TCAGGCCCTTCACGTGAGGTCTGTGAGGACCAATTTGTGGGTAGTTCATCTTCCCTCGAT
TGGTTAACTCCTTAGTTTCAGACCACAGACTCAAGATTGGCTCTTCCCAGAGGGCAGCAG
ACAGTCACCCCAAGGCAGGTGTAGGGAGCCCAGGGAGGCCAATCAGCCCCCTGAAGACTC
TGGTCCCAGTCAGCCTGTGGCTTGTGGCCTGTGACCTGTGACCTTCTGCCAGAATTGTCA
TGCCTCTGAGGCCCCCTCTTACCACACTTTACCAGTTAACCCTGAAGCCCCCAATTCCC
ACAGCTTTTCCATTAAAATGCAAATGGTGGTGGTTCAATCTAATCTGATATTGACATATT
AGAAGGCAATTAGGGTGTTTCCTTAAACAACCTCTTTCCAAGGATCAGCCCTGAGAGCAG
GTTGGTGACTTTGAGGAGGGCAGTCCTCTGTCCAGATTGGGGTGGGAGCAAGGGACAGGG
AGCAGGGCAGGGGCTGAAAGGGGCACCTGATTGAGACCAGGGAGGCAACTACACACCAACA
TGCTGGCTTTAGAATAAAAGCACCAACTGAAAAA

180/392

FIGURE 172

MRGATRVSIMLLLVTVSDCAVITGACERDVQCGAGTCCAISLWLRGLRMCTPLGREGEEC
HPGSHKVPFFRKRKHHTCPCLPNLLCSRFPDGRYRCSMDLKNINF

Signal peptide:
amino acids 1-19

Tyrosine kinase phosphorylation site:
amino acids 88-95

N-myristoylation sites:
amino acids 33-39, 35-41, 46-52

181/392

FIGURE 173

AGCGCCCGGGCGTCGGGGCGGTAAAAGGCCGGCAGAAGGGAGGCACTTGAGAAATGTCTT
TCCTCCAGGACCCAAGTTTCTTCACCATGGGGATGTGGTCCATTGGTGCAGGAGCCCTGG
GGGCTGCTGCCTTGGCATTGCTGCTTGCCAACACAGACGTGTTTCTGTCCAAGCCCCAGA
AAGCGGCCCTGGAGTACCTGGAGGATATAGACCTGAAAACACTGGAGAAGGAACCAAGGA
CTTTCAAAGCAAAGGAGCTATGGGAAAAAAATGGAGCTGTGATTATGGCCGTGCGGAGGC
CAGGCTGTTTCTCTGTCTGAGAGGAAGCTGCGGATCTGTCTCCCTGAAAAGCATGTTGG
ACCAGCTGGGCGTCCCCCTCTATGCAGTGGTAAAGGAGCACATCAGGACTGAAGTGAAGG
ATTTCCAGCCTTATTTCAAAGGAGAAATCTTCCTGGATGAAAAGAAAAAGTTCTATGGTC
CACAAAGGCGGAAGATGATGTTTATGGGATTTATCCGTCTGGGAGTGTGGTACAACCTTCT
TCCGAGCCTGGAACGGAGGCTTCTCTGGAACCTGGAAGGAGAAGGCTTCATCCTTGGGG
GAGTTTTCGTGGTGGGATCAGGAAAGCAGGGCATTCTTCTTGAGCACCGAGAAAAAGAAT
TTGGAGACAAAGTAAACCTACTTTCTGTTCTGGAAGCTGCTAAGATGATCAAACCAACAGA
CTTTGGCCTCAGAGAAAAAAATGATTTGTGTGAAACTGCCAGCTCAGGGATAACCAGGGAC
ATTCACCTGTGTTTCATGGGATGTATTGTTTCCACTCGTGTCCCTAAGGAGTGAGAAACCC
ATTTATACTCTACTCTCAGTATGGATTATTAATGTATTTTAATATTCTGTTTAGGCCAC
TAAGGCAAAATAGCCCCAAAACAAGACTGACAAAAATCTGAAAACTAATGAGGATTATT
AAGCTAAAACCTGGGAAATAGGAGGCTTAAAATTGACTGCCAGGCTGGGTGCAGTGGCTC
ACACCTGTAATCCCAGCACTTTGGGAGGCCAAGGTGAGCAAGTCACTTGAGGTCGGGAGT
TCGAGACCAGCCTGAGCAACATGGCGAAACCCCGTCTCTACTAAAAATACAAAAATCACC
CGGGTGTGGTGGCAGGCACCTGTAGTCCCAGCTACCCGGGAGGCTGAGGCAGGAGAATCA
CTTGAACCTGGGAGGTGGAGGTTGCGGTGAGCTGAGATCACACCACTGTATTCCAGCCTG
GGTGA CTGAGACTCTAACTAA

182/392

FIGURE 174

MSFLQDPSFFTGMWWSIGAGALGAAALALLANTDVFLSKPQKALEYLEDIDLKTLEKE
PRTFKAKELWEKNGAVIMAVRRPGCFLCREEAADLSSLKSMLDQLGVPLYAVVKEHIRTE
VKDFQPYFKGEIFLDEKKKFYGPQRRKMMFMGFIRLGWYNFFRAWNGGFSGNLEGEGLFI
LGGVFVVGSGKQGILLEHREKEFGDKVNLLSVLEAAKMIKPQTLASEKK

183/392

FIGURE 175

GACAGTGGAGGGCAGTGGAGAGGACCGCGCTGTCCTGCTGTCACCAAGAGCTGGAGACAC
CATCTCCCACCGAGAGTCAATGGCCCCATTGGCCCTGCACCTCCTCGTCCTCGTCCCCATC
CTCCTCAGCCTGGTGGCCTCCCAGGACTGGAAGGCTGAACGCAGCCAAGACCCCTTCGAG
AAATGCATGCAGGATCCTGACTATGAGCAGCTGCTCAAGGTGGTGACCTGGGGGCTCAAT
CGGACCCTGAAGCCCCAGAGGGTGATTGTGGTTGGCGCTGGTGTGGCCGGGCTGGTGGCC
GCCAAGGTGCTCAGCGATGCTGGACACAAGGTCACCATCCTGGAGGCAGATAACAGGATC
GGGGGCCGCATCTTACCTACCGGGACCAGAACACGGGCTGGATTGGGGAGCTGGGAGCC
ATGCGCATGCCCAGCTCTCACAGGATCCTCCACAAGCTCTGCCAGGGCCTGGGGCTCAAC
CTGACCAAGTTCACCCAGTACGACAAGAACACGTGGACGGAGGTGCACGAAGTGAAGCTG
CGCAACTATGTGGTGGAGAAGGTGCCCGAGAAGCTGGGCTACGCCTTGCGTCCCCAGGAA
AAGGGCCACTCGCCCGAAGACATCTACCAGATGGCTCTCAACCAGGGCCCTCAAAGACCTC
AAGGCACTGGGCTGCAGAAAGGCGATGAAGAAGTTTGAAAGGCACACGCTCTTGGAATAT
CTTCTCGGGGAGGGGAACCTGAGCCGGCCGGCCGTGCAGCTTCTGGGAGACGTGATGTCC
GAGGATGGCTTCTTCTATCTCAGCTTCGCCGAGGCCCTCCGGGCCCACAGCTGCCTCAGC
GACAGACTCCAGTACAGCCGCATCGTGGGTGGCTGGGACCTGCTGCCGCGCGCGCTGCTG
AGCTCGCTGTCCGGGCTTGTGCTGTTGAACGCGCCCCGTGGTGGCGATGACCCAGGGACCG
CACGATGTGCACGTGCAGATCGAGACCTCTCCCCCGCGCGGAATCTGAAGGTGCTGAAG
GCCGACGTGGTGCTGCTGACGGCGAGCGGACCGGCGGTGAAGCGCATCACCTTCTCGCCG
CCGCTGCCCCGCCACATGCAGGAGGCGCTGCGGAGGCTGCACTACGTGCCGGCCACCAAG
GTGTTCCCTAAGCTTCCGCAGGCCCTTCTGGCGCGAGGAGCACATTGAAGGCGGCCACTCA
AACACCGATCGCCCGTCGCGCATGATTTTCTACCCGCCGCCGCGCGAGGGCGCGCTGCTG
CTGGCCTCGTACACGTGGTTCGGACGCGGCGGCAGCGTTTCGCCGGCTTGAGCCGGGAAGAG
GCGTTGCGCTTGGCGCTCGACGACGTGGCGGCATTGCACGGGCCTGTCTGTGCGCCAGCTC
TGGGACGGCACCCGGCGTTCGTCAAGCGTTGGGCGGAGGACCAGCACAGCCAGGGTGGCTTT
GTGGTACAGCCGCCGGCGCTCTGGCAAACCGAAAAGGATGACTGGACGGTCCCTTATGGC
CGCATCTACTTTGCCGGCGAGCACACCGCCTACCCGCACGGCTGGGTGGAGACGGCGGTC
AAGTCGGCGCTGCGCGCCGCCATCAAGATCAACAGCCGGAAGGGGCTGCATCGGACACG
GCCAGCCCCGAGGGGCACGCATCTGACATGGAGGGGCAGGGGCATGTGCATGGGGTGGCC
AGCAGCCCCCTCGCATGACCTGGCAAAGGAAGAAGGCAGCCACCCTCCAGTCCAAGGCCAG
TTATCTCTCCAAAACACGACCCACACGAGGACCTCGCATTAAAGTATTTTCGGAAAAAAA
AAA

184/392

FIGURE 176

MAPLALHLLVLVPILLSLVASQDWKAERSQDPFEKCMQDPDYEQLLKVVTWGLNRTLKPQ
RVIVVGAGVAGLVAAKVLSDAGHKVTILEADNRIGGRIFTYRDQNTGWIGELGAMRMPSS
HRILHKLCQGLGLNLTKFTQYDKNTWTEVHEVKLRNYVVEKVPEKLGALRPQEKGHSPE
DIYQMALNQALKDLKALGCRKAMKKFERHTLLEYLLGEGNLSRPAVQLLGDMSEDGFFY
LSFAEALRAHSCLSDRLQYSRIVGGWDLPRALLSSLSGLVLLNAPVVAMTQGPHDVHVQ
IETSPPARNLKVLKADVLLTASGPAVKRITFSPPLPRHMQEALRRLHYVPATKVFLSFR
RPFWREEHIEGGHSNTDRPSRMIFYPPPREGALLLASYTWSDAFAAGLSREEALRLAL
DDVAALHGPVVRQLWDGTGVVKRWAEDQHSQGGFVVQPPALWQTEKDDWTVPYGRIYFAG
EHTAYPHGWVETAVKSALRAAIKINSRKGPASDTASPEGHASDMEGQGHVHGVASSPSHD
LAKEEGSHPPVQGQLSLQNTTHTRTSH

Signal peptide:
amino acids 1-21

185/392

FIGURE 177

CCGGGGAGGGGAGGGCCCCGTCCCGCCCCCTCCCCGTCTCTCCCCGCCCCCTCCCCGTCCCTC
CCGCCGAAGCTCCGTCCCGCCCCGCGGGCCGGCTCCGCCCTCACCTCCCGGCCGCGGCTGC
CCTCTGCCCCGGGTTGTCCAAGATGGAGGGCGCTCCACCGGGGTGCTCGCCCTCCGGCTC
CTGCTGTTCTGTGGCGCTACCCGCCTCCGGCTGGCTGACGACGGGCGCCCCCGAGCCGCCG
CCGCTGTCCGGAGCCCCACAGGACGGCATCAGAATTAATGTAACTACACTGAAAGATGAT
GGGGACATATCTAAACAGCAGGTTGTTCTTAACATAACCTATGAGAGTGGACAGGTGTAT
GTAAATGACTTACCTGTAAATAGTGGTGTAAACCCGAATAAGCTGTCAGACTTTGATAGTG
AAGAATGAAAATCTTGAAAATTTGGAGGAAAAAGAATATTTTGGAATTGTCAGTGTAAAG
ATTTTAGTTTCATGAGTGGCCTATGACATCTGGTTCAGTTTGCAACTAATTGTCATTCAA
GAAGAGGTAGTAGAGATTGATGGAAAACAAGTTTCAAGCAAAAGGATGTCAGTCAAATTGAT
ATTTTAGTTAAGAACCGGGGAGTACTCAGACATTCAAACCTATACCCCTCCCTTTGGAAGAA
AGCATGCTCTACTCTATTTCTCGAGACAGTGACATTTTATTTACCCTTCCTAACCTCTCC
AAAAAAGAAAGTGTTAGTTCACTGCAAACCACTAGCCAGTATCTTATCAGGAATGTGGAA
ACCACTGTAGATGAAGATGTTTTACCTGGCAAGTTACCTGAAACTCCTCTCAGAGCAGAG
CCGCCATCTTCATATAAGGTAATGTGTGTCAGTGGATGGAAAAGTTTAGAAAAGATCTGTGT
AGGTTCTGGAGCAACGTTTTCCAGTATTCTTTCAGTTTTTGAACATCATGGTGGTTGGA
ATTACAGGAGCAGCTGTGGTAATAACCATCTTAAAGGTGTTTTCCAGTTTCTGAATAC
AAAGGAATTCCTCAGTTGGATAAAGTGGACGTCATACCTGTGACAGCTATCAACTTATAT
CCAGATGGTCCAGAGAAAAGAGCTGAAAACCTTGAAGATAAAACATGTATTTAAAACGCC
ATCTCATATCATGGACTCCGAAGTAGCCTGTTGCCTCAAATTTGCCACTTGAATATAAT
TTTCTTTAAATCGTT

186/392

FIGURE 178

MEGAPPGSLALRLLLFVALPASGWLTGAPPEPPPLSGAPQDGIRINVTTLKDDGDISKQQ
VVLNITYESGQVYVNDLPVNSGVTRISCQTLIVKNENLENLEEKEYFGIVSVRILVHEWP
MTSGSSLQLIVIQEEVVEIDGKQVQQKDVTEIDILVKNRGVLRHSNYTLPLEESMLYSIS
RDS DILFTLPNLSKKESVSSLQTTSQYLIRNVETTVDVLPGLPETPLRAEPPSSYKV
MCQWMEKFRKDL CRFWSNVFPVFFQFLNIMVVGITGA AVVITILKVFFPVSEYKGILQLD
KVDVIPVTAINLYPDGPEKRAENLEDKTCI

Signal peptide:

1-23

Transmembrane domain:

266-284

Leucine zipper pattern:

155-177

N-glycosylation site:

46-50, 64-68, 166-170, 191-195

Motif name: N-myristoylation site:

3-9, 42-48, 273-279

187/392

FIGURE 179

CTCCTTAGGTGGAAACCTGGGAGTAGAGTACTGACAGCAAAGACCGGGAAAGACCATAC
GTCCCCGGGCAGGGGTGACAAACAGGTGTCATCTTTTTTGATCTCGTGTGTGGCTGCCTTCC
TATTTCAAGGAAAGACGCCAAGGTAATTTTGACCCAGAGGAGCAATGATGTAGCCACCTC
CTAACCTTCCCTTCTTGAACCCCCAGTTATGCCAGGATTTACTAGAGAGTGTCAACTCAA
CCAGCAAGCGGCTCCTTCGGCTTAACCTGTGGTTGGAGGAGAGAACCTTTGTGGGGCTGC
GTTCTCTTAGCAGTGCTCAGAAGTGACTTGCCCTGAGGGTGGACCAGAAGAAAGGAAAGGT
CCCCCTCTTGCTGTTGGCTGCACATCAGGAAGGCTGTGATGGGAATGAAGGTGAAAACCTTG
GAGATTTCACTTCAGTCATTGCTTCTGCCTGCAAGATCATCCTTTAAAAGTAGAGAAGCT
GCTCTGTGTGGTGGTTAACTCCAAGAGGCAGAACTCGTTCTAGAAGGAAATGGATGCAAG
CAGCTCCGGGGGCCCCAAACGCATGCTTCTGTGGTCTAGCCCAGGGAAGCCCTTCCGTG
GGGGCCCCGGCTTTGAGGGATGCCACCGGTTCTGGACGCATGGCTGATTCTGAATGATG
ATGGTTTCGCCGGGGGCTGCTTGCCTGGATTTCGCCGGTGGTGGTTTTGCTGGTGTCTCTC
TGCTGTGCTATCTCTGTCTGTACATGTTGGCCTGCACCCCCAAAAGGTGACGAGGAGCAG
CTGGCACTGCCCAGGGCCAACAGCCCCACGGGGAAGGAGGGGTACCAGGCCGTCTTTCAG
GAGTGGGAGGAGCAGCACCGCAACTACGTGAGCAGCCTGAAGCGGCAGATCGCACAGCTC
AAGGAGGAGCTGCAGGAGAGGAGTGAGCAGCTCAGGAATGGGCAGTACCAAGCCAGCGAT
GCTGCTGGCCTGGGTCTGGACAGGAGCCCCCAGAGAAAACCCAGGCCGACCTCCTGGCC
TTCCTGCACTCGCAGGTGGACAAGGCAGAGGTGAATGCTGGCGTCAAGCTGGCCACAGAG
TATGCAGCAGTGCCTTTCGATAGCTTTACTCTACAGAAGGTGTACCAGCTGGAGACTGGC
CTTACCCGCCACCCCGAGGAGAAGCCTGTGAGGAAGGACAAGCGGGATGAGTTGGTGGAA
GCCATTGAATCAGCCTTGGAGACCCTGAACAATCCTGCAGAGAACAGCCCCAATCACCGT
CCTTACACGGCCTCTGATTTTCATAGAAGGGATCTACCGAACAGAAAGGGACAAAGGGACA
TTGTATGAGCTCACCTTCAAAGGGGACCACAAACACGAATTCAAACGGCTCATCTTATTT
CGACCATTTCAGCCCCATCATGAAAAGTGAAAAATGAAAAGCTCAACATGGCCAACACGCTT
ATCAATGTTATCGTGCCTCTAGCAAAAAGGGTGGACAAGTTCCGGCAGTTCATGCAGAAT
TTCAGGGAGATGTGCATTGAGCAGGATGGGAGAGTCCATCTCACTGTTGTTTACTTTGGG
AAAGAAGAAATAAATGAAGTCAAAGGAATACCTGAAAACACTTCCAAAGCTGCCAATTTC
AGGAACTTTACCTTCATCCAGCTGAATGGAGAATTTTCTCGGGGAAAGGGACTTGATGTT
GGAGCCCGCTTCTGGAAGGGAAGCAACGTCTTCTCTTTTTCTGTGATGTGGACATCTAC
TTCACATCTGAATTCCTCAATACGTGTAGGCTGAATACACAGCCAGGGAAGAAGGTATTT
TATCCAGTTCCTTTTCAGTCAGTACAATCCTGGCATAATATACGGCCACCATGATGCAGTC
CCTCCCTTGGAACAGCAGCTGGTCATAAAGAAGGAACTGGATTTTGGAGAGACTTTGGA
TTTGGGATGACGTGTGAGTATCGGTGAGACTTCATCAATATAGGTGGGTTTGATCTGGAC
ATCAAAGGCTGGGGCGGAGAGGATGTGCACCTTTATCGCAAGTATCTCCACAGCAACCTC
ATAGTGGTACGGACGCCTGTGCGAGGACTCTTCCACCTCTGGCATGAGAAGCGCTGCATG
GACGAGCTGACCCCCGAGCAGTACAAGATGTGCATGCAGTCCAAGGCCATGAACGAGGCA
TCCCACGGCCAGCTGGGCATGCTGGTGTTCAGGCACGAGATAGAGGCTCACCTTCGAAA
CAGAAACAGAAGACAAGTAGCAAAAAACATGAACTCCAGAGAAGGATTGTGGGAGACA
CTTTTTCTTCTTTTGAATTAAGTGGCTGCAACAGAGAAAAGACTTCCATAAA
GGACGACAAAAGAAATTGGACTGATGGGTGAGAGATGAGAAAGCCTCCGATTTCTCTCTGT
TGGGCTTTTTTACAACAGAAATCAAAATCTCCGCTTTGCCTGCAAAAGTAACCCAGTTGCA
CCCTGTGAAGTGTCTGACAAAGGCAGAAATGCTTGTGAGATTATAAGCCTAATGGTGTGGA
GGTTTTGATGGTGTTTACAATACACTGAGACCTGTTGTTTTGTGTGCTCATTGAAATATT
CATGATTTAAGAGCAGTTTTGTAAAAAATTCATTAGCATGAAAGGCAAGCATATTTCTCC
TCATATGAATGAGCCTATCAGCAGGGCTCTAGTTTCTAGGAATGCTAAAAATATCAGAAGG
CAGGAGAGGAGATAGGCTTATTATGATACTAGTGAGTACATTAAGTAAAAATAAATGGAC
CAGAAAAGAAAAGAAACCATAAATATCGTGTATATTTTCCCCAAGATTAACCAAAAATA

188/392

ATCTGCTTATCTTTTTGGTTGTCCTTTTAACTGTCTCCGTTTTTTTTCTTTTATTTAAAAA
TGCACTTTTTTTCCCTTGTGAGTTATAGTCTGCTTATTTAATTACCACTTTGCAAGCCTT
ACAAGAGAGCACAAAGTTGGCCTACATTTTTATATTTTTTAAGAAGATACTTTGAGATGCA
TTATGAGAACTTTCAGTTCAAAGCATCAAATTGATGCCATATCCAAGGACATGCCAAATG
CTGATTCTGTCAGGCACTGAATGTCAGGCATTGAGACATAGGGAAGGAATGGTTTGTACT
AATACAGACGTACAGATACTTTCTCTGAAGAGTATTTTCGAAGAGGAGCAACTGAACACT
GGAGGAAAAGAAAATGACACTTTCTGCTTTACAGAAAAGGAACTCATTGAGACTGGTGA
TATCGTGATGTACCTAAAAGTCAGAAACCACATTTTCTCCTCAGAAGTAGGGACCGCTTT
CTTACCTGTTTAAATAAACCAAAGTATACCGTGTGAACCAAACAATCTCTTTTCAAAACA
GGGTGCTCCTCCTGGCTTCTGGCTTCCATAAGAAGAAATGGAGAAAAATATATATATATA
TATATATATTGTGAAAGATCAATCCATCTGCCAGAATCTAGTGGGATGGAAGTTTTTGCT
ACATGTTATCCACCCAGGCCAGGTGGAAGTAACTGAATTATTTTTTAAATTAAGCAGTT
CTACTCAATCACCAAGATGCTTCTGAAAATTGCATTTTATTACCATTTCAAACATTTTTT
TAAAAATAAATACAGTTAACATAGAGTGGTTTCTTCATTGATGTGAAAATTATTAGCCAG
CACCAGATGCATGAGCTAATTATCTCTTTGAGTCCTTGCTTCTGTTTGCTCACAGTAAAC
TCATTGTTTAAAAGCTTCAAGAACATTCAAGCTGTTGGTGTGTTAAAAAATGCATTGTAT
TGATTTGTACTGGTAGTTTATGAAATTTAATTAAACACAGGCCATGAATGGAAGGTGGT
ATTGCACAGCTAATAAAATATGATTTGTGGATATGAA

189/392

FIGURE 180

MMVRRGLLAWISRVVLLVLLCCAISVLYMLACTPKGDEEQLALPRANSPTGKEGYQAV
LQEWEEQHRNYVSSLKRQIAQLKEELQERSEQLRNGQYQASDAAGLGLDRSPPEKTQADL
LAFLHSQVDKAEVNAGVKLATEYA AVPFDSFTLQKVYQLETGLTRHPPEKPVRKDKRDEL
VEAIESALETLNPAENSPNHRPYTASDFIEGIYRTERDKGTLYELTFKGDHKHEFKRLI
LFRPFSPIMKVKNEKLNMAN TLINVIVPLAKRVDKFRQFMQNFREMCIEQDGRVHLTVVY
FGKEEINEVKGILENTSKAANFRNFTFIQLNGEFSRGKGLDVGARFWKGSNVLLFFCDVD
IYFTSEFLNTCRLNTQPGKKVFYPVLF SQYNPGIIYGHDAVPPLEQQQLVIKKETGFWRD
FGFGMTCQYRSDFINIGGFDLDIKGWGGEDVHLYRKYLHSNLIVVRTPV RGLFHLWHEKR
CMDELTPEQYKMC MQSKAMNEASHGQLGMLVFRHEIEAHLRKQKQKTSSKKT

190/392

FIGURE 181

CGTCTCTGCGTTTCGCCATGCGTCCCGGGGCGCCAGGGCCACTCTGGCCTCTGCCCTGGGG
GGCCCTGGCTTGGGCCGTGGGCTTCGTGAGCTCCATGGGCTCGGGGAACCCCGCGCCCGG
TGGTGT'TTGCTGGCTCCAGCAGGGCCAGGAGGCCACCTGCAGCCTGGTGCTCCAGACTGA
TGTCACCCGGGCCGAGTGCTGTGCCTCCGGCAACATTGACACCGCCTGGTCCAACCTCAC
CCACCCGGGGAAACAAGATCAACCTCCTCGGCTTCTTGGGCCTTGTCCTACTGCCTTCCCTG
CAAAGATTTCGTGCGACGGCGTGAGTGCGGCCCGGGCAAGGCGTGCCGCATGCTGGGGGG
CCGCCCGCGCTGCGAGTGCGCGCCCGACTGCTCGGGGCTCCCGGCGCGGCTGCAGGTCTG
CGGCTCAGACGGCGCCACCTACCGCGACGAGTGCGAGCTGCGCGCCGCGCGCTGCCGCGG
CCACCCGGACCTGAGCGTCATGTACCGGGGCCGCTGCCGCAAGTCCTGTGAGCACGTGGT
GTGCCCGCGGCCACAGTCGTGCGTCTGGACCAGACGGGCAGCGCCCACTGCGTGGTGTG
TCGAGCGGCGCCCTGCCCTGTGCCCTCCAGCCCCGGCCAGGAGCTTTGCGGCAACAACAA
CGTCACCTACATCTCCTCGTGCCACATGCGCCAGGCCACCTGCTTCTGGGCCGCTCCAT
CGGCGTGCGCCACGCGGGCAGCTGCGCAGGCACCCCTGAGGAGCCGCCAGGTGGTGAGTC
TGCAGAAGAGGAAGAGAACTTCGTGTGAGCCTGCAGGACAGGCCTGGGCCTGGTGCCCGA
GGCCCCCATCATCCCCCTGTTATTTATTGCCACAGCAGAGTCTAATTTATATGCCACGGA
CACTCCTTAGAGCCCGGATTTCGACCACTTGGGGATCCCAGAACCTCCCTGACGATATCC
TGGAAGGACTGAGGAAGGGAGGCCTGGGGGCCGGCTGGTGGGTGGGATAGACCTGCGTTC
CGGACACTGAGCGCCTGATTTAGGGGCCCTTCTCTAGGATGCCCCAGCCCCCTACCCTAAGA
CCTATTGCCGGGGAGGATTCCACACTTCCGCTCCTTTGGGGATAAACCTATTAATTATTG
CTACTATCAAGAGGGGCTGGGCATTCTCTGCTGGTAATTCTGAAGAGGCATGACTGCTTT
TCTCAGCCCCAAGCCTCTAGTCTGGGTGTGTACGGAGGGTCTAGCCTGGGTGTGTACGGA
GGGTCTAGCCTGGGTGAGTACGGAGGGTCTAGCCTGGGTGAGTACGGAGGGTCTAGCCTG
GGTGAGTACGGAGGGTCTAGCCTGGGTGTGTATGGAGGATCTAGCCTGGGTGAGTATGGA
GGGTCTAGCCTGGGTGAGTATGGAGGGTCTAGCCTGGGTGTGTATGGAGGGTCTAGCCTG
GGTGAGTATGGAGGGTCTAGCCTGGGTGTGTATGGAGGGTCTAGCCTGGGTGAGTATGGA
GGGTCTAGCCTGGGTGTGTACGGAGGGTCTAGTCTGAGTGCGTGTGGGGACCTCAGAACA
CTGTGACCTTAGCCCAGCAAGCCAGGCCCTTCATGAAGGCCAAGAAGGCTGCCACCATTCT
CCTGCCAGCCCAAGAACTCCAGCTTCCCCACTGCCTCTGTGTGCCCCCTTTGCGTCCTGTG
AAGGCCATTGAGAAATGCCCAGTGTGCCCCCTGGGAAAGGGCACGGCCTGTGCTCCTGAC
ACGGGCTGTGCTTGGCCACAGAACCACCCAGCGTCTCCCCTGCTGCTGTCCACGTCAGTT
CATGAGGCAACGTCGCGTGGTCTCAGACGTGGAGCAGCCAGCGGCAGCTCAGAGCAGGGC
ACTGTGTCCGGCGGAGCCAAGTCCACTCTGGGGGAGCTCTGGCGGGGACCACGGGCCACT
GCTCACCCACTGGCCCCGAGGGGGGTGTAGACGCCAAGACTCACGCATGTGTGACATCCG
GAGTCCTGGAGCCGGGTGTCCCAGTGGCACCCTAGGTGCCTGCTGCCTCCACAGTGGGG
TTCACACCCAGGGCTCCTTGGTCCCCACAACCTGCCCCGGCCAGGCCTGCAGACCCAGA
CTCCAGCCAGACCTGCCTCACCCACCAATGCAGCCGGGGCTGGCGACACCAGCCAGGTGC
TGGTCTTGGGCCAGTTCTCCACGACGGCTCACCCCTCCCCTCCATCTGCGTTGATGCTCA
GAATCGCCTACCTGTGCCTGCGTGTAACACAGCCTCAGACCAGCTATGGGGAGAGGAC
AACACGGAGGATATCCAGCTTCCCCGGTCTGGGGTGAGGAATGTGGGGAGCTTGGGCATC
CTCCTCCAGCCTCCTCCAGCCCCCAGGCAGTGCCCTTACCTGTGGTGCCAGAAAAGTGCC
CCTAGGTTGGTGGGTCTACAGGAGCCTCAGCCAGGCAGCCACCCACCCCTGGGGCCCTG
CCTCACCAAGGAAATAAAGACTCAAGCCATAAAAAAA

191/392

FIGURE 182

MRPGAPGPLWPLPWGALAWAVGFVSSMSGGNPAPGGVCWLQQGQEATCSLVLQTDVTRAE
CCASGNIDTAWSNLTHPGNKINLLGFLGLVHCLPCKDSCDGVECGPGKACRMLGGRPRCE
CAPDCSGLPARLQVCGSDGATYRDECELRAARCRGHPDLSVMYRGRCRKSCHEVVCPRPQ
SCVVDQTGSAHCVVCRAAPCPVPSSPGQELCGNNNVTYISSCHMRQATCFLGRSIGVRHA
GSCAGTPEEPPGGESAEEEENFV

Important features:**Signal peptide:**

amino acids 1-20

N-glycosylation sites:

amino acids 73-77, 215-219

Osteonectin domain proteins:

amino acids 97-130, 169-202

192/392

FIGURE 183

CACTCATTTCATTCCAAAGGGTCTCTCAAGGCAATGGTAATGTGCAAGGAGGTGATACCTA
AATGAATGACCAAAAGAACATGCTTCTGCTTTTGTGTGTCTCCTACATTTTAGACATTTG
TTTGTTTTCTCTTGGTAGCCTTTAAATTCCTTGAAGCCCAGGACCATGTCTCACTTACCTT
TGTGTTTTCCACTAACTAGTCTACCTCCTGGAATTGGCAGATACTCAGTGAAAGCCTGTGA
AATAAGTGATGTCTATTTCTAGCATATTATTCTGAGATTTAATGATAGATTTAGTGATTG
AATGAGATTTCCATTTTCAAATACAGCAAAAGCATAACTATTTTCATTTCATTTCATATTCA
TTCAACTTCATTCTCAAATTAGGTCTGAGTTAACTAATAATTACCTTTGAAATGTGTG
GGTTATTTGAGGCAATCAGGTGGTGACATTGAGCTCTCAGCCAGAGTTTGTCTGGAAT
TGATTCAGTTCATTTGCATTGATTTTGTCTCAGAAGCCAAGGTTTCCCATGAAAAATC
ATTCCCCTTGAATTGGGCTGTGATTTCTGCTGCGTTTAAAGTAAAGGAAGCCTCTTGGTT
CTAGTTCTGCAAACTTACACACTGAACTGGGACAAGTTTTTGTCTTAGAGTAATGGCTGGG
AAAAGAGGAACCTTTCATTTTATTTCAGAAGTCAAAAACAAAGGCCTCCAGCCACCTGGA
GATGTTTTGTTCAGACACCAGCCTGGCTCTGTCTTTATGCCTAACAATTGAGCATCCAG
TCTTCTTTGTGCTGGGACCATTGCTCAGCTCTGCAAGGGGAAAAGAGGGAGAAAGCCAGA
GCTGCCAGGCTTCTTGCACTGGGGCCGGGGAGGGTTTCTGGGAAGCAGGTGCTCTCTGG
CTTCTTGGTACGTGAGGCTCTCGGAGCTGCCTCTCCTCTGACCCTCAGGTCCTCACCGAG
TTTGCTCCAGGAGTATATTGAAAACATACCCAGTGCTCTCTCAAGCACCCACTGCTTAGA
GGGCCCAGATTTCTTTTCTTCTTCCCTTGCAAGCTGGAGACTGCATCGGGCATCTGG
TGTTTAAACTAAACAGGAAAAGTGAATAAGGTCCACAGTGCTCATTGTGTAGACTAGCT
GCCCTCCGATGGGTGCTCTGATTATCAGTGGTTCCAGTGCAAGGCCTGTCACTAAACAGG
CCTCACTTCCTCCTTGGGGGCTTTCCCATGGGAGGTGTGGCTTTTTACTCTACATGGAAA
TGACTCTCTGCAGCCACAGAACACAGTCATTTTCTGAATTATCCAGTCTCTCATGCGCC
CTGGATTCTCCTCAGATGCCTTATATCTCTTGTGCAAAGTTGTCTAAATTTGGTTCCCAG
CTTCCAAGCCTTGCCTTTTGGCCTTCTGGAAGTATTTTTGTTGATGAGTCGTCTGTCTAT
TATTCTCTAAATGATTTGCTTTTTGTTTCTTTCATTTCCTATTTCCACCCACATATACA
CACATGCTTCTTAACTTAGGGGATTACATGCCAATAAATCTATTGTTGAAAATGCACTAA
TACTATCGCAAAGACGAAAATTCACAGGCTGAACCGTTGTAAGTCCATATGCTCCTCAAC
TTACATGTGTGATGGAGTTATGCCCAAATAAGTCCATCGTCAAGTTGAAAAATCAAAATC
AAGCCATCTTAGGTTGAGGACCATTGTGTTGTACCTCCAAGATGTCATATCTTTAAACA
TACTCCCTAGCTTTTCTTTTACTTTTTATTTTGAAGTAATTATAGAATCACAGAAAGTT
GCAAAAAA

193/392

FIGURE 184

MGALIIISGSSAGPVTQASLPPWGLSHGRCGFLLYMENTLCSHRTQSFSELSQSLMRPGF
LQMPYISCAKLSKIWFPAKPCLLAFLEVFLLMSRLSLFSKMICFLFLSFLFPPHIYTHAS

Important features of the protein:

Signal peptide:

amino acids 1-41

Transmembrane domain:

amino acids 88-107

Casein kinase II phosphorylation site:

amino acids 47-50

N-myristoylation site:

amino acids 24-29

194/392

FIGURE 185

AACTCAAACCTCTCTCTCTGGGAAAACGCGGTGCTTGCTCCTCCCGGAGTGGCCTTGGCA
GGGTGTTGGAGCCCTCGGTCTGCCCCGTCCGGTCTCTGGGGCCAAGGCTGGGTTTCCCTC
ATGTATGGCAAGAGCTCTACTCGTGCGGTGCTTCTTCTCCTTGGCATACAGCTCACAGCT
CTTTGGCCTATAGCAGCTGTGGAAATTTATACCTCCCGGGTGCTGGAGGCTGTTAATGGG
ACAGATGCTCGGTTAAAATGCACTTTCTCCAGCTTTGCCCCCTGTGGGTGATGCTCTAACA
GTGACCTGGAATTTTCTGTCCTCTAGACGGGGGACCTGAGCAGTTTGTATTCTACTACCAC
ATAGATCCCTTCCAACCCATGAGTGGGCGGTTTAAGGACCGGGTGTCTTGGGATGGGAAT
CCTGAGCGGTACGATGCCTCCATCCTTCTCTGGAACTGCAGTTCGACGACAATGGGACA
TACACCTGCCAGGTGAAGAACCCACCTGATGTTGATGGGGTGATAGGGGAGATCCGGCTC
AGCGTCGTGCACACTGTACGCTTCTCTGAGATCCACTTCCTGGCTCTGGCCATTGGCTCT
GCCTGTGCACTGATGATCATAATAGTAATTGTAGTGGTCCTCTTCCAGCATTACCGGAAA
AAGCGATGGGCCGAAAGAGCTCATAAAGTGGTGGAGATAAAATCAAAAAGAAGAGGAAAGG
CTCAACCAAGAGAAAAAGGTCTCTGTTTATTTAGAAGACACAGACTAACAATTTTAGATG
GAAGCTGAGATGATTTCCAAGAACAAGAACCTAGTATTTCTTGAAGTTAATGGAACTT
TTCTTTGGCTTTTCCAGTTGTGACCCGTTTTCCAACCAGTTCTGCAGCATATTAGATTCT
AGACAAGCAACACCCCTCTGGAGCCAGCACAGTGCTCCTCCATATCACCAGTCATACACA
GCCTCATTATTAAGGTCTTATTTAATTTTCAAGAGTGTAATTTTTTCAAGTGCTCATTAGG
TTTTATAACAAGAAGCTACATTTTTTGCCCTTAAGACACTACTTACAGTGTTATGACTTG
TATACACATATATTGGTATCAAAGGGGATAAAAGCCAATTTGTCTGTTACATTTCTTTTC
ACGTATTTCTTTTAGCAGCACTTCTGCTACTAAAGTTAATGTGTTTACTCTCTTTCTTC
CCACATTCTCAATTAAAAGGTGAGCTAAGCCTCCTCGGTGTTTCTGATTAAACAGTAAATC
CTAAATTCAAACGTGTTAAATGACATTTTTATTTTTATGTCTCTCCTTAACATATGAGACAC
ATCTTGTTTTACTGAATTTCTTTCAATATTCCAGGTGATAGATTTTTTGTCG

195/392

FIGURE 186

MYGKSSTRAVLLLLGIQLTALWPAAVEIYTSRVLEAVNGTDARLKCTFSSFAPVGDALT
VTWNFRPLDGGPEQFVFYHIDPFQPMSEGRFKDRVSWDGNPERYDASILLWKLQFDDNGT
YTCQVKNPPDVGVIIEIRLSVVHTVRFSEIHFLALALIGSACALMIIIVIVVVLFFQHYRK
KRWAEARAHKVVEIKSKEEERLNQEKVSVYLETD

196/392

FIGURE 187

GCATTTTTGTCTGTGCTCCCTGATCTTCAGGTCACCACCATGAAGTTCTTAGCAGTCCTG
GTACTCTTGGGAGTTTCCATCTTTCTGGTCTCTGCCAGAATCCGACAACAGCTGCTCCA
GCTGACACGTATCCAGCTACTGGTCCTGCTGATGATGAAGCCCCTGATGCTGAAACCACT
GCTGCTGCAACCACTGCGACCACTGCTGCTCCTACCACTGCAACCACCGCTGCTTCTACC
ACTGCTCGTAAAGACATTCCAGTTTTACCCAAATGGGTGGGGATCTCCCGAATGGTAGA
GTGTGTCCCTGAGATGGAATCAGCTTGAGTCTTCTGCAATTGGTCACAACCTATTCATGCT
TCCTGTGATTTTCATCCAACCTACTTACCTTGCTACGATATCCCCTTTATCTCTAATCAGT
TTATTTTCTTTCAAATAAAAAATAACTATGAGCAACATAAAAAAAAAAAAAA

197/392

FIGURE 188

MKFLAVLVLLGVSI FLVSAQNPTTAAPADTYPATGPADDEAPDAETTAAATTATTAAPT
ATTAASTTARKDIPVLPKWVGDL PNGRVCP

198/392

FIGURE 189

GAGCGAACATGGCAGCGCGTTGGCGGTTTTGGTGTGTCTCTGTGACCATGGTGGTGGCGC
TGCTCATCGTTTTGCGACGTTCCCTCAGCCTCTGCCCAAAGAAAGAAGGAGATGGTGTAT
CTGAAAAGGTTAGTCAGCTGATGGAATGGACTAACAAAAGACCTGTAATAAGAATGAATG
GAGACAAGTTCGGTCGCCTTGTGAAAGCCCCACCGAGAAATTACTCCGTTATCGTCATGT
TCACTGCTCTCCAACTGCATAGACAGTGTGTCTGTTTGAAGCAAGCTGATGAAGAATTCC
AGATCCTGGCAAACCTCTGGCGATACTCCAGTGCATTACCAACAGGATATTTTTTGGCCA
TGGTGGATTTTGATGAAGGCTCTGATGTATTTTCAAGTGTAAACATGAATTCAGCTCCAA
CTTTCATCAACTTTCTGCAAAAGGGAAACCCAAACGGGGTGATACATATGAGTTACAGG
TGCGGGGTTTTTCAGCTGAGCAGATTGCCCGGTGGATCGCCGACAGAACTGATGTCAATA
TTAGAGTGATTAGACCCCCAAATTATGCTGGTCCCTTATGTTGGGATTGCTTTTGGCTG
TTATTGGTGGACTTGTGTATCTTCGAAGAAGTAATATGGAATTTCTCTTAATAAACTG
GATGGGCTTTTGCAGCTTTGTGTTTTGTGCTTGCTATGACATCTGGTCAAATGTGGAACC
ATATAAGAGGACCACCATATGCCATAAGAATCCCCACACGGGACATGTGAATTATATCC
ATGGAAGCAGTCAAGCCCAGTTTGTAGCTGAAACACACATTGTTCTTCTGTTTAATGGTG
GAGTTACCTTAGGAATGGTGCCTTTTATGTGAAGCTGCTACCTCTGACATGGATATTGGAA
AGCGAAAGATAATGTGTGTGGCTGGTATTGGACTTGTTGTATTATTCTTCAGTTGGATGC
TCTCTATTTTTAGATCTAAATATCATGGCTACCCATACAGCTTTCTGATGAGTTAAAAAG
GTCCCAGAGATATATAGACACTGGAGTACTGGAAATTGAAAAACGAAAATCGTGTGTGTT
TGAAAAGAAGAATGCAACTTGTATATTTTTGTATTACCTCTTTTTTTTCAAGTGATTTAAAT
AGTTAATCATTTAAACCAAAGAAGATGTGTAGTGCCTTAACAAGCAATCCTCTGTCAAAAT
CTGAGGTATTTGAAAATAATTATCCTCTTAACCTTCTCTTCCCAGTGAACTTTATGGAAC
ATTTAATTTAGTACAATTAAGTATATTATAAAAAATTGTAAACTACTACTTTGTTTTAGT
TAGAACAAAGCTCAAACTACTTTAGTTAACTTGGTCATCTGATTTTATATTGCCCTTATC
CAAAGATGGGGAAAGTAAGTCCTGACCAGGTGTTCCACATATGCCTGTTACAGATAACT
ACATTAGGAATTCATTCTTAGCTTCTTCATCTTTGTGTGGATGTGTATACTTTACGCATC
TTTCCTTTTGTAGTAGAGAAATTATGTGTGTGTCATGTGGTCTTCTGAAAATGGAACACCATT
CTTCAGAGCACACGTCTAGCCCTCAGCAAGACAGTTGTTTCTCCTCCTCCTTGCATATTT
CCTACTGCGCTCCAGCCTGAGTGATAGAGTGAGACTCTGTCTCAAAAAAAGTATCTCTA
AATACAGGATTATAATTTCTGCTTGAGTATGGTGTAACTACCTTGTATTTAGAAAGATT
TCAGATTCATTCCATCTCCTTAGTTTTCTTTTAAGGTGACCCATCTGTGATAAAAATATA
GCTTAGTGCTAAAATCAGTGTAACCTTATACATGGCCTAAAATGTTTCTACAAATTAGAGT
TTGTCACTTATTCATTTGTACCTAAGAGAAAAATAGGCTCAGTTAGAAAAGGACTCCCT
GGCCAGGCGCAGTGACTTACGCCTGTAATCTCAGCACTTTGGGAGGCCAAGGCAGGCAGA
TCACGAGGTCAGGAGTTTCGAGACCATCCTGGCCAAACATGGTGAAACCCCGTCTCTACTAA
AAATATAAAAAATTAGCTGGGTGTGGTGGCAGGAGCCTGTAATCCCAGCTACACAGGAGGC
TGAGGCACGAGAATCACTTGAACCTCAGGAGATGGAGGTTTCAGTGAGCCGAGATCACGCC
ACTGCACTCCAGCCTGGCAACAGAGCGAGACTCCATCTCAAAAAAAAAAAAAA

199/392

FIGURE 190

MAARWRFWCVSVTMVVALLIVCDVPSASAQRKKEMVLSEKVSQLMWETNKRPVIRMNGDK
FRRLVKAPPRNYSVIVMFTALQLHRQCVVCKQADEEFQILANSWRYSSAFTNRIFFAMVD
FDEGSDVFQMLNMNSAPTFINFPAGKPKRGDTYELQVRGFSAEQIARWIADRTDVNIRV
IRPPNYAGPLMLGLLLAVIGGLVYLRRSNMEFLFNKTGWAFALCFVLAMTSGQMWNHIR
GPPYAHKNPHTGHVNYIHGSSQAQFVAETHIVLLFNNGGVTLGMVLLCEAATSDMDIGKRK
IMCVAGIGLVVLFFSWMLSIFRSKYHGYPSFLMS

Signal peptide:

amino acids 1-29

Transmembrane domains:

amino acids 183-205, 217-237, 217-287, 301-321

200/392

FIGURE 191

GAGAGAAAGTCAGCCTGGCAGAGAGACTCTGAAATGAGGGATTAGAGGTGTTCAAGGAGCA
AGAGCTTCAGCCTGAAGACAAGGGAGCAGTCCCTGAAGACGCTTCTACTGAGAGGTCTGC
CATGGCCTCTCTTGGCCTCCAACTTGTGGGCTACATCCTAGGCCTTCTGGGGCTTTTGGG
CACACTGGTTGCCATGCTGCTCCCCAGCTGGAACAAGTTCTTATGTCGGTGCCAGCAT
TGTGACAGCAGTTGGCTTCTCCAAGGGCCTCTGGATGGAATGTGCCACACACAGCACAGG
CATCACCCAGTGTGACATCTATAGCACCTTCTGGGCCTGCCGCTGACATCCAGGCTGC
CCAGGCCATGATGGTGACATCCAGTGCAATCTCCTCCCTGGCCTGCATTATCTCTGTGGT
GGGCATGAGATGCACAGTCTTCTGCCAGGAATCCCGAGCCAAAGACAGAGTGGCGGTAGC
AGGTGGAGTCTTTTTTCATCCTTGGAGGCCTCCTGGGATTCATTCCTGTTGCCTGGAATCT
TCATGGGATCCTACGGGACTTCTACTCACCCTGGTGCCTGACAGCATGAAATTTGAGAT
TGGAGAGGCTCTTTACTTGGGCATTATTTCTTCCCTGTTCTCCCTGATAGCTGGAATCAT
CCTCTGCTTTTCTGCTCATCCCAGAGAAATCGCTCCAATACTACGATGCCTACCAAGC
CCAACCTCTTGCCACAAGGAGCTCTCCAAGGCCTGGTCAACCTCCCAAAGTCAAGAGTGA
GTTCAATTCTACAGCCTGACAGGGTATGTGTGAAGAACCAGGGGCCAGAGCTGGGGGGT
GGCTGGGTCTGTGAAAAACAGTGGACAGCACCCCGAGGGCCACAGGTGAGGGACACTACC
ACTGGATCGTGTGAGAAGGTGCTGCTGAGGATAGACTGACTTTGGCCATTGGATTGAGCA
AAGGCAGAAATGGGGGCTAGTGTAACAGCATGCAGGTTGAATTGCCAAGGATGCTCGCCA
TGCCAGCCTTTCTGTTTTCTCACCTTGCTGCTCCCCTGCCCTAAGTCCCCAACCTCAA
CTTGAAACCCCATTCCTTAAGCCAGGACTCAGAGGATCCCTTTGCCCTCTGGTTTACCT
GGGACTCCATCCCCAAACCCACTAATCACATCCCCTGACTGACCCTCTGTGATCAAAGA
CCCTCTCTCTGGCTGAGGTTGGCTCTTAGCTCATTGCTGGGGATGGGAAGGAGAAGCAGT
GGCTTTTGTGGGCATTGCTCTAACCTACTTCTCAAGCTTCCCTCCAAAGAACTGATTGG
CCCTGGAACCTCCATCCCCTCTTGTTATGACTCCACAGTGTCCAGACTAATTTGTGCAT
GAACTGAAATAAAACCATCCTACGGTATCCAGGGAACAGAAAGCAGGATGCAGGATGGGA
GGACAGGAAGGCAGCCTGGGACATTTAAAAAATA

201/392

FIGURE 192

MASLGLQLVGYYLGLLGLLGLTLVAMLLPSWKTSSYVGASIVTAVGFSKGLWMECATHSTG
ITQCDIYSTLLGLPADIQAAQAMMVTSSAIISSLACIISVVGMRCTVFCQESRAKDRVAVA
GGVFFILGGLLGFIPVAWNLHGILRDFYSPLVPDSMKFEIGEALYLGIISSLFSLIAGII
LCFSCSSQRNRSNYYDAYQAQPLATRSSPRPGQPPKVKSEFNSSYSLTGYV

Important features of the protein:

Signal peptide:

amino acids 1-24

Transmembrane domains:

amino acids 82-102, 117-140, 163-182

N-glycosylation site:

amino acids 190-193

PMP-22 / EMP / MP20 family proteins:

amino acids 46-59

202/392

FIGURE 193

CTCCACTGCAACCACCCAGAGCCATGGCTCCCCGAGGCTGCATCGTAGCTGTCTTTGCCA
TTTTCTGCATCTCCAGGCTCCTCTGCTCACACGGAGCCCCAGTGGCCCCCATGACTCCTT
ACCTGATGCTGTGCCAGCCACACAAGAGATGTGGGGACAAGTTCTACGACCCCCTGCAGC
ACTGTTGCTATGATGATGCCGTCGTGCCCTTGGCCAGGACCCAGACGTGTGGAAACTGCA
CCTTCAGAGTCTGCTTTGAGCAGTGCTGCCCCTGGACCTTCATGGTGAAGCTGATAAACC
AGAACTGCGACTCAGCCCGGACCTCGGATGACAGGCTTTGTGCGCAGTGTCAGCTTAATGGA
ACATCAGGGGAACGATGACTCCTGGATTCTCCTTCCTGGGTGGGCCTGGAGAAAGAGGCT
GGTGTACCTGAGATCTGGGATGCTGAGTGGCTGTTTGGGGGCCAGAGAAACACACACTC
AACTGCCCACTTCATTCTGTGACCTGTCTGAGGCCACCCTGCAGCTGCCCTGAGGAGGC
CCACAGGTCCCCTTCTAGAATTCTGGACAGCATGAGATGCGTGTGCTGATGGGGGCCAG
GGACTCTGAACCCCTCCTGATGACCCCTATGGCCAACATCAACCCGGCACCCCCCAAGGC
TGGCTGGGGAACCCCTCACCCCTTCTGTGAGATTTTCCATCATCTCAAGTTCTTTCTATC
CAGGAGCAAAGCACAGGATCATAATAAATTTATGTACTTTATAAATGAAAA

203/392

FIGURE 194

MAPRGCIVAVFAIFCISRLLC SHGAPVAPMTPYLM LCQPHKRCGDKFYDPLQHCCYDDAV
VPLARTQTCGNCTFRVCFEQCCPWT FMVKLINQNCD SARTSDDRLCRSVS

Signal peptide:
amino acids 1-24

204/392

FIGURE 195

CATTTCCAACAAGAGCACTGGCCAAGTCAGCTTCTTCTGAGAGAGTCTCTAGAAGACATG
ATGCTACACTCAGCTTTGGGTCTCTGCCTCTTACTCGTCACAGTTTCTTCCAACCTTGCC
ATTGCAATAAAAAAGGAAAAGAGGCCTCCTCAGACACTCTCAAGAGGATGGGGAGATGAC
ATCACTTGGGTACAACTTATGAAGAAGGTCTCTTTTATGCTCAAAAAAGTAAGAAGCCA
TTAATGGTTATTCATCACCTGGAGGATTGTCAATACTCTCAAGCACTAAAGAAAGTATTT
GCCCAAATGAAGAAATACAAGAAATGGCTCAGAATAAGTTCATCATGCTAAACCTTATG
CATGAAACCACTGATAAGAATTTATCACCTGATGGGCAATATGTGCCTAGAATCATGTTT
GTAGACCCTTCTTTAACAGTTAGAGCTGACATAGCTGGAAGATACTCTAACAGATTGTAC
ACATATGAGCCTCGGGATTTACCCCTATTGATAGAAAACATGAAGAAAGCATTAAAGACTT
ATTCAGTCAGAGCTATAAGAGATGATGGAAAAAAGCCTTCACTTCAAAGAAGTCAAATTT
CATGAAGAAAACCTCTGGCACATTGACAAATACTAAATGTGCAAGTATATAGATTTTGTGTA
ATATTACTATTTAGTTTTTTTTAATGTGTTTGCAATAGTCTTATTAAAATAAATGTTTTTT
AAATCTGA

205/392

FIGURE 196

MMLHSALGLCLLLVTVSSNLAIKKEKRPPQTLSRGWGDDITWVQTYEEGLFYAQKSKK
PLMVIHHLEDCQYSQALKKVFAQNEEIQEMAQNKFIMLNLMHETTDKNLSPDGQYVPRIM
FVDPSLTVRADIAGRYSNRLYTYEPRDLPLLIENMKKALRLIQSEL

Important features:

Signal peptide:

amino acids 1-23

N-myristoylation site:

amino acids 51-57

206/392

FIGURE 197

GGGGGCGGGTGCCTGGAGCACGGCGCTGGGGCCGCCCCGAGCGCTCACTCGCTCGCACTC
AGTCGCGGGAGGCTTCCCCGCGCCGCGCGTCCCCCGCTCCCCGGCACCAGAAGTTC
CTCTGCGCGTCCGACGGCGACATGGGCGTCCCCACGGCCCTGGAGGCCGGCAGCTGGCGC
TGGGGATCCCTGCTCTTCGCTCTCTTCCTGGCTGCGTCCCTAGGTCCGGTGGCAGCCTTC
AAGGTCGCCACGCCGTATTCCCTGTATGTCTGTCCCGAGGGGCAGAACGTCACCTCACC
TGCAGGCTCTTGGGCCCTGTGGACAAAGGGCACGATGTGACCTTCTACAAGACGTGGTAC
CGCAGCTCGAGGGGCGAGGTGCAGACCTGCTCAGAGCGCCGGCCCATCCGCAACCTCACG
TTCCAGGACCTTCACCTGCACCATGGAGGCCACCAGGCTGCCAACACCAGCCACGACCTG
GCTCAGCGCCACGGGCTGGAGTCGGCCTCCGACCACCATGGCAACTTCTCCATCACCATG
CGCAACCTGACCCTGCTGGATAGCGCCTCTACTGCTGCCTGGTGGTGGAGATCAGGCAC
CACCCTCGGAGCACAGGGTCCATGGTGCCATGGAGCTGCAGGTGCAGACAGGCAAAGAT
GCACCATCCAAGTGTGTGGTGTACCCATCCTCCTCCAGGATAGTGAAAACATCACGGCT
GCAGCCCTGGCTACGGGTGCCTGCATCGTAGGAATCCTCTGCCTCCCCCTCATCCTGCTC
CTGGTCTACAAGCAAAGGCAGGCAGCCTCCAACCGCCGTGCCAGGAGCTGGTGC GGATG
GACAGCAACATTCAAGGGATTGAAAACCCCGGCTTTGAAGCCTCACCACCTGCCAGGGG
ATACCCGAGGCCAAAGTCAGGCACCCCTGTCTTATGTGGCCAGCGGCAGCCTTCTGAG
TCTGGGCGGCATCTGCTTTCGGAGCCCAGCACCCCTGTCTCCTCCAGGCCCGGAGAC
GTCTTCTTCCCATCCCTGGACCTGTCCCTGACTCTCCAAACTTTGAGGTCATCTAGCCC
AGCTGGGGGACAGTGGGCTGTTGTGGCTGGGTCTGGGGCAGGTGCATTTGAGCCAGGGCT
GGCTCTGTGAGTGGCCTCCTTGGCCTCGGCCCTGGTTCCTCCTCCTGCTCTGGGCTCA
GATACTGTGACATCCCAGAAGCCCAGCCCCCTCAACCCCTCTGGATGCTACATGGGGATGC
TGGACGGCTCAGCCCCCTGTTCCAAGGATTTTGGGGTGCTGAGATTCTCCCCTAGAGACCT
GAAATTCACCAGCTACAGATGCCAAATGACTTACATCTTAAGAAGTCTCAGAACGTCCAG
CCCTTCAGCAGCTCTCGTTCTGAGACATGAGCCTTGGGATGTGGCAGCATCAGTGGGACA
AGATGGACACTGGGCCACCCTCCCAGGCACCAGACACAGGGCACGGTGGAGAGACTTCTC
CCCCGTGGCCGCCTTGGCTCCCCCGTTTTGCCCGAGGCTGCTCTTCTGTGCACTTCCCTC
TTTGTACCACAGTGGCTCTGGGGCCAGGCCTGCCTGCCCACTGGCCATCGCCACCTTCCC
CAGCTGCCTCCTACCAGCAGTTTCTCTGAAGATCTGTCAACAGGTAAAGTCAATCTGGGG
CTTCCACTGCCTGCATTCCAGTCCCAGAGCTTGGTGGTCCCGAAACGGGAAGTACATAT
TGGGGCATGGTGGCCTCCGTGAGCAAATGGTGTCTTGGGCAATCTGAGGCCAGGACAGAT
GTTGCCCCACCCACTGGAGATGGTGTCTGAGGGAGGTGGGTGGGGCCTTCTGGGAAGGTGA
GTGGAGAGGGGCACCTGCCCCCGCCCTCCCATCCCTACTCCCACTGCTCAGCGCGGG
CCATTGCAAGGGTGCCACACAATGTCTTGTCCACCCTGGGACACTTCTGAGTATGAAGCG
GGATGCTATTAAAACTACATGGGGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AGA

207/392

FIGURE 198

MGVPTALEAGSWRWGSLFLFALFLAASLGPVAAFQVATPYSLYVCPEGQNVTLTCRLLGPV
DKGHDVTFYKWTYRSSRGEVQTCSERRPIRNLTFQDLHLHHGGHQAANTSHDLAQRHGLE
SASDHGNGFSITMRNLTLDDSGLYCCLVVEIRHHHSEHRVHGAMELQVQTGKDAPSNCVV
YPSSSQDSENITAAALATGACIVGILCLPLILLLVYKQRQAASNRRAQELVRMDSNIQGI
ENPGFEASPPAQGIPEAKVRHPLSYVAQRQPSESGRHLLSEPSTPLSPPGPGDVFFPSLD
PVPDSPNFEVI

Signal peptide:
amino acids 1-28

Transmembrane domain:
amino acids 190-216

208/392

FIGURE 199

CTAGCCTGCGCCAAGGGGTAGTGAGACCGCGCGGCAACAGCTTGCGGGCTGCGGGGAGCTC
CCGTGGGCGCTCCGCTGGCTGTGCAGGCGGCCATGGATTCCCTGCGGAAAATGCTGATCT
CAGTCGCAATGCTGGGCGCAGGGGCTGGCGTGGGCTACGCGCTCCTCGTTATCGTGACCC
CGGGAGAGCGGCGGAAGCAGGAAATGCTAAAGGAGATGCCACTGCAGGACCCAAGGAGCA
GGGAGGAGGCGGCCAGGACCCAGCAGCTATTGCTGGCCACTCTGCAGGAGGCAGCGACCA
CGCAGGAGAACGTGGCCTGGAGGAAGAACTGGATGGTTGGCGGCGAAGGCGGCGCCAGCG
GGAGGTCAACGTGAGACCGGACTTGCCCTCCGTGGGCGCCGGACCTTGGCTTGGGCGCAGG
AATCCGAGGCAGCCTTTCTCCTTCGTGGGCCCAGCGGAGAGTCCGGACCGAGATAACCATG
CCAGGACTCTCCGGGGTCTGTGAGCTGCCGTCCGGTGAGCACGTTTCCCCCAAACCTG
GACTGACTGCTTTAAGGTCCGCAAGGCGGGCCAGGGCCGAGACGCGAGTCGGATGTGGTG
AACTGAAAGAACCAATAAAATCATGTTCTCCAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

209/392

FIGURE 200

MDSLRLKMLISVAMLGAGAGVGYALLVIVTPGERRKQEMLKEMPLQDPRSREEAARTQQLL
LATLQEAATTQENVAWRKNWMVGEGGASGRSP

Signal peptide:
amino acids 1-18

210/392

FIGURE 201

GACAGCTGTGTCTCGATGGAGTAGACTCTCAGAACAGCGCAGTTTGGCCCTCCGCTCACGC
AGAGCCTCTCCGTGGCTTCCGCACCTTGAGCATTAGGCCAGTTCTCCTCTTCTCTCTAAT
CCATCCGTCACCTCTCCTGTCATCCGTTTCCATGCCGTGAGGTCCATTACAGAACACAT
CCATGGCTCTCATGCTCAGTTTGGTTCTGAGTCTCCTCAAGCTGGGATCAGGGCAGTGGC
AGGTGTTTGGGCCAGACAAGCCTGTCCAGGCCCTTGGTGGGGAGGACGCAGCATTCTCCT
GTTTCCTGTCTCCTAAGACCAATGCAGAGGCCATGGAAGTGCAGTTCTTCAAGGGCCAGT
TCTCTAGCGTGGTCCACCTCTACAGGGACGGGAAGGACCAGCCATTTATGCAGATGCCAC
AGTATCAAGGCAGGACAAAACTGGTGAAGGATTCTATTGCGGAGGGGCGCATCTCTCTGA
GGCTGGAAAACATTACTGTGTTGGATGCTGGCCTCTATGGGTGCAGGATTAGTTCCCAGT
CTTACTACCAGAAGGCCATCTGGGAGCTACAGGTGTCAGCACTGGGCTCAGTTCTCTCA
TTTCCATCACGGGATATGTTGATAGAGACATCCAGCTACTCTGTCAGTCTCGGGCTGGT
TCCCCCGGGCCACAGCGAAGTGGAAGGTCCACAAGGACAGGATTTGTCCACAGACTCCA
GGACAAACAGAGACATGCATGGCCTGTTTGATGTGGAGATCTCTCTGACCGTCCAAGAGA
ACGCCGGGAGCATATCCTGTTCCATGCGGCATGCTCATCTGAGCCGAGAGGTGGAATCCA
GGGTACAGATAGGAGATACCTTTTTTCGAGCCTATATCGTGGCACCTGGCTACCAAAGTAC
TGGGAATACTCTGCTGTGGCCTATTTTTTGGCATTGTTGGACTGAAGATTTTCTTCTCCA
AATTCCAGTGGAATAATCCAGGCGGAACTGGACTGGAGAAGAAAGCACGGACAGGCAGAAT
TGAGAGACGCCCGGAAACACGCAGTGGAGGTGACTCTGGATCCAGAGACGGCTCACCCGA
AGCTCTGCGTTTTCTGATCTGAAAACTGTAACCCATAGAAAAGCTCCCCAGGAGGTGCCTC
ACTCTGAGAAGAGATTTTACAAGGAAGAGTGTGGTGGCTTCTCAGAGTTTCCAAGCAGGGA
AACATTACTGGGAGGTGGACGGAGGACACAATAAAAGGTGGCGCGTGGGAGTGTGCCGGG
ATGATGTGGACAGGAGGAAGGAGTACGTGACTTTGTCTCCCGATCATGGGTACTGGGTCC
TCAGACTGAATGGAGAACATTTGTATTTTACATTAAATCCCCGTTTTATCAGCGTCTTCC
CCAGGACCCCACCTACAAAAATAGGGGTCTTCTTGACTATGAGTGTGGGACCATCTCCT
TCTTCAACATAAATGACCAGTCCCTTATTTATACCCTGACATGTCGTTTTGAAGGCTTAT
TGAGGCCCTACATTGAGTATCCGTCTATAATGAGCAAAATGGAATCCCATAGTCATCT
GCCCAGTCACCCAGGAATCAGAGAAAGAGGCCTCTTGCAAAAGGGCCTCTGCAATCCCAG
AGACAAGCAACAGTGAGTCTCCTCACAGGCAACCACGCCCTTCTCCCCAGGGGTGAAA
TGAGGATGAATCACATCCCACATTCTTCTTTAGGGATATTAAGGTCTCTCTCCAGATC
CAAAGTCCCGCAGCAGCCGGCCAAGGTGGCTTCCAGATGAAGGGGGACTGGCCTGTCCAC
ATGGGAGTCAGGTGTCATGGCTGCCCTGAGCTGGGAGGGAAGAAGGCTGACATTACATTT
AGTTTGCTCTCACTCCATCTGGCTAAGTGATCTTGAAATACCACCTCTCAGGTGAAGAAC
CGTCAGGAATTCCCATCTCACAGGCTGTGGTGTAGATTAAGTAGACAAGGAATGTGAATA
ATGCTTAGATCTTATTGATGACAGAGTGTATCCTAATGGTTTGTTCATTATATTACACTT
TCAGTAAAAAAA

211/392

FIGURE 202

MALMLSLVLSLLKLGSGQWQVFGPDKPVQALVGEDAAFSCFLSPKTNAEAMEVRFFRGQF
SSVHLYRDGKDQPFMQMPQYQGRTKLVKDSIAEGRISLRLENITVLDAGLYGCRISQS
YYQKAIWELQVSALGSVPLISITGYVDRDIQLLCQSSGWFPRPTAKWKGPQGQDLSTDSR
TNRDMHGLFDVEISLTVQENAGSISCSMRHAHLSREVESRVQIGDTFFEPISWHLATKVL
GILCCGLFFGIVGLKIFFSKFQWKIQAELDWRRKHGQAEIRDARKHAVEVTLDPETAHPK
LCVSDLKTVTHRKAQEVPHSEKRFTRKSVVASQSFGAGKHYWEVDGGHNKRWRVGVCRD
DVDRRKEYVTLSPDHGYWVLRNLNGEHLYFTLNPRFISVFPRTPTKIGVFLDYECGTISF
FNINDQSLIYTLTCRFEGLLRPYIEYPSYNEQNGTPIVICPVTQESEKEASWQRASAIPE
TSNSESSSQATTFFLPRGEM

Signal peptide:
amino acids 1-17

Transmembrane domain:
amino acids 239-255

212/392

FIGURE 203

TGCGGCGCAGTG TAGACCTGGGAGGATGGGCGGCCTGCTGCTGGCTGCTTTTCTGGCTTT
GGTCTCGGTGCCCAGGGCCCAGGCCGTGTGGTTGGGAAGACTGGACCCTGAGCAGCTTCT
TGGGCCCTGGTACGTGCTTGC GGTGGCCTCCCGGGAAAAGGGCTTTGCCATGGAGAAGGA
CATGAAGAACGTCGTGGGGGTGGTGGTGACCCTCACTCCAGAAAACAACCTGCGGACGCT
GTCCTCTCAGCACGGGCTGGGAGGGTGTGACCAGAGTGTCATGGACCTGATAAAGCGAAA
CTCCGGATGGGTGTTTGAGAATCCCTCAATAGGCGTGCTGGAGCTCTGGGTGCTGGCCAC
CAACTTCAGAGACTATGCCATCATCTTCACTCAGCTGGAGTTCGGGGACGAGCCCTTCAA
CACCGTGGAGCTGTACAGTCTGACGGAGACAGCCAGCCAGGAGGCCATGGGGCTCTTCAC
CAAGTGGAGCAGGAGCCTGGGCTTCCTGTACAGTAGCAGGCCCAGCTGCAGAAGGACCT
CACCTGTGCTCACAAGATCCTTCTGTGAGTGCTGCGTCCCCAGTAGGGATGGCGCCCA
GGGTCTGTGACCTCGGCCAGTGTCACCCACCTCGCTCAGCGGCTCCCGGGGCCAGCA
CCAGCTCAGAATAAAGCGATTCCACAGCA

213/392

FIGURE 204

MGGLLLAAFLALVSVPRQA VWLGRLDPEQLLGPWYVLAVASREKGFAMEKDMKNVVGVV
VTLTPENNLRTLSSQHGLGGCDQSVMDLIKRN SGWVFENPSIGVLELWVLATNFRDYAI I
FTQLEFGDEPFNTVELYSLTETASQEAMGLFTKWSRSLGFLSQ

Signal peptide:
amino acids 1-20

214/392

FIGURE 205

GACGCCCAGTGACCTGCCGAGGTCGGCAGCACAGAGCTCTGGAGATGAAGACCCTGTTCC
TGGGTGTCACGCTCGGCCTGGCCGCTGCCCTGTCCTTCACCCTGGAGGAGGAGGATATCA
CAGGGACCTGGTACGTGAAGGCCATGGTGGTCGATAAGGACTTTCCGGAGGACAGGAGGC
CCAGGAAGGTGTCCCCAGTGAAGGTGACAGCCCTGGGCGGTGGGAAGTTGGAAGCCACGT
TCACCTTCATGAGGGAGGATCGGTGCATCCAGAAGAAAATCCTGATGCGGAAGACGGAGG
AGCCTGGCAAATACAGCGCCTATGGGGGCAGGAAGCTCATGTACCTGCAGGAGCTGCCCCA
GGAGGGACCACTACATCTTTTACTGCAAAGACCAGCACCATGGGGGCCTGCTCCACATGG
GAAAGCTTGTGGGTAGGAATTCTGATACCAACCGGGAGGCCCTGGAAGAATTTAAGAAAT
TGGTGCAGCGCAAGGGACTCTCGGAGGAGGACATTTTCACGCCCCTGCAGACGGGAAGCT
GCGTTCCCGAACACTAGGCAGCCCCCGGGTCTGCACCTCCAGAGCCCACCCTACCACCAG
ACACAGAGCCCGGACCACCTGGACCTACCCTCCAGCCATGACCCTTCCCTGCTCCCACCC
ACCTGACTCCAAATAAAGTCCTTTTCCCCCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAA

215/392

FIGURE 206

MKTLFLGVTLGLAAALSFTLEEDITGTWYVKAMVVDKDFPEDRRPRKVSPVKVTALGGG
KLEATFTFMREDRCIQKKILMRKTEEPGKYSAYGGRKLMYLQELPRRDHYIFYCKDQHHG
GLLHMGKLVGRNSDTNREALEEFKKLVQRKGLSEEDIFTPLQTGSCVPEH

Important features:

Signal peptide:

amino acids 1-17

216/392

FIGURE 207

GTTCCGCAGATGCAGAGGTTGAGGTGGCTGCGGGACTGGAAGTCATCGGGCAGAGGTCTC
ACAGCAGCCAAGGAACCTGGGGCCCGCTCCTCCCCCTCCAGGCCATGAGGATTCTGCAG
TTAATCCTGCTTGCTCTGGCAACAGGGCTTGTAGGGGGAGAGACCAGGATCATCAAGGGG
TTCGAGTGCAAGCCTCACTCCCAGCCCTGGCAGGCAGCCCTGTTGAGAAGACGCGGCTA
CTCTGTGGGGCGACGCTCATCGCCCCAGATGGCTCCTGACAGCAGCCCACTGCCTCAAG
CCCCGCTACATAGTTACCTGGGGCAGCACAACCTCCAGAAGGAGGAGGGCTGTGAGCAG
ACCCGGACAGCCACTGAGTCCTTCCCCACCCCGGCTTCAACAACAGCCTCCCCAACAAA
GACCACCGCAATGACATCATGCTGGTGAAGATGGCATCGCCAGTCTCCATCACCTGGGCT
GTGCGACCCCTCACCTCTCCTCACGCTGTGTCACTGCTGGCACCAGCTGCCTCATTTCC
GGCTGGGGCAGCACGTCCAGCCCCAGTTACGCCTGCCTCACACCTTGCGATGCGCCAAC
ATCACCATCATTGAGCACCAGAAGTGTGAGAACGCCTACCCCGGCAACATCACAGACACC
ATGGTGTGTGCCAGCGTGCAGGAAGGGGGCAAGGACTCCTGCCAGGGTGACTCCGGGGGC
CCTCTGGTCTGTAACCAGTCTCTTCAAGGCATTATCTCCTGGGGCCAGGATCCGTGTGCG
ATCACCCGAAAGCCTGGTGTCTACACGAAAGTCTGCAAATATGTGGACTGGATCCAGGAG
ACGATGAAGAACAATTAGACTGGACCCACCCACCACAGCCCATCACCTCCATTTCCACT
TGGTGTTTGGTTCCTGTTCACTCTGTTAATAAGAAACCCTAAGCCAAGACCCTCTACGAA
CATTCCTTTGGGCCTCCTGGACTACAGGAGATGCTGTCACTTAATAATCAACCTGGGGTTC
GAAATCAGTGAGACCTGGATTCAAATTCTGCCTTGAAATATTGTGACTCTGGGAATGACA
ACACCTGGTTTGTCTCTGTTGTATCCCAGCCCCAAAGACAGCTCCTGGCCATATATCA
AGGTTTCAATAAATATTTGCTAAATGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAA

217/392

FIGURE 208

MRILQLILLALATGLVGGETRIIKGFECKPHSQPWQAALFEKTRLLCGATLIAPRWLLTA
AHCLKPRYIVHLGQHNLOKEEGCEQTRTATESFPHPGFNNSLPNKDHRNDIMLVKMASPV
SITWAVRPLTLSSRCVTAGTSCLISGWGSTSSPQLRLPHTLRCANITITIEHQKCENAYPG
NITDTMVCASVQEGGKDSCQGDSSGGLVLCNQSLOGIISWGQDPCAITRKPGVYTKVCKYV
DWIQETMKNN

Important features:**Signal peptide:**

amino acids 1-18

Serine proteases, trypsin family, histidine active site:

amino acids 58-63

N-glycosylation sites:

amino acids 99-102, 165-168, 181-184, 210-213

Glycosaminoglycan attachment site:

amino acids 145-148

Kringle domain proteins:

amino acids 197-209, 47-64

Serine proteases, trypsin family, histidine protein:

amino acids 199-209, 47-63, 220-243

Apple domain proteins:

amino acids 222-249, 189-222

218/392

FIGURE 209

GCGGCCACACGCAGCTAGCCGGAGCCCGGACCAGGCGCCTGTGCCTCCTCCTCGTCCCTC
GCCGCGTCCGCGAAGCCTGGAGCCGGCGGGAGCCCCGCGCTCGCCATGTCGGGCGAGCTC
AGCAACAGGTTCCAAGGAGGGAAGGCGTTCGGCTTGCTCAAAGCCCGGCAGGAGAGGAGG
CTGGCCGAGATCAACCGGGAGTTTCTGTGTGACCAGAAGTACAGTGATGAAGAGAACCTT
CCAGAAAAGCTCACAGCCTTCAAAGAGAAGTACATGGAGTTTGACCTGAACAATGAAGGC
GAGATTGACCTGATGTCTTTAAAGAGGATGATGGAGAAGCTTGGTGTCCCCAAGACCCAC
CTGGAGATGAAGAAGATGATCTCAGAGGTGACAGGAGGGGTGAGTGACACTATATCCTAC
CGAGACTTTGTGAACATGATGCTGGGGAAACGGTCGGCTGTCTCAAGTTAGTCATGATG
TTTGAAGGAAAAGCCAACGAGAGCAGCCCCAAGCCAGTTGGCCCCCTCCAGAGAGAGAC
ATTGCTAGCCTGCCCTGAGGACCCCGCCTGGACTCCCCAGCCTTCCACCCCATACCTCC
CTCCCGATCTTGCTGCCCTTCTTGACACACTGTGATCTCTCTCTCTCATTGTGTTGGT
CATTGAGGGTTTGTGTTGTGTTTTTCATCAATGTCTTTGTAAAGCACAAATTATCTGCCTTA
AAGGGGCTCTGGGTCGGGGAATCCTGAGCCTTGGGTCCCCTCCCTCTCTTCTTCCCTCCT
TCCCCGCTCCCTGTGCAGAAGGGCTGATATCAAACCAAAAAGTAGAGGGGGCAGGGCCAG
GGCAGGGAGGCTTCCAGCCTGTGTTCCCCCTCACTTGGAGGAACCAGCACTCTCCATCCTT
TCAGAAAAGTCTCCAAGCCAAGTTCAGGCTCACTGACCTGGCTCTGACGAGGACCCAGGC
CACTCTGAGAAGACCTTGGAGTAGGGACAAGGCTGCAGGGCCTCTTTTCGGGTTTCTTTGG
ACAGTGCCATGGTTCCAGTGCTCTGGTGTCAACCAGGACACAGCCACTCGGGGCCCCGCT
GCCCCAGCTGATCCCCACTCATTCCACACCTCTTCTCATCCTCAGTGATGTGAAGGTGGG
AAGGAAAGGAGCTTGGCATTTGGGAGCCCTTCAAGAAGGTACCAGAAGGAACCCTCCAGTC
CTGCTCTCTGGCCACACCTGTGCAGGCAGCTGAGAGGCAGCGTGCAGCCCTACTGTCCCT
TACTGGGGCAGCAGAGGGCTTCGGAGGCAGAAGTGAGGCCTGGGGTTTGGGGGGAAAGGT
CAGCTCAGTGCTGTTCCACCTTTTAGGGAGGATACTGAGGGGACCAGGATGGGAGAATGA
GGAGTAAAATGCTCACGGCAAAGTCAGCAGCACTGGTAAGCCAAGACTGAGAAATACAAG
GTTGCTTGTCTGACCCCAATCTGCTTGAAAAAAAAAAAAAAAAAAAAA

219/392

FIGURE 210

MSGELSNRFQGGKAFGLLKARQERRLAEGINREFLCDQKYSDEENLPEKLTAFFKEKYMEFD
LNNEGEIDLMSLKRMMEKLGVPKTHLEMKKMISEVTGGVSDTISYRDFVNMMLGKRSAVL
KLVMMFEGKANESSPKPVGPPPERDIASLP

220/392

FIGURE 211

CTGGGATCAGCCACTGCAGCTCCCTGAGCACTCTCTACAGAGACGCGGACCCCAGACATG
AGGAGGCTCCTCCTGGTCACCAGCCTGGTGGTTGTGCTGCTGTGGGAGGCAGGTGCAGTC
CCAGCACCCAAGGTCCCTATCAAGATGCAAGTCAAACACTGGCCCTCAGAGCAGGACCCA
GAGAAGGCCTGGGGCGCCCGTGTGGTGGAGCCTCCGGAGAAGGACGACCAGCTGGTGGTG
CTGTTCCCTGTCCAGAAGCCGAAACTCTTGACCACCGAGGAGAAGCCACGAGGTCAGGGC
AGGGGCCCCATCCTTCCAGGCACCAAGGCCTGGATGGAGACCGAGGACACCCTGGGCCGT
GTCCTGAGTCCCGAGCCCGACCATGACAGCCTGTACCACCCTCCGCCTGAGGAGGACCAG
GGCGAGGAGAGGCCCCGGTTGTGGGTGATGCCAAATCACCAGGTGCTCCTGGGACCGGAG
GAAGACCAAGACCACATCTACCACCCCCAGTAGGGGCTCCAGGGGCCATCACTGCCCCCGC
CCTGTCCCAAGGCCCAGGCTGTTGGGACTGGGACCCTCCCTACCCTGCCCCAGCTAGACA
AATAAACCCCAGCAGGCAAAAAAAAAAAAAAAAAAAAAA

221/392

FIGURE 212

MRRLLLVTSLVVLLWEAGAVPAPKVPIKMQVKHWPSEQDPEKAWGARVVEPPEKDDQLV
VLFPVQKPKLLTTEEKPRGQGGRGPILPGTKAWMETEDTLGRVLSPEPDHDSLYHPPPEED
QGEERPRLWVMPNHQVLLGPEEDQDHIYHPQ

222/392

FIGURE 213

CAGGCAGAAGCGAACAAAGACCCAGCAAGAGAAGGCAGAGGCTAAGACCCATCCCGTATC
TGCTCTCCTGAAATAATTCTGGAGTCATGCCTGAAATGCCAGAGGACATGGAGCAGGAGG
AAGTTAACATCCCTAATAGGAGGGTTCTGGTTACTGGTGCCACTGGGCTTCTTGGCAGAG
CTGTACACAAAGAATTTTACGACAGAATAATTGGCATGCAGTTGGCTGTGGTTTCAGAAGAG
CAAGACCAAAATTTGAACAGGTAACTCTGTTGGATTCTAATGCAGTTCATCACATCATTC
ATGATTTTTCAGCCCCATGTTATAGTACATTGTGCAGCAGAGAGAAGACCAGATGTTGTAG
AAAATCAGCCAGATGCTGCCTCTCAACTTAATGTGGATGCTTCTGGGAATTTAGCAAAGG
AAGCAGCTGCTGTTGGAGCATTCTCATCTACATTAGCTCAGATTATGTATTTGATGGAA
CAAATCCACCTTACAGAGAGGAAGACATACCAGCTCCCCTAAATTTGTATGGCAAACAA
AATTAGATGGAGAAAAGGCTGTCTGGAGAACAACTTAGGAGCTGCTGTTTTGAGGATTC
CTATTCTGTATGGGGAAGTTGAAAAGCTCGAAGAAAAGTGCTGTGACTGTTATGTTTGATA
AAGTGCAGTTCAGCAACAAGTCAGCAACATGGATCACTGGCAGCAGAGGTTCCCCACAC
ATGTCAAAGATGTGGCCACTGTGTGCCGCGCAGCTAGCAGAGAAGAGAATGCTGGATCCAT
CAATTAAGGGAACCTTTCACTGGTCTGGCAATGAACAGATGACTAAGTATGAAATGGCAT
GTGCAATTGCAGATGCCTTCAACCTCCCAGCAGTCACCTAAGACCTATTACTGACAGCC
CTGTCTTAGGAGCACACGTCGAGAAATGCTCAGCTTGACTGCTCCAAATTGGAGACCT
TGGGCATTGGCCAACGAACACCATTTTGAATTTGGAATCAAAGAATCACTTTGGCCTTTCC
TCATTGACAAGAGATGGAGACAAACGGTCTTTCATTAGTTTTATTTGTGTTGGGTTCTTTT
TTTTTTTTAAATGAAAAGTATAGTATGTGGCACTTTTTAAAGAACAAAGGAAATAGTTTTG
TATGAGTACTTTAATTGTGACTCTTAGGATCTTTCAGGTAAATGATGCTCTTGCACTAGT
GAAATTGTCTAAAGAACTAAAGGGCAGTCATGCCCTGTTTGCAGTAATTTTTCTTTTAA
TCATTTTGTGTTGTCCTGGCTAACTTGGAGTTTGAGTATAGTAAATTATGATCCTTAAAT
ATTTGAGAGTCAGGATGAAGCAGATCTGCTGTAGACTTTTCAGATGAAATTGTTTATTCT
CGTAACCTCCATATTTTCAGGATTTTGAAGCTGTTGACCTTTTCATGTTGATTATTTTA
AATTGTGTGAAATAGTATAAAAATCATTGGTGTTCATTATTTGCTTTGCCTGAGCTCAGA
TCAAAATGTTTGAAGAAAGGAACCTTTATTTTGAAGTTACGTACAGTTTTTATGCTTGA
GATATTTCAACATGTTATGTATATTGGAACCTCTACAGCTTGATGCCTCCTGCTTTTATA
GCAGTTTATGGGGAGCACTTGAAAGAGCGTGTGTACATGTATTTTTTTCTAGGCAAACA
TTGAATGCAAACGTGTATTTTTTTAATATAAATATAAATGTCCTTTTCATCCCATGTT
GCCGCTAAGTGATATTTTCATATGTGTGGTTATACTCATAAATAATGGGCCTTGTAAGTCTT
TTCACCATTCATGAATAATAAATAATATGTACTGCTGGCATGTAATGCTTAGTTTTCTTG
TATTTACTTCTTTTTTTAAATGTAAGGACCAAACTTCTAACTAATTGTTCTTTTGTTC
TTTAATTTTTTAAAAATTACATTCTTCTGATGTAACATGTGATACATAAAAAGAATATAG
TTTAATATGTATTGAAATAAAACACAATAAAAT

223/392

FIGURE 214

MPEMPEDMEQEEVNIPNRRVLVTGATGLLGRAVHKEFQQNNWHAVGCGFRRARPKFEQVN
LLDSNAVHHIIHDFQPHVIVHCAAERRPDVVENQPDAAASQLNVDASGNLAKEAAAVGAFL
IYISSDYVFDGTNPPYREEDIAPAPLNLYGKTKLDGEKAVLENNLGA AVLRIPILYGEVEK
LEESAVTVMFDKVQFSNKSANMDHWQQRFP THVKDVATVCRQLAEKRMLDPSIKGTFHWS
GNEQMTKYEMACAIADAFNLPSSHLRPITDSPVLGAQRPRNAQLDCSKLET LGIGQRTPF
RIGIKESLWPFLIDKRWRQTVFH

Signal peptide:
amino acids 1-30

Transmembrane domain:
amino acids 105-127

N-glycosylation site:
amino acids 197-201

N-myristoylation site:
amino acids 303-309

Short-chain dehydrogenases/reductases family proteins:
amino acids 18-30

224/392

FIGURE 215

GTGAATGTGAGGGTTTGATGACTTTCAGATGTCTAGGAACCAGAGTGGGTGCAGGGGCCC
CAGGCAGGGCTGATTCTTGGGCGGAGGAGAGTAGGGTAAAGGGTTCTGCATGAGCTCCTT
AAAGGACAAAGGTAACAGAGCCAGCGAGAGAGCTCGAGGGGAGACTTTGACTTCAAGCCA
CAGAATTGGTGAAGTGTGCGCGCCGCCGCCGCTCGCTCCTGCAGCGCTGTGACCTA
GCCGCTAGCATCTTCCCGAGCACCGGGATCCCGGGGTAGGAGGCGACGCGGGCGAGCACC
AGCGCCAGCCGGCTGCGGCTGCCACACGGCTCACCATGGGCTCCGGGCGCCGGGCGCTG
TCCGCGGTGCCGGCCGTGCTGCTGGTCCTCACGCTGCCGGGGCTGCCCGTCTGGGCACAG
AACGACACGGAGCCCATCGTGCTGGAGGGCAAGTGTCTGGTGGTGTGCGACTCGAACCCG
GCCACGGACTCCAAGGGCTCCTCTTCCCTCCCGCTGGGGATATCGGTCCGGGCGGCCAAC
TCCAAGGTCGCCCTTCTCGGCGGTGCGGAGCACCAACCACGAGCCATCCGAGATGAGCAAC
AAGACGCGCATCATTTACTTCGATCAGATCCTGGTGAATGTGGGTAAATTTTTTTCATTG
GAGTCTGTCTTTGTAGCACCAAGAAAAGGAATTTACAGTTTCAGTTTTTCACGTGATTAAA
GTCTACCAGAGCCAACTATCCAGGTAACTTGATGTTAAATGGAAAAACCAGTAATATCT
GCCTTTGCGGGGGACAAAGATGTTACTCGTGAAAGCTGCCACGAATGGTGTCTGCTCTAC
CTAGATAAAGAGGATAAGGTTTACCTAAAACTGGAGAAAGGTAATTTGGTTGGAGGCTGG
CAGTATTCCACGTTTTCTGGCTTTCTGGTGTTCCTCCCTATAGGATTCAATTTCTCCATGA
TGTTTCATCCAGGTGAGGGATGACCCACTCCTGAGTTATTGGAAGATCATTTTTTTCATCAT
TGGATTGATGTCTTTTATTGGTTTCTCATGGGTGGATATGGATTCTAAGGATTCTAGCCT
GTCTGAACCAATACAAAATTTACAGATTATTTGTGTGTGTCTGTTTCAGTATATTTGGA
TTGGGACTCTAAGCAGATAATACCTATGCTTAAATGTAACAGTCAAAGCTGTCTGCAAG
ACTTATTCTGAATTTTCATTTCTGGGATTACTGAATTAGTTACAGATGTGGAATTTTATT
TGTTTAGTTTTTAAAGACTGGCAACCAGGTCTAAGGATTAGAAAACCTAAAGTTCTGAC
TTCAATCAACGGTTAGTGTGATACTGCCAAAGAACTGTATACTGTGTTAATATATTGATT
ATATTTGTTTTTATTCCTTTGGAATTAGTTTGTGTTGGTTCTTGTAAAAACTTGGAATTTT
TTTTTTTCAGTAACTGGTATTATGTTTTCTCTTAAATAAGGTAATGAATGGCTTGCCAC
AAATTTACCTTGACTACGATATCATCGACATGACTTCTCTCAAAAAAAGAATGCTTCA
TAGTTGTATTTTAATTGTATATGTGAAAGAGTCATATTTTCCAAGTTATATTTTCTAAGA
AGAAGAATAGATCATAAATCTGACAAGGAAAAAGTTGCTTACCCAAAATCTAAGTGCTCA
ATCCCTGAGCCTCAGCAAAACAGCTCCCCTCCGAGGGAAATCTTATACTTTATTGCTCAA
CTTTAATTAAATGATTGATAATAACCACTTTATTAAAAACCTAAGGTTTTTTTTTTTTC
CGTAGACATGACCACTTTATTAAGTGGTGGGATGCTGTTGTTTCTAATTATACCTAT
TTTTCAAGGCTTCTGTTGTATTTGAAGTATCATCTGGTTTTGCCTTAACTCTTTAAATTG
TATATATTTATCTGTTTAGCTAATATTAAATTCAAATATCCCATATCTAAATTTAGTGCA
ATATCTTGTCTTTTGTATAGGTCATATGAATTCATAAAATTATTTATGTCTGTTATAGAA
TAAAGATTAATATATGTTAAAAAAA

225/392

FIGURE 216

MMSGRRALSAVPAVLLVLTLPGLPVWAQNDTEPIVLEGKCLVVCDSNPATDSKGSSSSPL
GISVRAANSKVAFSAVRSTNHEPSEMSNKTRI IYFDQILVNVGNFFTLESVVFVAPRKG IY
SFSFHVIKVYQSQTIQVNLMLNGKPVISAFAGDKDVTREAAATNGVLLYLDKEDKVYLKLE
KGNLVGGWQYSTFSGFLVFPL

Signal peptide:
amino acids 1-27

226/392

FIGURE 217

CGGCAACCAGCCGCCGCCACCACCGCTGCCACTGCCGCCCTGCCGGGGCCATGTTTCGCTC
TGGGCTTGCCCTTCTTGGTGCTCTTGGTGGCCTCGGTTCGAGAGCCATCTGGGGGTTCTGG
GGCCCAAGAACGTCTCGCAGAAAGACGCCGAGTTTGAGCGCACCTACGTGGACGAGGTCA
ACAGCGAGCTGGTCAACATCTACACCTTCAACCATACTGTGACCCGCAACAGGACAGAGG
GCGTGCGTGTGTCTGTGAACGTCTGAACAAGCAGAAGGGGGCGCCGTTGCTGTTTGTGG
TCCGCCAGAAGGAGGCTGTGGTGTCTTCCAGGTGCCCCTAATCCTGCGAGGGATGTTTC
AGCGCAAGTACCTCTACCAAAAAGTGGAACGAACCCTGTGTGAGCCCCCACCAAGAATG
AGTCGGAGATTTCAGTTCTTCTACGTGGATGTGTCCACCCTGTCACCAGTCAACACCACAT
ACCAGCTCCGGGTCAGCCGCATGGACGATTTTGTGCTCAGGACTGGGGAGCAGTTCAGCT
TCAATACCACAGCAGCACAGCCCCAGTACTTCAAGTATGAGTTCCTGAAGGCGTGGACT
CGGTAATTGTCAAGGTGACCTCCAACAAGGCCCTTCCCCTGCTCAGTCATCTCCATTCAGG
ATGTGCTGTGTCTGTCTATGACCTGGACAACAACGTAGCCTTCATCGGCATGTACCAGA
CGATGACCAAGAAGGCGGCCATCACCGTACAGCGCAAAGACTTCCCCAGCAACAGCTTTT
ATGTGGTGGTGGTGGTGAAGACCGAAGACCAAGCCTGCGGGGGCTCCCTGCCTTTCTACC
CCTTCGCAGAAGATGAACCGGTGATCAAGGGCACCGCCAGAAAACCTGTGAGTGCTGG
TGTCTCAAGCAGTCACGTCTGAGGCATACGTGAGTGGGATGCTCTTTTGCCTGGGTATAT
TTCTCTCCTTTTACCTGCTGACCGTCCTCCTGGCCTGCTGGGAGAAGTGGAGGCAGAAGA
AGAAGACCCTGCTGGTGGCCATTGACCGAGCCTGCCAGAAAGCGGTACCCCTCGAGTCC
TGGCTGATTCTTTTCCCTGGCAGTTCCCCTTATGAGGGTTACAACCTATGGCTCCTTTGAGA
ATGTTTCTGGATCTACCGATGGTCTGGTTGACAGCGCTGGCACTGGGGACCTCTCTTACG
GTTACCAGGGCCGCTCCTTTGAACCTGTAGGTACTCGGCCCCGAGTGGACTCCATGAGCT
CTGTGGAGGAGGATGACTACGACACATTGACCGACATCGATTCCGACAAGAATGTCATTCT
GCACCAAGCAATACCTCTATGTGGCTGACCTGGCACGGAAGGACAAGCGTGTCTGCGGA
AAAAGTACCAGATCTACTTCTGGAACATTGCCACCATTGCTGTCTTCTATGCCCTTCCTG
TGGTGCAGCTGGTGATCACCTACCAGACGGTGGTGAATGTCACAGGGAATCAGGACATCT
GCTACTACAACCTTCTGCGCCCAACCACTGGGCAATCTCAGCGCCTTCAACAACATCC
TCAGCAACCTGGGGTACATCCTGCTGGGGCTGCTTTTCCCTGCTCATCATCCTGCAACGGG
AGATCAACCACAACCGGGCCCTGCTGCGCAATGACCTCTGTGCCCTGGAATGTGGGATCC
CCAAACACTTTGGGCTTTTCTACGCCATGGGCACAGCCCTGATGATGGAGGGGCTGCTCA
GTGCTTGCTATCATGTGTGCCCCAACTATAACCAATTTCCAGTTTGACACATCGTTCATGT
ACATGATCGCCGACTCTGCATGCTGAAGCTCTACCAGAAGCGGCACCCGGACATCAACG
CCAGCGCCTACAGTGCCCTACGCCTGCCTGGCCATTGTGTCATCTTCTTCTGTGCTGGGCG
TGGTCTTTGGCAAAGGGAACACGGCGTTCTGGATCGTCTTCTCCATCATTCACATCATCG
CCACCCTGCTCCTCAGCACGCAGCTCTATTACATGGGCCGGTGGAAACTGGACTCGGGGA
TCTTCCGCCGCATCCTCCACGTGCTCTACACAGACTGCATCCGGCAGTGCAGCGGGCCGC
TCTACGTGGACCGCATGGTGTGCTGGTTCATGGGCAACGTTCATCAACTGGTCGCTGGCTG
CCTATGGGCTTATCATGCGCCCCAATGATTTGCTTCTTACTTGTGGCCATTGGCATCT
GCAACCTGCTCCTTTACTTCGCCTTCTACATCATCATGAAGCTCCGGAGTGGGGAGAGGA
TCAAGCTCATCCCCCTGCTCTGCATCGTTTGACCTCCGTGGTCTGGGGCTTCGCGCTCT
TCTTCTTCTTCCAGGGACTCAGCACCTGGCAGAAAACCCCTGCAGAGTCGAGGGAGCACA
ACCGGGACTGCATCCTCCTCGACTTCTTTGACGACCACGACATCTGGCACTTCTCTCCT
CCATCGCCATGTTTCGGGTCTTCTCCTGGTGTGCTGACACTGGATGACGACCTGGATACTG
TGCAGCGGGACAAGATCTATGTCTTCTAGCAGGAGCTGGGCCCTTCGCTTCACCTCAAGG
GGCCCTGAGCTCCTTTGTGTTCATAGACCGGTCACTCTGTGCTGCTGTGGGGATGAGTCCC
AGCACCGCTGCCAGCACTGGATGGCAGCAGGACAGCCAGGTCTAGCTTAGGCTTGGCCT
GGGACAGCCATGGGGTGGCATGGAACCTTGCAGCTGCCCTCTGCCGAGGAGCAGGCCTGC
TCCCCTGGAACCCCCAGATGTTGGCCAAATTGCTGCTTCTTCTCAGTGTGGGGCCTTC

227/392

CATGGGCCCCCTGTCCTTTGGCTCTCCATTTGTCCCTTTGCAAGAGGAAGGATGGAAGGGA
CACCTCCCCATTTTCATGCCTTGCAATTTGCCCCGTCTCCTCCCCACAATGCCCCAGCCT
GGGACCTAAGGCCTCTTTTCTCTCCATACTCCCACTCCAGGGCCTAGTCTGGGGCCTGA
ATCTCTGTCCTGTATCAGGGCCCCAGTTCTCTTTGGGCTGTCCCTGGCTGCCATCACTGC
CCATTCCAGTCAGCCAGGATGGATGGGGGTATGAGATTTTGGGGGTGGCCAGCTGGTGC
CAGACTTTTGGTGCTAAGGCCTGCAAGGGGCCTGGGGCAGTGCGTATTCTCTTCCCTCTG
ACCTGTGCTCAGGGCTGGCTCTTTAGCAATGCGCTCAGCCCAATTTGAGAACCGCCTTCT
GATTCAAGAGGCTGAATTCAGAGGTCACCTCTTCATCCCATCAGCTCCCAGACTGATGCC
AGCACCAGGACTGGAGGGAGAAGCGCCTCACCCCTTCCCTTCCTTCTTTCCAGGCCCTTA
GTCTTGCCAAACCCCAGCTGGTGGCCTTTTCAGTGCCATTGACACTGCCCAAGAATGTCCA
GGGGCAAAGGAGGGATGATACAGAGTTCAGCCCGTTCTGCCTCCACAGCTGTGGGCACCC
CAGTGCCCTACCTTAGAAAGGGGCTTCAGGAAGGGATGTGCTGTTTCCCTCTACGTGCCCA
GTCCTAGCCTCGCTCTAGGACCCAGGGCTGGCTTCTAAGTTTCCGTCCAGTCTTCAGGCA
AGTTCTGTGTTAGTCATGCACACACATACCTATGAAACCTTGGAGTTTACAAAGAATTGC
CCCAGCTCTGGGCACCCCTGGCCACCCTGGTCCTTGATCCCCTTCGTCCACCTGGTCCA
CCCCAGATGCTGAGGATGGGGGAGCTCAGGCGGGGCCTCTGCTTTGGGGATGGGAATGTG
TTTTTCTCCCAAACCTTGTTTTTATAGCTCTGCTTGAAGGGCTGGGAGATGAGGTGGGTCT
GGATCTTTTCTCAGAGCGTCTCCATGCTATGGTTGCATTTCCGTTTTCTATGAATGAATT
TGCAATCAATAAACAACCAGACTCAAAAAAAAAAAAAA

228/392

FIGURE 218

MFALGLPFLVLLVASVESHLGVLGPKNVSQKDAEFERTYVDEVNSELVNIYTFNHTVTRN
RTEGVRVSVNVLNKQKGAPLLFVVRQKEAVVSFQVPLILRGMFQRKYLYQKVERTLCQPP
TKNESEIQFFYVDVSTLSPVNTTYQLRVSRMDDFVLRTEGEQFSFNTTAAQPQYFKYEFPE
GVDSVIVKVTSNKAFPCSVISIQDVLCPVYDLNNAVFIGMYQTMKKAAITVQRKDFPS
NSFYVVVVVKTEDQACGGS LPFYPPFAEDFVDQGHRQKTL SVLV SQAVTSEAYVSGMLFC
LGIFLSFYLLTVLLACWENWRQKKKTLLVAIDRACPESGHPRVLADSFPGSSPYEGYNYG
SFENVSGSTDGLVDSAGTGDLSYGYQGRSFEPVGTRPRVDSMSSVEEDDYDTLTDIDSDK
NVIRTKQYLYVADLARKDKRVLRRKKYQIYFVNIATIAVFYALPVVQLVITYQTVVNVNVTGN
QDICYYNFLCAHPLGNLSAFNNILSNLGYILLGLLFLLIILQREINHNRALLRNDLCALC
CGIPKHFGLFYAMGTALMMEGLLSACYHVC PNYTNFQFDTSFMYMIAGLCMLKLYQKRHP
DINASAYSAYACLAIVIFFSVLGVVFGKGNTAFWIVFSIIHIIATLLLSTQLYYMGRWKL
DSGIFRRILHVLTYTDCIRQCSGPLYVDRMVLVVMGNVINWSLAAYGLIMRPNDFASYLLA
IGICNLLLYFAFYIIMKLRSGERIKLIPLLCIVCTSVVWGFALFFFFQGLSTWQKTPAES
REHNRDCILLDFDDHDIWHFLSSIAMFGSFLVLLTLDDDDLDTVQRDKIYVF

Important features of the protein:**Signal peptide:**

amino acids 1-18

Transmembrane domains:

amino acids 292-317, 451-470, 501-520, 607-627, 751-770

Leucine zipper pattern:

amino acids 497-518

N-glycosylation sites:amino acids 27-30, 54-57, 60-63, 123-126, 141-144, 165-168,
364-367, 476-479, 496-499, 572-575, 603-606, 699-702

229/392

FIGURE 219

AATTTTTCACCAAGAGTAAACTTGGAGAAACCAACTGGACCTTGAGTATTGTACATTTTGCCTCGTGGACCCAAAGGTAGCAATCTGAAACATGAGGAGTACGATTCTACTGTTTTGTCTTCTAGGATCAACTCGGTCATTACCACAGCTCAAACCTGCTTTGGGACTCCCTCCACAAAAC TGGCTCCGGATCAGGGAACACTACCAAACCAACAGCAGTCAAATCAGGTCTTTCCTTCTTTAAGTCTGATACCATTAAACACAGATGCTCACACTGGGGCCAGATCTGCATCTGTTAAATCCTGCTGCAGGAATGACACCTGGTACCAGACCCACCCATTGACCTGGGAGGGTTGAATGTACAACAGCAACTGCACCCACATGTGTTACCAATTTTTTGTACACAACTTGGAGCCCAGG GCACTATCCTAAGCTCAGAGGAATTGCCACAAATCTTCACGAGCCTCATCATCCATTCTCTGTTCCCGGGAGGCATCCTGCCACCAGTCAGGCAGGGGCTAATCCAGATGTCCAGGATGGAAGCCTTCCAGCAGGAGGAGCAGGTGTAAATCCTGCCACCAGGGAACCCCAGCAGGCC GCCTCCCAACTCCAGTGGCACAGATGACGACTTTCGAGTGACCAACCCTGCAGGCATCCAAAGGAGCACACATGCCATCGAGGAAGCCACCACAGAATCAGCAAATGGAATTCAAGTAAGCTGTTTCAAATTTTTTTCAACTAAGCTGCCTCGAATTTGGTGATACATGTGAATCTTTATCATTTGATTATATTATGGAATAGATTGAGACACATTGGATAGTCTTAGAAGAAATTAATCTCTAATTTACCTGAAAATATTTCTTGAAATTTAGAAAATATGTTCTATGTAGAGAAATCCCAA CTTTTAAAAACAATAATTCAATGGATAAATCTGTCTTTGAAATATAACATTATGCTGCCTGGATGATATGCATATTTAAACATATTTTGGAAAACCTGGAAAAA

230/392

FIGURE 220

MRSTILLFCLLGSTRSLPQLKPALGLPPTKLAPDQGTLPNQQQSNQVFPSLSLIPLTQML
TLGPDHLHLLNPAAGMTPGTQTHPLTLGGLNVQQQLHPHVLPIFVTQLGAQGTTLSSEELP
QIFTSLIIHSLFPGGILPTSQAGANPDVQDGSLPAGGAGVNPATQGTPAGRLPTPSGTDD
DFAVTTPAGIQRSTHAIEEATTESANGIQ

Signal peptide:
amino acids 1-16

231/392

FIGURE 221

[illegible]

232/392

FIGURE 222

MARMSFVIAACQLVLGLLMTSLTESSIONSECPQLCVCEIRPWFTPQSTYREATTVDCND
LRLTRIPSNLSSDTQVLLQLQSNNIKTVDELQQLFNLTELDIFSQNNFTNIKEVGLANLTQ
LTTLHLEENQITEMTDYCLQDLSNLQELYINHNQISTISAHAFAGLKNLLRLHLNSNKLK
VIDSRWFDSTPNLEILMIGENPVIGILDMNFKPLANLRSLVLAGMYLTDIPGNALVGLDS
LESLSFYDNKLVKVPQLALQKVPNLKFLDLNKNPIHKIQEGDFKNMLRLKELGINNMGEL
VSVDRYALDNLPELTKLEATNNPKLSYIHLAFRSVPALSLMLNNNALNAIYQKTVESL
PNLREISISHSNPLRCDLCVIHWINSNKTNIIRFMEPLSMFCAMPPEYKQGHQVKEVLIQDSSE
QCLPMISHDSFPNRLNVDIGTTVFLDCRAMAEPEPEIYWVTPIGNKITVETLSDKYKLSS
EGTLEISNIQIEDSGRYTCVAQNVQGADTRVATIKVNGTLLDGTQVLKIYVKQTESHSL
VSWKVNSNVMTSNLKWSSATMKIDNPHITYTARVPVDVHEYNLTHLQPSDYEVCCLTVSN
IHQQTQKSCVNVTTKNAFAVDISDQETSTALAAVMGSMFAVISLASIAVYFAKRFRKN
YHHSLLKKYMQKTSSIPLNELYPPPLINLWEGDSEKDKDGSADTKPTQVDTSRSYMW

Important features:**Signal peptide:**

Amino acids 1-25

Transmembrane domain:

Amino acids 508-530

N-glycosylation sites:Amino acids 69-73;96-100;106-110;117-121;385-389;517-521;
582-586;611-615**Tyrosine kinase phosphorylation site:**

Amino acids 573-582

N-myristoylation sites:

Amino acids 16-22;224-230;464-470;637-643;698-704

233/392

FIGURE 223

[illegible]

234/392

FIGURE 224

MASYLYGVLFVAVGLCAPIYCVSPANAPSAYPRPSSTKSTPASQVYSLNTDFAFRLYRRLV
LETPSQNIFFSPVSVSTSLAMLSLGAHSVTKTQILQGLGFNLTHTPESAIHQGFQHLVHS
LTVPSKDLTLKMGSALFVKKELQLQANFLGNVKRLYEAEVFSTDFSNPSIAQARINSHVK
KKTQGKVVDIIQGLDLLTAMVLVNHIFFKAKWEKPFHLEYTRKNFPFLVGEQVTQVPMM
HQKEQFAFGVDTELNCFVLQMDYKGDVAFFVLPSKGKMRQLEQALSARTLIKWSHSLQK
RWIEVFIPRFSISASYNLETILPKMGIQNAFDKNADFSGIKRDSLQVSKATHKAVLDVS
EEGTEATAATTTKFIVRSKDGPSYFTVSNRRTFLMMITNKATDGILFLGKVENPTKS

Signal peptide:
amino acids 1-20

235/392

FIGURE 225

GGGAGAGAGGATAAATAGCAGCGTGGCTTCCCTGGCTCCTCTCTGCATCCTTCCCGACCT
TCCCAGCAATATGCATCTTGACAGTCTGGTCGGCTCCTGCTCCCTCCTTCTGCTACTGGG
GGCCCTGTCTGGATGGGCGGCCAGCGATGACCCATTGAGAAGGTCATTGAAGGGATCAA
CCGAGGGCTGAGCAATGCAGAGAGAGAGGTGGGCAAGGCCCTGGATGGCATCAACAGTGG
AATCACGCATGCCGGAAGGGAAGTGGAGAAGGTTTTCAACGGACTTAGCAACATGGGGAG
CCACACCGGCAAGGAGTTGGACAAAGGCGTCCAGGGGCTCAACCACGGCATGGACAAGGT
TGCCCATGAGATCAACCATGGTATTGGACAAGCAGGAAAGGAAGCAGAGAAGCTTGGCCA
TGGGGTCAACAACGCTGCTGGACAGGCCGGGAAGGAAGCAGACAAAGCGGTCCAAGGGTT
CCCACTGGGGTCCACCAGGCTGGGAAGGAAGCAGAGAACTTGGCCAAGGGGTCAACCA
TGCTGCTGACCAGGCTGGAAAGGAAGTGGAGAAGCTTGGCCAAGGTGCCCACCATGCTGC
TGGCCAGGCCGGGAAGGAGCTGCAGAATGCTCATAATGGGGTCAACCAAGCCAGCAAGGA
GGCCAACCAGCTGCTGAATGGCAACCATCAAAGCGGATCTTCCAGCCATCAAGGAGGGGC
CACAACCACGCCGTTAGCCTCTGGGGCCTCAGTCAACACGCCTTTCATCAACCTTCCCGC
CCTGTGGAGGAGCGTCGCCAACATCATGCCCTAAACTGGCATCCGGCCTTGCTGGGAGAA
TAATGTCGCCGTTGTCACATCAGCTGACATGACCTGGAGGGGTTGGGGGTGGGGGACAGG
TTTCTGAAATCCCTGAAGGGGGTTGTACTGGGATTTGTGAATAAACTTGATACACCA

236/392

FIGURE 226

MHLARLVGSCSLLLLLGALSGWAASDDPIEKVIEGINRGLSNAEREVGKALDGINSGITH
AGREVEKVFENGLSNMGSHGTGKELDKGVQGLNHGMDKVAHEINHGIGQAGKEAEKLGHG
VN
NAAGQAGKEADKAVQGFHTGVHQAGKEAEKLGQGVNHAADQAGKEVEKLGQGAHHAAGQA
GKELQNAHNGVNVQASKEANQLLNHGHQSGSSSHQGGATTTPLASGASVNTPFINLPALWR
SVANIMP

Important features of the protein:

Signal peptide:

amino acids 1-25

Homologous region to circumsporozoite (CS) repeats:

amino acids 35-225

237/392

FIGURE 227

GAAGTAGAGGTGTTGTGCTGAGCGGCGCTCGGCGAACTGTGTGGACCGTCTGCTGGGACT
CCGGCCCTGCGTCCGCTCAGCCCCGTGGCCCCGCGCACCTACTGCCATGGAGACGCGGCC
TCGTCTCGGGGCCACCTGTTTGCTGGGCTTCAGTTTCCTGCTCCTCGTCATCTCTTCTGA
TGGACATAATGGGCTTGGAAAGGGTTTTGGAGATCATATTCATTGGAGGACACTGGAAGA
TGGGAAGAAAGAAGCAGCTGCCAGTGGACTGCCCCCTGATGGTGATTATTCATAAATCCTG
GTGTGGAGCTTGCAAAGCTCTAAAGCCCAAATTTGCAGAATCTACGGAAATTTCAGAACT
CTCCCATAATTTTGTTATGGTAAATCTTGAGGATGAAGAGGAACCCAAAGATGAAGATTT
CAGCCCTGACGGGGGTTATATTCCACGAATCCTTTTTCTGGATCCCAGTGGCAAGGTGCA
TCCTGAAATCATCAATGAGAATGGAAACCCAGCTACAAGTATTTTTATGTCAGTGCCGA
GCAAGTTGTTCAAGGGATGAAGGAAGCTCAGGAAAGGCTGACGGGTGATGCCTTCAGAAA
GAAACATCTTGAAGATGAATTGTAACATGAATGTGCCCCTTCTTTCATCAGAGTTAGTGT
TCTGGAAGGAAAGCAGCAGGGAAGGGAATATTGAGGAATCATCTAGAACAATTAAGCCGA
CCAGGAAACCTCATTCCTACCTACACTGGAAGGAGCGCTCTCACTGTGGAAGAGTTCTGC
TAACAGAAGCTGGTCTGCATGTTTGTGGATCCAGCGGAGAGTGGCAGACTTTCTTCTCCT
TTCCCTCTCACCTAAATGTCAACTTGTCAATTGAATGTAAAGAATGAAACCTTCTGACAC
AAAA

238/392

FIGURE 228

METRPRLGATCLLGFSFLLLVISSDGHNLGKGFGDHIHWRTLEDGKKEAAASGLPLMVI
IHKSWCGACKALKPKFAESTEISELSHNFVMVNLEDEEPEKDEDFSPDGGYIPRILFLDP
SGKVHPEIINENGNPYKYFYVSAEQVVQGMKEAQERLTGDAFRKKHLEDEL

Signal peptide:
Amino acids 1-23

Thioredoxin family proteins Homology Block:
Amino acids 58-75

239/392

FIGURE 229

CCCACGCGTCCGCCCACGCGTCCGGGTGCCACTCGCGCGCCGGCCGCGCTCCGGGCTTCT
CTTTTCCCTCCGACGCGCCACGGCTGCCCAGACATTCCGGCTGCCGGGTCTGGAGAGCTC
CCCGAACCCTCCGCGGAGAGGAGCGAGGCGGCGCCAGGGTGGCCCCCGGGGCGCGCTTG
GTCTCGGAGAAGCGGGGACGAGGCCGGAGGATGAGCGACTGAGGGCGACGCGGGCACTGA
CGCGAGTTGGGGCCGCGACTACCGGCAGCTGACAGCGCGATGAGCGACTCCCCAGAGACG
CCCTAGCCCAGGTGTGCGCGCCAGGCGGAGCGCGCAGGTGGGGCTGGGCTGTTAGTGGTCC
GCCCCACGCGGGTCGCCCGCCGGCCCCAGGATGGGCGCTGGCAACCCGGGCCCCGCGCCCCG
CGCTGCTACCCCTGCGCCCCGCTGCGAGCCCGGCGTCCGGCCCCGCGCCCTGCGCTCATGGA
CGGCGGCTCCCGGCTGGCGGCGGCGCGCCCCCGGGCTGTGAATGCGACTCGCCCCCTCGGC
CGCGCTCCCCGCCCCGCCGCCCGGGACGTGGTAGGGGATGCCAGCTCCACTGCGAT
GGCAGTTGGCGCGCTCTCCAGTTCCCTCCTGGTCACTGCTGCCTGATGGTGGCTCTGTG
CAGTCCGAGCATCCCGCTGGAGAAGCTGGCCCAGGCACCAGAGCAGCCGGGCCAGGAGAA
GCGTGAGCACGCCACTCGGGACGGCCCGGGGCGGGTGAACGAGCTCGGGCGCCCGGCGAG
GGACGAGGGCGGCAGCGGCCGGGACTGGAAGAGCAAGAGCGGCCGTGGGCTCGCCGGCCG
TGAGCCGTGGAGCAAGCTGAAGCAGGCCTGGGTCTCCAGGGCGGGGGCGCCAAGGCCGG
GGATCTGCAGGTCCGGCCCCGCGGGGACACCCCGCAGGCGGAAGCCCTGGCCGCAGCCGC
CCAGGACGCGATTGGCCCGGAACCTCGCGCCCACGCCCGAGCCACCCGAGGAGTACGTGTA
CCCGGACTACCGTGGCAAGGGCTGCGTGGACGAGAGCGGCTTCGTGTACGCGATCGGGGA
GAAGTTCGCGCCGGGCCCCCTCGGCCTGCCCGTGCCTGTGCACCGAGGAGGGGCGCTGTG
CGCGCAGCCCGAGTGCCCGAGGCTGCACCCGCGCTGCATCCACGTCGACACGAGCCAGTG
CTGCCCCGAGTGCAAGGAGAGGAAGAACTACTGCGAGTTCGGGGCAAGACCTATCAGAC
TTTGAGGAGTTCGTGGTGTCTCCATGCGAGAGGTGTCGCTGTGAAGCCAACGGTGAGGT
GCTATGCACAGTGTGAGCGTGTCCCCAGACGGAGTGTGTGGACCCTGTGTACGAGCCTGA
TCAGTGCTGTCCCATCTGCAAAAATGGTCCAAACTGCTTTGCAGAAACCGCGGTGATCCC
TGCTGGCAGAGAAGTGAAGACTGACGAGTGCACCATATGCCACTGTACTTATGAGGAAGG
CACATGGAGAATCGAGCGGCAGGCCATGTGCACGAGACATGAATGCAGGCAAATGTAGAC
GCTTCCCAGAACACAAACTCTGACTTTTTCTAGAACATTTTACTGATGTGAACATTCTAG
ATGACTCTGGGAACATCAGTCAAAGAAGACTTTTGATGAGGAATAATGAAAATTGTTG
GTACTTTTCTTTTCTTGATAACAGTTACTACAACAGAAGGAAATGGATATATTTCAAAA
CATCAACAAGAACTTTGGGCATAAAATCCTTCTCTAAATAAATGTGCTATTTTACAGTA
AGTACACAAAAGTACACTATTATATATCAAATGTATTTCTATAATCCCTCATTAGAGAG
CTTATATAAGTGTTTTCTATAGATGCAGATTAAAAATGCTGTGTTGTCAACCGTCAAAAA
AAAAAAAAAAAAAAAAAAAAA

240/392

FIGURE 230

MPSSTAMAVGALSSSLLVTCCLMVALCSPSIPLEKLAQAPEQPGQEKREHATRDGPGRVN
ELGRPARDEGGSGRDWKS KSGRGLAGREPWSKLKQAWVSQGGGAKAGDLQVRPRGDTPOA
EALAAAAQDAIGPELAPTPEPPPEEYVYPDYRGKGCVDSESGFVYAIGEFAPGPSACPCLC
TEEGPLCAQPECPRLHPRCIHVDTSQCCPQCKERKNYCEFRGKTYQTLEEFVVSPCERC
CEANGEVLCTVSACPQTECVDPVYEPDQCCPICKNGPNCFAETAVIPAGREVKTDECTIC
HCTYEEGTWRIERQAMCTRHECRQM

Important features of the protein:**Signal peptide:**

amino acids 1-27

Transmembrane domain:

amino acids 11-30

Glycosaminoglycan attachment site:

amino acids 80-83

N-myristoylation sites:

amino acids 10-15, 102-107, 103-108

Cell attachment sequence:

amino acids 114-117

EGF-like domain cysteine pattern signature:

amino acids 176-187

241/392

FIGURE 231

GGCCGGACGCCTCCGCGTTACGGGATGAATTAACGGCGGGTTCCGCACGGAGGTTGTGAC
CCCTACGGAGCCCCAGCTTGCCACGCACCCACTCGGCGTCGCGCGGCGTGCCCTGCTT
GTCACAGGTGGGAGGCTGGAACATCAGGCTGAAAAACAGAGTGGGTACTCTCTTCTGGG
AAGCTGGCAACAAATGGATGATGTGATATATGCATTCCAGGGGAAGGGAATTGTGGTGC
TTCTGAACCCATGGTCAATTAACGAGGCAGTTTCTAGCTACTGCACGTACTTCATAAAGC
AGGACTCTAAAAGCTTTGGAATCATGGTGTGATGGAAAGGGATTTACTTTTATACTGACTC
TGTTTTGGGGAAGCTTTTTTGGGAAGCATTTCATGCTGAGTCCCTTTTTACCTTTGATGT
TTGTAAACCCATCTTGGTATCGCTGGATCAACAACCGCCTTGTGGCAACATGGCTCACCC
TACCTGTGGCATTATTGGAGACCATGTTTGGTGTAAAAGTGATTATAACTGGGGATGCAT
TTGTTCCCTGGAGAAAGAAGTGTCAATTATCATGAACCATCGGACAAGAATGGACTGGATGT
TCCTGTGGAATTGCCTGATGCGATATAGCTACCTCAGATTGGAGAAAAATTTGCCTCAAAG
CGAGTCTCAAAGGTGTTCCCTGGATTGTTGGTGGGCCATGCAGGCTGCTGCCTATATCTTCA
TTCATAGGAAATGGAAGGATGACAAGAGCCATTTTGAAGACATGATTGATTACTTTTGTG
ATATTCACGAACCACTTCAACTCCTCATATTTCCAGAAGGGACTGATCTCACAGAAAACA
GCAAGTCTCGAAGTAATGCATTTGCTGAAAAAATGGACTTCAGAAATATGAATATGTTT
TACATCCAAGAACTACAGGCTTTACTTTTGTGGTAGACCGTCTAAGAGAAGGTAAGAACC
TTGATGCTGTCCATGATATCACTGTGGCGTATCCTCACAAACATTCCTCAATCAGAGAAGC
ACCTCCTCCAAGGAGACTTTCCAGGGGAAATCCACTTTCACGTCCACCGGTATCCAATAG
ACACCTTCCCCACATCCAAGGAGGACCTTCAACTCTGGTGCCACAAACGGTGGGAAGAGA
AAGAAGAGAGGCTGCGTTCCTTCTATCAAGGGGAGAAGAATTTTTATTTTACCGGACAGA
GTGTCATTCCACCTTGCAAGTCTGAACTCAGGGTCCTTGTGGTCAAATTGCTCTCTATAC
TGTATTGGACCCTGTTTCAAGCCCTGCAATGTGCCTACTCATATATTTGTACAGTCTTGTTA
AGTGGTATTTTATAATCACCATTGTAATCTTTGTGCTGCAAGAGAGAATATTTGGTGGAC
TGGAGATCATAGAACTTGATGTTACCGACTTTTACACAAACAGCCACATTTAAATTCAA
AGAAAAATGAGTAAGATTATAAGGTTTGCCATGTGAAAACCTAGAGCATATTTTGGAAT
GTTCTAAACCTTTCTAAGCTCAGATGCATTTTGCATGACTATGTCGAATATTTCTTACT
GCCATCATATTTTGTAAAGATATTTTGCATTAATTTTGTGGGAAAAATATTGCTACAA
TTTTTTTTTAATCTCTGAATGTAATTTGATACTGTGTACATAGCAGGAGTGATCGGGGT
GAAATAACTTGGGCCAGAATATTATTAAACAATCATCAGGCTTTTAA

242/392

FIGURE 232

MHSRGREIVVLLNPWSINEAVSSYCTYFIKQDSKSFSGIMVSWKGIYFILTLFWGSFFGSI
FMLSPFLPLMFVNPSWYRWINNRLVATWLTLPVALLETMFGVKVIITGDAFVPGERSVII
MNHRTMDWMFLWNCLMRYSYLRLEKICLKASLKGVPFGWAMQAAAYIFIHRKWKDDKS
HFEDMIDYFCDIHEPLQLLIFPEGTDLTENSKSRSNFAEKNGLQKYEYVLHPRTTGFTF
VVDRLREGKNLDAVHDITVAYPHNIPQSEKHLQGDFFPREIHFHVHRYPIDTLPTSKEDL
QLWCHKRWEEKERLRSFYQGEKNFYFTGQSVIPPCKSELRVLVVKLLSILYWTLFSPAM
CLLIYLYSLVKWYFIITIVIFVLQERIFGGLEIIEIACRYLLHKQPHLNSKKNE

Important features of the protein:

Signal peptide:

amino acids 1-22

Transmembrane domains:

amino acids 44-63, 90-108, 354-377

243/392

FIGURE 233

CGGCTCGAGTGCAGCTGTGGGGAGATTTCAAGTGCATTGCCTCCCCTGGGTGCTCTTCATC
TTGGATTTGAAAAGTTGAGAGCAGCATGTTTTGCCCACTGAAACTCATCCTGCTGCCAGTG
TACTGGATTATTCCTTGGGCCTGAATGACTTGAATGTTTTCCCGCCTGAGCTAACAGTC
CATGTGGGTGATTCACTCTGATGGGATGTGTTTTCCAGAGCACAGAAGACAAATGTATA
TTCAAGATAGACTGGACTCTGTACCAGGAGAGCACGCCAAGGACGAATATGTGCTATAC
TATTACTCCAATCTCAGTGTGCCTATTGGGCGCTTCCAGAACC CGGTACACTTGATGGGG
GACATCTTATGCAATGATGGCTCTCTCCTGCTCCAAGATGTGCAAGAGGCTGACCAGGGA
ACCTATATCTGTGAAATCCGCCTCAAAGGGGAGAGCCAGGTGTTCAAGAAGGCGGTGGTA
CTGCATGTGCTTCCAGAGGAGCCCAAAGAGCTCATGGTCCATGTGGGTGGATTGATTAG
ATGGGATGTGTTTTCCAGAGCACAGAAGTGAACACGTGACCAAGGTAGAATGGATATTT
TCAGGACGGCGCGCAAAGGAGGAGATTGTATTTCTGTTACTACCACAACTCAGGATGTCT
GTGGAGTACTCCAGAGCTGGGGCCACTTCCAGAATCGTGTGAACCTGGTGGGGGACATT
TTCCGCAATGACGGTTCATCATGCTTCAAGGAGTGAGGGAGTCAGATGGAGGAACTAC
ACCTGCAGTATCCACCTAGGGAACCTGGTGTTCAGAAAACCATTGTGCTGCATGTCAGC
CCGGAAGAGCCTCGAACACTGGTGACCCCGGCAGCCCTGAGGCCTCTGGTCTTGGGTGGT
AATCAGTTGGTGATCATTGTGGGAATTGTCTGTGCCACAATCCTGCTGCTCCCTGTTCTG
ATATTGATCGTGAAGAAGACCTGTGGAAATAAGAGTTCAGTGAATTCTACAGTCTTGGTG
AAGAACACGAAGAAGACTAATCCAGAGATAAAAGAAAAACCCTGCCATTTTGAAAGATGT
GAAGGGGAGAAACACATTTACTCCCAATAATTGTACGGGAGGTGATCGAGGAAGAAGAA
CCAAGTGAAAAATCAGAGGCCACCTACATGACCATGCACCCAGTTTGGCCTTCTCTGAGG
TCAGATCGGAACAACTCACTTGAAAAAAGTCAGGTGGGGGAATGCCAAAAACACAGCAA
GCCTTTTTGAGAAGAATGGAGAGTCCCTTCATCTCAGCAGCGGTGGAGACTCTCTCCTGTG
TGTGTCCTGGGCCACTCTACCAGTGATTTCAAGACTCCCGCTCTCCCAGCTGTCCTCCTGT
CTCATTGTTTGGTCAATACACTGAAGATGGAGAATTTGGAGCCTGGCAGAGAGACTGGAC
AGCTCTGGAGGAACAGGCCTGCTGAGGGGAGGGGAGCATGGACTTGGCCTCTGGAGTGGG
ACACTGGCCCTGGGAACCAGGCTGAGCTGAGTGGCCTCAAACCCCCCGTTGGATCAGACC
CTCCTGTGGGCAGGGTTCTTAGTGGATGAGTTACTGGGAAGAATCAGAGATAAAAACCAA
CCCAAATCAA

244/392

FIGURE 234

MFCPLKLILLPVLLDYSLGLNDLNVSPPELTVHVGDSALMGCVFQSTEDKCIFKIDWTLS
PGEHAKDEYVLYYYSNLSVPIGRFQNRVHLMGDILCNDGSLLLQDVQEADQGTYICEIRL
KGESQVFKKAVVLHVLPEEPKELMVHVGGLIQMGCVFQSTEVKHVTKVEWIFSGRRAKEE
IVFRYYHKL RMSVEYSQSWGHFQNRVNLVGDI FRNDGSIMLQGVRES DGGNYTCSIHLGN
LVFKKTI VLHVSPEEPRTLVT PAALRPLVLGGNQLVI IVGIVCATILLLPVLILIVKKTC
GNKSSVNSTVLVKNTKKTNPEIKEKPCHFERCEGEKHIYSPIIVREVIEEEEPSEKSEAT
YMTMHPVWPSLRSDRNNSLEKKSGGMPKTQQAF

245/392

FIGURE 235

TAAACAGCTACAATATTCCAGGGCCAGTCACTTGCCATTTCTCATAACAGCGTCAGAGA
GAAAGAACTGACTGAAACGTTTGAGATGAAGAAAGTTCTCCTCCTGATCACAGCCATCTT
GGCAGTGGCTGTTGGTTTCCAGTCTCTCAAGACCAGGAACGAGAAAAAGAAGTATCAG
TGACAGCGATGAATTAGCTTCAGGGTTTTTTGTGTTCCCTTACCCATATCCATTTGCCCC
ACTTCCACCAATTCCATTTCCAAGATTTCCATGGTTTAGACGTAATTTTCTATTCCAAT
ACCTGAATCTGCCCCTACAACCTCCCCTTCTAGCGAAAAGTAAACAAGAAGGATAAGTCA
CGATAAACCTGGTCACCTGAAATTGAAATTGAGCCACTTCCTTGAAGAATCAAAATTCCT
GTTAATAAAAGAAAAACAAATGTAATTGAAATAGCACACAGCATTTCTCTAGTCAATATCT
TTAGTGATCTTCTTTAATAAACATGAAAGCAAAGATTTTGGTTTCTTAATTTCCACA

246/392

FIGURE 236

MKKVLLLLITAILAVAVGFPVSQDQEREKRSISDSDELASGFFVFPYPYPFRPLPPIPFPR
FPWFRRNFPIPIPIESAPTTPLPSEK

Important features of the protein:

Signal peptide:

amino acids 1-17

Homologous region to B3-hordein:

amino acids 47-85

247/392

FIGURE 237

TCGCCATGGCCTCTGCCGGAATGCAGATCCTGGGAGTCGTCCTGACACTGCTGGGCTGGG
TGAATGGCCTGGTCTCCTGTGCCCTGCCCATGTGGAAGGTGACCGCTTTCATCGGCAACA
GCATCGTGGTGGCCCAGGTGGTGTGGGAGGGCCTGTGGATGTCTTGCGTGGTGCAGAGCA
CCGGCCAGATGCAGTGCAAGGTGTACGACTCACTGCTGGCGCTGCCACAGGACCTGCAGG
CTGCACGTGCCCTCTGTGTTCATCGCCCTCCTTGTGGCCCTGTTGCGCTTGCTGGTCTACC
TTGCTGGGGCCAAGTGTACCACCTGTGTGGAGGAGAAGGATTCCAAGGCCCGCCTGGTGC
TCACCTCTGGGATTGTCTTTGTTCATCTCAGGGGTCCTGACGCTAATCCCCGTGTGCTGGA
CGGCGCATGCCATCATCCGGGACTTCTATAACCCCCCTGGTGGCTGAGGCCCAAAAGCGGG
AGCTGGGGGCCTCCCTCTACTTGGGCTGGGCGGCCTCAGGCCTTTTGTGCTGGGTGGGG
GGTTGCTGTGCTGCACTTGCCCCCTCGGGGGGGTCCCAGGGCCCCAGCCATTACATGGCCC
GCTACTCAACATCTGCCCTGCCATCTCTCGGGGGCCCTCTGAGTACCCTACCAAGAATT
ACGTCTGACGTGGAGGGGAATGGGGGCTCCGCTGGCGCTAGAGCCATCCAGAAGTGGCAG
TGCCCAACAGCTTTGGGATGGGTTTCGTACCTTTTGTCTTGCTCCTGCTATTTTTCTTT
TGACTGAGGATATTTAAAAATTCATTTGAAAACTGAGCCAAGGTGTTGACTCAGACTCTCA
CTTAGGCTCTGCTGTTTTCTCACCTTGGATGATGGAGCCAAAGAGGGGATGCTTTGAGAT
TCTGGATCTTGACATGCCCATCTTAGAAGCCAGTCAAGCTATGGAACATAATGCGGAGGCT
GCTTGCTGTGCTGGCTTTGCAACAAGACAGACTGTCCCCAAGAGTTCCTGCTGCTGCTGG
GGGCTGGGCTTCCCTAGATGTCACTGGACAGCTGCCCCCATCCTACTCAGGTCTCTGGA
GCTCCTCTCTTCACCCCTGGAAAAACAAATCATCTGTAAACAAAGGACTGCCCACCTCCG
GAACTTCTGACCTCTGTTTCCTCCGTCTGATAAGACGTCCACCCCCCAGGGCCAGGTCC
CAGCTATGTAGACCCCCGCCCCACCTCCAACACTGCACCCTTCTGCCCTGCCCCCCTCG
TCTCACCCCTTTTACACTCACATTTTTTATCAAATAAAGCATGTTTTGTAGTGCA

248/392

FIGURE 238

MASAGMQILGVVLTLLGWVNGLVSCALPMWKVTAFIGNSIVVAQVWVWGLWMSCVVQSTG
QMCKVYDSLALPQDLQAARALCVIALLVALLFGLLVYLAGAKCTTCVEEKDSKARLVLT
SGIVFVISGVLTLIPVCWTAHAIRDFYNPLVAEAQKRELGASLYLGWAASGLLLLGGGL
LCCTCPSSGGSQGPPSHYMARYSTSAPAISRGPPSEYPTKNYV

Transmembrane domains:

amino acids 8-30 (type II), 82-102, 121-140, 166-186

249/392

FIGURE 239

AGTGACAATCTCAGAGCAGCTTCTACACCACAGCCATTTCCAGCATGAAGATCACTGGGG
GTCTCCTTCTGCTCTGTACAGTGGTCTATTTCTGTAGCAGCTCAGAAGCTGCTAGTCTGT
CTCCAAAAAAGTGGACTGCAGCATTTACAAGAAGTATCCAGTGGTGGCCATCCCCTGCC
CCATCACATACCTACCAGTTTGTGGTTCTGACTACATCACCTATGGGAATGAATGTCACT
TGTGTACCGAGAGCTTGAAAAGTAATGGAAGAGTTCAGTTTCTTCACGATGGAAGTTGCT
AAATTCTCCATGGACATAGAGAGAAAGGAATGATATTCTCATCATCATCTTCATCATCCC
AGGCTCTGACTGAGTTTCTTTCAGTTTACTGATGTTCTGGGTGGGGGACAGAGCCAGAT
TCAGAGTAATCTTGACTGAATGGAGAAAGTTTCTGTGCTACCCCTACAAACCCATGCCTC
ACTGACAGACCAGCATTTTTTTTTTAACACGTCAATAAAAAATAATCTCCCAGA

250/392

FIGURE 240

MKITGGLLLLCTVVYFCSSSEAASLSPKKVDCSIYKKYPVVAIPCPITYLPVCGSDYITY
GNECHLCTESLKSNGRVQFLHDGSC

Signal peptide:
amino acids 1-19

251/392

FIGURE 241

CCCGCGCCCGGTTCTCCCTCGCAGCACCTCGAAGTGCGCCCCCTCGCCCTCCTGCTCGCGC
CCCGCCGCCATGGCTGCCTCCCCGCGCGGCCTGCTGTCCTGGCCCTGACCGGGCTGGCG
CTGCTCCTGCTCCTGTGCTGGGGCCCAGGTGGCATAAGTGGAAATAAACTCAAGCTGATG
CTTCAAAAACGAGAAGCACCTGTTCCAACCTAAGACTAAAGTGGCCGTTGATGAGAATAAA
GCCAAAGAATTCTTTGGCAGCCTGAAGCGCCAGAAGCGGCAGCTGTGGGACCGGACTCGG
CCCGAGGTGCAGCAGTGGTACCAGCAGTTTCTCTACATGGGCTTTGATGAAGCGAAATTT
GAAGATGACATCACCTATTGGCTTAACAGAGATCGAAATGGACATGAATACTATGGCGAT
TACTACCAACGTCACTATGATGAAGACTCTGCAATTGGTCCCCGGAGCCCCTACGGCTTT
AGGCATGGAGCCAGCGTCAACTACGATGACTACTTAACCATGACTTGCCACACGCTGTACA
AGAAGCAAATAGCGATTCTCTTCATGTATCTCCTAATGCCTTACACTACTTGGTTTCTGA
TTTGCTCTATTTTCAGCAGATCTTTTCTACCTACTTTGTGTGATCAAAAAAGAAGAGTTAA
AACAAACACATGTAAATGCCTTTTGATATTTTCATGGGAATGCCTCTCATTTAAAAATAGAA
ATAAAGCATTTTGTAAAAAGA

252/392

FIGURE 242

MAASPARPAVLALTGLALLLLLCWGPGGISGNKLKMLQKREAPVPTKTKVAVDENKAKE
FLGSLKRQKRQLWDRTRPEVQQWYQQFLYMGFDEAKFEDDITYWLNDRDRNGHEYYGDYYQ
RHYEDSAIGPRSPYGFRHGASVNYDDY

Signal peptide:
amino acids 1-30

253/392

FIGURE 243

CTCCATTAAACCACCACCAGCTCCCCAAGCCACCCCTTCAGCCATGAAGTTCCTGCTCCT
GGTCTTGGCAGCCCTCGGATTCTTGACCCAGGTGATCCCAGCCAGTGCAGGTGGGTCAAA
ATGTGTGAGTAACACCCCGAGGATACTGCAGGACATGTTGCCACTGGGGGGAGACAGCATT
GTTTCATGTGCAACGCTTCCAGAAAATGCTGCATCAGCTACTCCTTCCTGCCGAAGCCTGA
CCTACCACAGCTCATCGGTAACCACTGGCAATCAAGGAGAAGAAACACACAAAGGAAAGA
CAAGAAGCAACAAACGACCGTAACATCATATAAACCCTGCTATCGCCTCCACCAACTCA
GAGAAATATCATTTCCACAGTTCCAATTCTCTTACATTGCTGAGTACTAGCCAAGGCTC
CTCTTTTATGGGGCAGATATCTATAGCCAACCCCAAACTTCTGTCTTCTATCATTCTGTC
ATTTCATCTAGTAACTAATTTGGAGTTTGTATCTATCTTACGAGAACAATCATCATGCAGA
TTCGTCCACAGGGGATCTGTGAGTTTGGGTCTCCAAATGAAAAATGTCAAGACAGAATT
GGACATGCAAAAGATTGACTGGGAGAACACACCTCTGATGGACAAAGGTGAGACAGAGCA
GCCACAGGCAGGGAGAGCCTTCAGACTGCAACGCTGGCCTGATACGTGTCAAAGGAGAGA
GGGATAGAGGAGGATTGAATAGAAGGAGACTAAGACTGCAGCTCTAAGAAAGTCTCAGCC
AAACAGATGGGGAGGCCCAAAGCAAGGCTTGCCCCCTCAGAGGAGCTCACGCAGGGCAGGA
ATAGCCAGGTTCTCATATCCCAGGGGTTTCAGACTTGGCTGAGAACAGCCCCCTGGAGAACA
TGGGGTGACTGCTACCATAGGTCTGGAAGTATGAGGCTGTCCACCAACTATCCCCTTGAA
GCAAGTTCTCTTGAAAGGAAATCTAAACAGTGCACCCCCATGGCTGCCACGGAGTATAAG
GAGGGAGAGAAAGGAGCTGAAAGTCTAGGTTTGGCCAGCTAGGTAGACTGACTTGTGAGG
TATTTATTTATTTCATTTGAGTAACAAAGCAGACAGAATACATAGCCACCATTGGTAGTAC
ACCCCAAAAGCAAGGATGGCATGATGCTGGTGACTCAAACGTGCCTACTCATGGTGTCAA
ATTGGCATAATCCTCTTGGAAGCTGTGTGGAATAAGCACAGAGAAGCAGAACTCTAAT
TGCTTAATCCACTAAACATTACTTCTGGAATTGGCTCATCATAAATTATCCAAGAGAAA
GCACAAAGTTATGGGCACAAAGGTTTTCCATATAATATTATTTAAAATGCTGAGAAAATG
AAAAAATCTAAATGGTGAAATATATACTAATGCCATCTATAAATACAAACAAATAGAATG
TTTATAGAATAATGGAACATAATAACATTATTCAAATTTGCATTTATGCTATAGTTGTCA
AAATTGTCTCCTTATATGATACAAAACCTCATGAAAATTATGACTTTTTTGTGTTGGTTGGA
AAGCAGAATTATGCATAAATTTCTCTTACAGTTCGATGCCATTAGTTTTATATAACAT
TTATTTGACACGTAAGTCTGACTTCTATCTGAGAAGAACAAACCAAAACACTCAGGCCTAAT
AATTAAAAACGGTCTTAAAACTAGCAAACCAGATAAGAAAAGATGTTAATGCCCATTC
CTAACTTATGTCTTAGACCAAATTAATTCTAGATGGTTTTTAAAATGACAGTGTAAAAGT
AAAGTATTAAAAGATTGTGTGGTCAAATATTCAATTTAAGAGCAAGGAAATCTTATAAA
TATAACAATAGAGGCAGAACTCATGTAAGAATAAATTGATTAGGTGGTATTAAATATTAA
GTTCTTATGTATGTCAAAGATATCATTTTGAATTCATCCATCTTATTGGGTATTGCAG
GAGTTCATTCTTTTTTGTGTTATAAATACTCTCCGTCATATGAATAGTATTCAATTTGTAT
ACTGGTTTGTGATGGACATTGGGTTGTTCCAGTTTATGGCTATTACAAATAAAGCTT
CTATGAACATTTATGTACA

254/392

FIGURE 244

MKFLLLVLAALGFLTQVIPASAGGSKCVSNTPGYCRTCCCHWGETALFMCNASRKCCISYS
FLPKPDLPLQLIGNHWQSRRRNTQRKDKKQQTTVTS

Important features of the protein:

Signal peptide:

amino acids 1-16

Transmembrane domain:

amino acids 1-22

N-glycosylation site:

amino acids 50-53

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 79-82

N-myristoylation site:

amino acids 23-28

255/392

FIGURE 245

GGAGAGAGGCGCGCGGGTGAAAGGCGCATTGATGCAGCCTGCGGCGGCCTCGGAGCGCGG
CGGAGCCAGACGCTGACCACGTTCTCTCCTCGGTCTCCTCCGCCTCCAGCTCCGCGCTG
CCCGGCAGCCGGGAGCCATGCGACCCAGGGCCCCGCCGCCTCCCCGCAGCGGCTCCGCG
GCCTCCTGCTGCTCCTGCTGCTGCAGCTGCCCGCGCCGTCGAGCGCCTCTGAGATCCCCA
AGGGGAAGCAAAAGGCGCAGCTCCGGCAGAGGGAGGTGGTGGACCTGTATAATGGAATGT
GCTTACAAGGGCCAGCAGGAGTGCTGGTTCGAGACGGGAGCCCTGGGGCCAATGTTATTC
CGGGTACACCTGGGATCCCAGGTCCGGATGGATTCAAAGGAGAAAAGGGGGAATGTCTGA
GGGAAAGCTTTGAGGAGTCCTGGACACCCAACTACAAGCAGTGTTTCATGGAGTTCATTGA
ATTATGGCATAGATCTTGGGAAAATTGCGGAGTGTACATTTACAAAGATGCGTTCAAATA
GTGCTCTAAGAGTTTTGTTTCAGTGGCTCACTTCGGCTAAAATGCAGAAATGCATGCTGTC
AGCGTTGGTATTTTCACATTCAATGGAGCTGAATGTTTCAGGACCTCTTCCCATTGAAGCTA
TAATTTATTTGGACCAAGGAAGCCCTGAAATGAATTCAACAATTAATATTCATCGCACTT
CTTCTGTGGAAGGACTTTGTGAAGGAATTGGTGTGATTAGTGGATGTTGCTATCTGGG
TTGGCACTTGTTTCAGATTACCCAAAAGGAGATGCTTCTACTGGATGGAATTCAGTTTCTC
GCATCATTATTGAAGAACTACCAAAATAAATGCTTTAATTTTCATTTGCTACCTCTTTTT
TTATTATGCCTTGGAATGGTTCACTTAAATGACATTTTAAATAAGTTTATGTATACATCT
GAATGAAAAGCAAAGCTAAATATGTTTACAGACCAAAGTGTGATTTACACTGTTTTTAA
ATCTAGCATTATTCATTTTGCTTCAATCAAAGTGGTTTCAATATTTTTTTTAGTTGGTT
AGAATACTTTCTTCATAGTCACATTCTCTCAACCTATAATTTGGAATATTGTTGTGGTCT
TTTGTTTTTTCTCTTAGTATAGCATTTTTTAAAAAATATAAAAGCTACCAATCTTGTAC
AATTTGTAAATGTTAAGAATTTTTTTTATATCTGTTAAATAAAAATTATTTCCAACA

256/392

FIGURE 246

MRPQGPAASPQRLRGLLLLLLLQLPAPSSASEIPKGKQKAQLRQREVVDLYNGMCLQGPA
GVPGRDGSPGANVIPGTPGIPGRDGFKEKGECLRESFEESWTPNYKQCSWSSLNYGIDL
GKIAECTFTKMRSNSALRVLFSGSLRLKCRNACCQRWYFTFNGAECSGPLPIEAIYLDQ
GSPEMNSTINIHRTSSVEGLCEGIGAGLVDVAIWVGTCSDYPKGDASTGWNSVSRIIIEE
LPK

Signal peptide:
amino acids 1-30

Transmembrane domain:
amino acids 195-217

257/392

FIGURE 247

GGCCGTTGGTTGGTGC GCGGCTGAAGGGTGTGGCGCGAGCAGCGTCGTTGGTTGGCCGGC
GGCGGGCCGGGACGGGCATGGCCCTGCTGCTGTGCCTGGTGTGCCTGACGGCGGCGCTGG
CCCACGGCTGTCTGCACTGCCACAGCAACTTCTCCAAGAAGTTCTCCTTCTACCGCCACC
ATGTGAACTTCAAGTCTTGGTGGGTGGGCGACATCCCCGTGTCAGGGGCGCTGCTCACC
ACTGGAGCGACGACACGATGAAGGAGCTGCACCTGGCCATCCCCGCCAAGATCACCCGGG
AGAAGCTGGACCAAGTGGCGACAGCAGTGTACCAGATGATGGATCAGCTGTACCAGGGGA
AGATGTACTTCCCCGGGTATTTCCCCAACGAGCTGCGAAACATCTTCCGGGAGCAGGTGC
ACCTCATCCAGAACGCCATCATCGAAAGGCACCTGGCACCAGGCAGCTGGGGAGGAGGGC
AGCTCTCCAGGGAGGGACCCAGCCTAGCACCTGAAGGATCAATGCCATCACCCCGCGGGG
ACCTCCCCTAAGTAGCCCCCAGAGGCGCTGGGAGTGTTGCCACCGCCCTCCCTGAAGTT
TGCTCCATCTCACGCTGGGGGTCAACCTGGGGACCCCTTCCCTCCGGGCCATGGACACAC
ATACATGAAAACCAGGCCGCATCGACTGTGACACCGCTGTGGCATCTTCCAGTACGAGA
CCATCTCCTGCAACAACTGCACAGACTCGCACGTGCGCTGCTTTGGCTATAACTGCGAGT
AGGGCTCAGGCATCACACCCACCCGTGCCAGGGCCCTACTGTCCCTGGGGTCCCAGGCTC
TCCTTGAGGGGGCTCCCCGCTTCCACCTGGCTGTCATCGGGTAGGGCGGGGCCGTGGG
TTCAGGGGCGCACCACTTCCAAGCCTGTGTCCACAGGTCTCTCGGCGCAGTGGAAGTCAG
CTGTCCAGGGCCTCCTGAACTACATAAATAACTGGCACAAAGTAAGTCCCCTCCTCAAACC
AACACAGGCAGTGTGTGTATGTGAGCACCTCGTGGGTGAGTATGTGTGGGGCACAGGCTG
GCTCCCTCAGCTCCCACGTCTTAGAGGGGCTCCCGAGGAGGTGGAACCTCAACCCAGCTC
TGCGCAGGAGGCGGCTGCAGTCCTTTTCTCCCTCAAAGGTCTCCGACCCCTCAGCTGGAGG
CGGGCATCTTTCTAAAGGGTCCCCATAGGGTCTGGTTCCACCCCATCCCAGGTCTGTGG
TCAGAGCCTGGGAGGGTTCCCTACGATGGTTAGGGGTGCCCCATGGAGGGGCTGACTGCC
CCACATTGCCCTTTCAGACAGGACACGAGCATGAGGTAAGGCCGCCCTGACCTGGACTTCA
GGGGGAGGGGGTAAAGGGAGAGAGGAGGGGGGCTAGGGGGTCTCTAGATCAGTGGGGC
ACTGCAGGTGGGGCTCTCCCTATACCTGGGACACCTGCTGGATGTACCTCTGCAACCAC
ACCCATGTGGTGGTTTCATGAACAGACCACGCTCCTCTGCCTTCTCCTGGCCTGGGACAC
ACAGAGCCACCCCGCCTTGTGAGTGACCCAGAGAAGGGAGGCCCTCGGGAGAAGGGGTGC
TCGTAAGCCAACACCAGCGTGCCGCGGCCCTGCACACCCTTCGGACATCCCAGGCACGAGG
GTGTGCTGGATGTGGCCACACATAGGACCACACGTCCCAGCTGGGAGGAGAGGCCTGGGG
CCCCCAGGGAGGGAGGCAGGGGGTGGGGGACATGGAGAGCTGAGGCAGCCTCGTCTCCCC
GCAGCCTGGTATCGCCAGCCTTAAGGTGTCTGGAGCCCCCACACTTGGCCAACCTGACCT
TGGAAGATGCTGCTGAGTGTCTCAAGCAGCACTGACAGCAGCTGGGCCTGCCCCAGGGCA
ACGTGGGGGCGGAGACTCAGCTGGACAGCCCCTGCCTGTCACTCTGGAGCTGGGCTGCTG
CTGCCTCAGGACCCCTCTCCGACCCCGGACAGAGCTGAGCTGGCCAGGGCCAGGAGGGC
GGGAGGGAGGGAATGGGGGTGGGCTGTGCGCAGCATCAGCGCCTGGGCAGGTCCGCAGAG
CTGCGGGATGTGATTAAAGTCCCTGATGTTTCTC

258/392

FIGURE 248

MALLLCVLCLTAALAHGCLHCHSNFSKKFSFYRHHVNFKSWWVGDI PVSGALLTDWSDDT
MKELHLAIPAKITREKLDQVATAVYQMMDQLYQGKMYFFPGYFPNELRNIFREQVHLIQNA
IIERHLAPGSWGGGQLSREGPSLAPEGSMPSPRGDL P

Signal peptide:
amino acids 1-15

259/392

FIGURE 249

CGACGATGCTACGCGCGCCCGGCTGCCTCCTCCGGACCTCCGTAGCGCCTGCCGCGGCC
TGGCTGCGGCGCTGCTCTCGTCGCTTGCGCGCTGCTCTCTTCTAGAGCCGAGGGACCCGG
TGGCCTCGTCGCTCAGCCCCCTATTTTCGGCACCAAGACTCGCTACGAGGATGTCAACCCCG
TGCTATTGTGCGGGCCCCGAGGCTCCGTGGCGGGACCCCTGAGCTGCTGGAGGGGACCTGCA
CCCCGGTGCAGCTGGTCGCCCTCATTTCGCCACGGCACCCGCTACCCACGGTCAAACAGA
TCCGCAAGCTGAGGCAGCTGCACGGGTTGCTGCAGGCCCGCGGGTCCAGGGATGGCGGGG
CTAGTAGTACCGGCAGCCGCGACCTGGGTGCAGCGCTGGCCGACTGGCCTTTGTGGTACG
CGGACTGGATGGACGGGCAGCTAGTAGAGAAGGGACGGCAGGATATGCGACAGCTGGCGC
TGCGTCTGGCCTCGCTCTTCCCGGCCCTTTTCAGCCGTGAGAACTACGGCCGCCTGCGGC
TCATCACCAGTTCCAAGCACCGCTGCATGGATAGCAGCGCCGCCTTCTGACGGGGCTGT
GGCAGCACTACCACCCTGGCTTGCCGCCGCCGGACGTGCGAGATATGGAGTTTGGACCTC
CAACAGTTAATGATAAACTAATGAGATTTTTTGTACTGTGAGAAGTTTTTAACTGAAG
TAGAAAAAATGCTACAGCTCTTTATCACGTGGAAGCCTTCAAACTGGACCAGAAATGC
AGAACATTTTAAAAAAGTTGCAGCTACTTTGCAAGTGCCAGTAAATGATTTAAATGCAG
ATTTAATTCAAGTAGCCTTTTTTACCTGTTTCAATTTGACCTGGCAATTAAAGGTGTAAAT
CTCCTTGGTGTGATGTTTTTGACATAGATGATGCAAAGGTATTAGAATATTTAAATGATC
TGAAACAATATTGGAAAAGAGGATATGGGTATACTATTAACAGTCGATCCAGCTGCACCT
TGTTTCAGGATATCTTTTCAGCACTTGGACAAAGCAGTTGAACAGAAACAAAGGTCTCAGC
CAATTTCTTCTCCAGTCATCCTCCAGTTTGGTTCATGCAGAGACTCTTCTTCCACTGCTTT
CTCTCATGGGCTACTTCAAAGACAAGGAACCCCTAACAGCGTACAATTACAAAAACAAA
TGCATCGGAAGTTCCGAAGTGGTCTCATTGTACCTTATGCCTCGAACCTGATATTTGTGC
TTTACCCTGTGAAAATGCTAAGACTCCTAAAGAACAATTCCGAGTGCAGATGTTATTAA
ATGAAAAGGTGTTACCTTTGGCTTACTCACAAGAACTGTTTCATTTTATGAAGATCTGA
AGAACCCTACAAGGACATCCTTCAGAGTTGTCAAACCAGTGAAGAATGTGAATTAGCAA
GGGCTAACAGTACATCTGATGAACATGAGTAAGTGAAGAACATTTTTAATTCTTTAGGA
ATCTGCAATGAGTGATTACATGCTTGTAATAGGTAGGCAATTCCTTGATTACAGGAAGCT
TTTATATTACTTGAGTATTTCTGTCTTTTACAGAAAAACATTGGGTTTCTCTCTGGGTT
TGGACATGAAATGTAAGAAAAGATTTTTTCACTGGAGCAGCTCTCTTAAGGAGAAACAAAT
CTATTTAGAGAAACAGCTGGCCCTGCAAATGTTTACAGAAATGAAATTCTTCTACTTAT
ATAAGAAATCTCACACTGAGATAGAATTGTGATTTTATAATAACACTTGAAAAGTGCTGG
AGTAACAAAATATCTCAGTTGGACCATCCTTAACCTTGATTGAACTGTCTAGGAACCTTAC
AGATTGTTCTGCAGTTCTCTCTTCTTTTCTCAGGTAGGACAGCTCTAGCATTTTCTTAA
TCAGGAATATTGTGGTAAGCTGGGAGTATCACTCTGGAAGAAAGTAACATCTCCAGATGA
GAATTTGAAACAAGAAACAGAGTGTGTAAGGACACCTTCACTGAAGCAAGTCGGAAA
GTACAATGAAAATAAATATTTTTGGTATTTATTTATGAAATATTTGAACATTTTTTCAAT
AATTCCTTTTTTACTTCTAGGAAGTCTCAAAGACCATCTTAAATTATTATATGTTTGGAC
AATTAGCAACAAGTCAGATAGTTAGAATCGAAGTTTTTCAAATCCATTGCTTAGCTAACT
TTTTCAATCTGTCACTTGGCTTCGATTTTTATATTTTCTATTATGAAATGTATCTTT
TGGTTGTTTGATTTTTCTTCTTTCTTTGTAAATAGTTCTGAGTTCTGTCAAATGCCGTG
AAAGTATTTGCTATAATAAAGAAAATTCTTGTGACTTTAAAAA

260/392

FIGURE 250

MLRAPGCLLRRTSVAPAAALAAALLSSLARCSLLEPRDPVASSLSPYFGTKTRYEDVNPVL
LSGPEAPWRDPELLEGTCTPVQLVALIRHGTRYPTVKQIRKLRQLHGLLQARGSRDGGAS
STGSRDLGAALADWPLWYADWMDGQLVEKGRQDMRQLALRLASLFPALFSRENYGRLRLI
TSSKHRCMDSSAAFLQGLWQHYHPGLPPPVDADMEFGPPTVNDKLMRFFDHCEKFLTEVE
KNATALYHVEAFKTGPENILKKVAATLQVPVNDLNADLIQVAFFTCSDLAIKGVKSP
WCDVFDIDDAKVLEYLNDLKQYWKRGYGYTTINSRSSCTLFQDIFQHLDKAVEQKQRSQPI
SSPVILQFGHAETLLPLLSLMGYFKDKEPLTAYNYKKQMRKFRSGLIVPYASNLI FVLY
HCENAKTPKEQFRVQMLLNEKVLPLAYSQETVSFYEDLKNHYKDILQSCQTSEECELARA
NSTSDEL

Important features:**Signal sequence**

amino acids 1-30

N-glycosylation sites:

amino acids 242-246, 481-485

N-myristoylation sites.

amino acids 107-113, 113-119, 117-123, 118-124, 128-134

Endoplasmic reticulum targeting sequence:

amino acids 484-489

261/392

FIGURE 251

GGAGAGCCGCGGCTGGGACCGGAGTGGGGAGCGCGGCGTGGAGGTGCCACCCGGCGCGGG
TGGCGGAGAGATCAGAAGCCTCTTCCCCAAGCCGAGCCAACCTCAGCGGGGACCCGGGGCT
CAGGGACGCGGCGGCGGCGGCGGCGGCGACTGCAGTGGCTGGACGATGGCAGCGTCCGCCGGA
GCCGGGGCGGTGATTGCAGCCCCAGACAGCCGGCGCTGGCTGTGGTTCGGTGTCTGGCGGCG
GCGCTTGGGCTCTTGACAGCTGGAGTATCAGCCTTGGAAGTATATACGCCAAAAGAAATC
TTCGTGGCAAATGGTACACAAGGGAAGCTGACCTGCAAGTTCAAGTCTACTAGTACGACT
GGCGGGTTGACCTCAGTCTCCTGGAGCTTCCAGCCAGAGGGGGCCGACACTACTGTGTCTG
TTTTTCCACTACTCCCAAGGGCAAGTGTACCTTGGGAATTATCCACCATTTAAAGACAGA
ATCAGCTGGGCTGGAGACCTTGACAAGAAAGATGCATCAATCAACATAGAAAATATGCAG
TTTATACACAATGGCACCTATATCTGTGATGTCAAAAACCTCCTGACATCGTTGTCCAG
CCTGGACACATTAGGCTCTATGTCTGTAGAAAAAGAGAATTTGCCTGTGTTTCCAGTTTGG
GTAGTGGTGGGCATAGTTACTGCTGTGGTCTTAGGTCTCACTCTGCTCATCAGCATGATT
CTGGCTGTCTCTATAGAAAGGAAAACTCTAAACGGGATTACACTGGCTGCAGTACATCA
GAGAGTTTGTCACCAGTTAAGCAGGCTCCTCGGAAGTCCCCCTCCGACACTGAGGGTCTT
GTAAAGAGTCTGCCTTCTGGATCTCACCAGGGCCCAGTCATATATGCACAGTTAGACCAC
TCCGGCGGACATCACAGTGACAAGATTAACAAGTCAGAGTCTGTGGTGTATGCGGATATC
CGAAAGAATTAAAGAGAATACCTAGAACATATCCTCAGCAAGAAACAAAACCAAACCTGGAC
TCTCGTGCAGAAAATGTAGCCCATTAACCATGTAGCCTTGGAGACCCAGGCAAGGACAA
GTACACGTGTACTCACAGAGGGAGAGAAAAGATGTGTACAAAGGATATGTATAAATATTCT
ATTTAGTCATCCTGATATGAGGAGCCAGTGTTCATGATGAAAAGATGGTATGATTCTAC
ATATGTACCCATTGTCTTGCTGTTTTTGTACTTTCTTTTCAGGTCAATTACAAATTGGGAG
ATTTCAGAAACATTTCCTTTCACCATCATTAGAAATGGTTTGCCTTAATGGAGACAATAG
CAGATCCTGTAGTATTTCCAGTAGACATGGCCTTTTAAATCTAAGGGCTTAAGACTGATTA
GTCTTAGCATTTACTGTAGTTGGAGGATGGAGATGCTATGATGGAAGCATAACCCAGGGTG
GCCTTTAGCACAGTATCAGTACCATTTATTTGTCTGCCGCTTTTAAAAAATACCCATTGG
CTATGCCACTTGAAAACAATTTGAGAAGTTTTTTTTGAAGTTTTTCTACTAAAATATGGG
GCAATTGTTAGCCTTACATGTTGTGTAGACTTACTTTAAGTTTTCACCCCTTGAAATGTGT
CATATCAATTTCTGGATTCAATAAGATTAGCAAAGGATAAATGCCGAAGGTCACT
TCATTCTGGACACAGTTGGATCAATACTGATTAAGTAGAAAATCCAAGCTTTGCTTGAGA
ACTTTTTGTAACGTGGAGAGTAAAAAGTATCGGTTTTTA

262/392

FIGURE 252

MAASAGAGAVIAAPDSRRWLWSVLAAALGLLTAGVSALEVYTPKEIFVANGTQGKLTCKF
KSTSTTGGLTSVSWSFQPEGADTTVSFFHYSQGQVYLGNYPPFKDRISWAGDLDDKKDASI
NIENMQFIHNGTYICDVKNPPDIVVQPGHIRLYVVEKENLPVFPVWVVVGIVTAVVLGLT
LLISMILAVLYRRKNSKRDTGCGSTSESLSPVKQAPRKSPSDTEGLVKSLPSGSHQGPVI
YAQLDHSGGHHSDKINKSESVVYADIRKN

Signal peptide:
amino acids 1-37

Transmembrane domain:
amino acids 161-183

263/392

FIGURE 253

GTGACACTATAGAAGAGCTATGACGTCGCATGCACGCGTACGTAAGCTCGGAATTCGGCT
CGAGGCTGGTGGGAAGAAGCCGAGATGGCGGCAGCCAGCGCTGGGGCAACCCGGCTGCTC
CTGCTCTTGCTGATGGCGGTAGCAGCGCCAGTCGAGCCCGGGGCAGCGGCTGCCGGGCC
GGGACTGGTGC GCGAGGGGCTGGGGCGGAAGGTCGAGAGGGCGAGGCCTGTGGCACGGTG
GGGCTGCTGCTGGAGCACTCATTTGAGATCGATGACAGTGCCAACTTCCGGAAGCGGGGC
TCACTGCTCTGGAACCAGCAGGATGGTACCTTGTCCCTGTCACAGCGGCAGCTCAGCGAG
GAGGAGCGGGGCCGACTCCGGGATGTGGCAGCCCTGAATGGCCTGTACCGGGTCCGGATC
CCAAGGCGACCCGGGGCCCTGGATGGCCTGGAAGCTGGTGGCTATGTCTCCTCCTTTGTC
CCTGCGTGCTCCCTGGTGGAGTCGCACCTGTTCGGACCAGCTGACCCTGCACGTGGATGTG
GCCGGCAACGTGGTGGGCGTGTCCGTGGTGACGCACCCCGGGGGCTGCCGGGGCCATGAG
GTGGAGGACGTGGACCTGGAGCTGTTCAACACCTCGGTGCAGCTGCAGCCGCCACCACA
GCCCCAGGCCCTGAGACGGCGGCCTTCATTGAGCGCCTGGAGATGGAAACAGGCCCAGAAG
GCCAAGAACCCCCAGGAGCAGAAGTCCTTCTTCGCCAAATACTGGATGTACATCATTCCC
GTCGTCCTGTTCCCTCATGATGTCAGGAGCGCCAGACACCGGGGGCCAGGGTGGGGGTGGG
GGTGGGGGTGGTGGTGGGGGTAGTGGCCTTTGCTGTGTGCCACCCTCCCTGTAAAGTCTAT
TTAAAAACATCGACGATACATTGAAATGTGTGAACGTTTTGAAAAGCTACAGCTTCCAGC
AGCCAAAAGCAACTGTTGTTTTGGCAAGACGGTCCTGATGTACAAGCTTGATTGAAATTC
ACTGCTCACTTGATACGTTATTCAGAAACCCAAGGAATGGCTGTCCCCATCCTCATGTGG
CTGTGTGGAGCTCAGCTGTGTTGTGTGGCAGTTTATTAACTGTCCCCCAGATCGACACG
CAAAAAAAAA

264/392

FIGURE 254

MAAASAGATRLLLLLLMAVAAPSRARGSGCRAGTGARGAGAEGREGEACGTVGLLLEHSF
EIDDSANFRKRGSLWNQDGTLSLSQRQLSEEERGLRDVAALNGLYRVRIIPRRPGALD
GLEAGGYVSSFVPACSLVESHLSDQLTLHVDVAGNVVGVSVVTHPGGCRGHEVEDVDLEL
FNTSVQLQPPTTAPGPETAAFIERLEMEQAQKAKNPQEQKSFFAKYWYIIPVVLFLMMS
GAPDTGGQGGGGGGGGGGGSGLCCVPPSL

Signal peptide:
amino acids 1-24

Transmembrane domain:
amino acids 226-243

265/392

FIGURE 255

GCGACGCGCGGCGGGGCGGCGAGAGGAAACGCGGCGCCGGGCGGGCCCGGCCCTGGAGA
TGGTCCCCGGCGCCGCGGGCTGGTGTGTCTCGTGCTCTGGCTCCCCGCGTGCGTCGCGG
CCCACGGCTTCCGTATCCATGATTATTTGTACTTTCAAGTGCTGAGTCCTGGGGACATTC
GATACATCTTCACAGCCACACCTGCCAAGGACTTTGGTGGTATCTTTCACACAAGGTATG
AGCAGATTCACCTTGTCCCCGCTGAACCTCCAGAGGCCTGCGGGGAACCTCAGCAACGGTT
TCTTCATCCAGGACCAGATTGCTCTGGTGGAGAGGGGGGGCTGCTCCTTCCTCTCCAAGA
CTCGGGTGGTCCAGGAGCACGGCGGGCGGGCGGTGATCATCTCTGACAACGCAGTTGACA
ATGACAGCTTCTACGTGGAGATGATCCAGGACAGTACCCAGCGCACAGCTGACATCCCCG
CCCTCTTCCTGCTCGGCCGAGACGGCTACATGATCCGCCGCTCTCTGGAACAGCATGGGC
TGCCATGGGCCATCATTTCCATCCCAGTCAATGTCACCAGCATCCCCACCTTTGAGCTGC
TGCAACCGCCCTGGACCTTCTGGTTAGAAAGAGTTTGTCCCACATTCCAGCCATAAGTGA
CTGAGCTGGGAAGGGGAAACCCAGGAATTTTGCTACTTGGAATTTGGAGATAGCATCTGG
GGACAAGTGGAGCCAGGTAGAGGAAAAGGGTTTGGGCGTTGCTAGGCTGAAAGGGAAGCC
ACACCACTGGCCTTCCCTTCCCCAGGGCCCCCAAGGGTGTCTCATGCTACAAGAAGAGGC
AAGAGACAGGCCCCAGGGCTTCTGGCTAGAACCCGAAACAAAAGGAGCTGAAGGCAGGTG
GCCTGAGAGCCATCTGTGACCTGTCACACTCACCTGGCTCCAGCCTCCCCTACCAGGGT
CTCTGCACAGTGACCTTCACAGCAGTTGTTGGAGTGGTTTAAAGAGCTGGTGTTTGGGGA
CTCAATAAACCTCACTGACTTTTTAGCAATAAAGCTTCTCATCAGGGTTGCAAAAAAAA
AAAAAAAAAAAAAAAAAAAA

266/392

FIGURE 256

MVPGAAGWCCLVLWLPACVAAHGFRIDYLYFQVLSPGDIRYIFTATPAKDFGGIFHTRY
EQIHLVPAEPPEACGELSNGFFIQDQIALVERGGCSFLSKTRVVQEHGGRAVIISDNAV
NDSFYVEMIQDSTQRTADIPALFLLGRDGYMIRRSLEQHGLPWAIISIPVNVTSIPTFEL
LQPPWTFW

Signal peptide:
amino acids 1-20

267/392

FIGURE 257

CTCGCTTCTTCCTTCTGGATGGGGGGCCAGGGGGCCAGGAGAGTATAAAGGCGATGTGG
AGGGTGCCCGGCACAACCAGACGCCCAGTCACAGGCGAGAGCCCTGGGATGCAACCGGCCA
GAGGCCATGCTGCTGCTGCTCACGCTTGCCCTCCTGGGGGGCCCCACCTGGGCAGGGAAG
ATGTATGGCCCTGGAGGAGGCAAGTATTTTACAGCACTGAAGACTACGACCATGAAATC
ACAGGGCTGCGGGTGTCTGTAGGTCTTCTCCTGGTGAAAAGTGTCCAGGTGAAACTTGGA
GACTCCTGGGACGTGAAACTGGGAGCCTTAGGTGGGAATACCCAGGAAGTCACCCTGCAG
CCAGGCGAATACATCACAAAAGTCTTTGTGCGCTTCCAAGCTTTCCTCCGGGGTATGGTC
ATGTACACCAGCAAGGACCGCTATTTCTATTTTGGGAAGCTTGATGGCCAGATCTCCTCT
GCCTACCCAGCCAAGAGGGGCAGGTGCTGGTGGGCATCTATGGCCAGTATCAACTCCTT
GGCATCAAGAGCATTGGCTTTGAATGGAATTATCCACTAGAGGAGCCGACCACTGAGCCA
CCAGTTAATCTCACATACTCAGCAAACCTACCCGTGGGTGCTAGGGTGGGGTATGGGGC
CATCCGAGCTGAGGCCATCTGTGTGGTGGTGGCTGATGGTACTGGAGTAACTGAGTCGGG
ACGCTGAATCTGAATCCACCAATAAATAAAGCTTCTGCAGAAA

268/392

FIGURE 258

MHRPEAML¹LL²LL³TL⁴ALL⁵GGPTWAGKMYGPGGGKYFSTTEDYDHEITGLRVSVGLLLVKS²⁰VQ
VKLGDSWDVKLGALGGNTQEVTLPGEYITKVFAFQAFLRGMVMYTSKDRYFYFGKLDG
QISSAYPSQEGQVLVGIYGQYQLLGIKSIGFEWNYPLEEPTTEPPVNL⁴⁰TYSANSPVGR

Signal peptide:
amino acids 1-22

269/392

FIGURE 259

CAGACATGGCTCAGTCACTGGCTCTGAGCCTCCTTATCCTGGTTCTGGCCTTTGGCATCC
CCAGGACCCAAGGCAGTGATGGAGGGGCTCAGGACTGTTGCCTCAAGTACAGCCAAAGGA
AGATTCCCGCCAAGGTTGTCCGCAGCTACCGGAAGCAGGAACCAAGCTTAGGCTGCTCCA
TCCCAGCTATCCTGTTCTTGCCCCGCAAGCGCTCTCAGGCAGAGCTATGTGCAGACCCAA
AGGAGCTCTGGGTGCAGCAGCTGATGCAGCATCTGGACAAGACACCATCCCCACAGAAAC
CAGCCCAGGGCTGCAGGAAGGACAGGGGGGCCTCCAAGACTGGCAAGAAAGGAAAGGGCT
CCAAAGGCTGCAAGAGGACTGAGCGGTCACAGACCCCTAAAGGGCCATAGCCCAGTGAGC
AGCCTGGAGCCCTGGAGACCCACACAGCCTCACCAGCGCTTGAAGCCTGAACCCAAGATG
CAAGAAGGAGGCTATGCTCAGGGGCCCTGGAGCAGCCACCCCATGCTGGCCTTGCCACAC
TCTTTCTCCTGCTTTAACCACCCCATCTGCATTCCCAGCTCTACCCTGCATGGCTGAGCT
GCCACAGCAGGCCAGGTCCAGAGAGACCGAGGAGGGAGAGTCTCCAGGGAGCATGAGA
GGAGGCAGCAGGACTGTCCCCTTGAAGGAGAATCATCAGGACCCTGGACCTGATACGGCT
CCCCAGTACACCCACCTCTTCCTTGTAATATGATTTATACCTAACTGAATAAAAAGCT
GTTCTGTCTTCCCNCCCA

270/392

FIGURE 260

MAQSLALSLLILVLAFGIPRTQGS D GGAQDCCLKYSQRKIPAKVVRSYRKQEPSLGCSIP
AILFLPRKRSQAELCADPKELWVQQLMQHLDKTPSPQKPAQGCRKDRGASKTGKKKGKGSK
GCKRTERSQTPKGP

Important features of the protein:

Signal peptide:

amino acids 1-17

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 67-71

N-myristoylation sites:

amino acids 17-23, 23-29, 27-33, 108-114, 118-124, 121-127

Amidation site:

amino acids 112-116

Small cytokines:

amino acids 51-91

271/392

FIGURE 261

GGGACTACAAGCCGCGCCGCGCTGCCGCTGGCCCCCTCAGCAACCCTCGACATGGCGCTGA
GGCGGCCACCGCGACTCCGGCTCTGCGCTCGGCTGCCTGACTTCTTCCTGCTGCTGCTTT
TCAGGGGCTGCCTGATAGGGGCTGTAAATCTCAAATCCAGCAATCGAACCCAGTGGTAC
AGGAATTTGAAAGTGTGGAAGTGTCTTGCATCATTACGGATTTCGAGACAAGTGACCCCA
GGATCGAGTGGAAGAAAATTCAAGATGAACAAACCACATATGTGTTTTTTTGACAACAAAA
TTCAGGGAGACTTGGCGGGTCGTGCAGAAATACTGGGGAAGACATCCCTGAAGATCTGGA
ATGTGACACGGAGAGACTCAGCCCTTTATCGCTGTGAGGTCGTTGCTCGAAATGACCGCA
AGGAAATTGATGAGATTGTGATCGAGTTAACTGTGCAAGTGAAGCCAGTGACCCCTGTCT
GTAGAGTGCCGAAGGCTGTACCAGTAGGCAAGATGGCAACACTGCACTGCCAGGAGAGTG
AGGGCCACCCCGGCCTCACTACAGCTGGTATCGCAATGATGTACCACTGCCACGGATT
CCAGAGCCAATCCCAGATTTTCGAATTTCTTCTTTCCACTTAAACTCTGAAACAGGCACTT
TGGTGTTCAGTGTCTGTTCAAGGACGACTCTGGGCAGTACTACTGCATTGCTTCCAATG
ACGCAGGCTCAGCCAGGTGTGAGGAGCAGGAGATGGAAGTCTATGACCTGAACATTGGCG
GAATTATTGGGGGGGTTCTGGTTGTCCTTGCTGTACTGGCCCTGATCACGTTGGGCATCT
GCTGTGCATACAGACGTGGCTACTTCATCAACAATAAACAGGATGGAGAAAGTTACAAGA
ACCCAGGGAAACCAGATGGAGTTAACTACATCCGCACTGACGAGGAGGGCGACTTCAGAC
ACAAGTCATCGTTTTGTGATCTGAGACCCGCGGTGTGGCTGAGAGCGCACAGAGCGCACGT
GCACATACCTCTGCTAGAAACTCCTGTCAAGGCAGCGAGAGCTGATGCACTCGGACAGAG
CTAGACACTCATTGAGAAAGCTTTTCGTTTTGGCCAAAGTTGACCACTACTCTTCTTACTC
TAACAAGCCACATGAATAGAAGAATTTTCCTCAAGATGGACCCGGTAAATATAACCACAA
GGAAGCGAAACTGGGTGCGTTCACTGAGTTGGGTTCCCTAATCTGTTTTCTGGCCTGATTCC
CGCATGAGTATTAGGGTGATCTTAAAGAGTTTGCTCACGTAAACGCCCGTGCTGGGCCCT
GTGAAGCCAGCATGTTCACTACTGGTTCGTTGAGCAGCCACGACAGCACCATGTGAGATGG
CGAGGTGGCTGGACAGCACCAGCAGCGCATCCCGGCGGGAACCCAGAAAAGGCTTCTTAC
ACAGCAGCCTTACTTCATCGGCCACAGACACCACCGCAGTTTTCTTCTTAAAGGCTCTGC
TGATCGGTGTTGCAGTGTCCATTGTGGAGAAGCTTTTTTGATCAGCATTGTTGTAATAACA
ACCAAAATCAGGAAGGTAAATGGTTGCTGGAAGAGGGATCTTGCCTGAGGAACCTTGCT
TGTCCAACAGGGTGTGAGGATTTAAGGAAAACCTTCGTCTTAGGCTAAGTCTGAAATGGT
ACTGAAATATGCTTTTCTATGGGTCTTGTTTATTTTATAAAATTTTACATCTAAATTTTT
GCTAAGGATGTATTTTGATTATTGAAAAGAAAATTTCTATTTAACTGTAAATATATTGT
CATACAATGTAAATAACCTATTTTTTTTAAAAAAGTTCAACTTAAGGTAGAAGTTCCAAG
CTACTAGTGTAAATGGAATAATCAATAATTAAGAGTATTTTACCCAAGGAATCCTCT
CATGGAAGTTTACTGTGATGTTCTTTTCTCACACAAGTTTTAGCCTTTTTCACAAGGGA
ACTCATACTGTCTACACATCAGACCATAGTTGCTTAGGAAACCTTTAAAAATTCAGTTA
AGCAATGTTGAAATCAGTTTGCATCTCTTCAAAAGAAACCTCTCAGGTTAGCTTTGAACT
GCCTCTTCTGAGATGACTAGGACAGTCTGTACCCAGAGGCCACCCAGAAGCCCTCAGAT
GTACATACACAGATGCCAGTCAGCTCCTGGGGTTGCGCCAGGCGCCCCCGCTCTAGCTCA
CTGTTGCCTCGCTGTCTGCCAGGAGGCCCTGCCATCCTTGGGCCCCTGGCAGTGGCTGTGT
CCCAGTGAGCTTTACTCACGTGGCCCTTGCTTCATCCAGCACAGCTCTCAGGTGGGCACT
GCAGGGACACTGGTGTCTTCCATGTAGCGTCCCAGCTTTGGGCTCCTGTAACAGACCTCT
TTTTGGTTATGGATGGCTCACAAAATAGGGCCCCCAATGCTATTTTTTTTTTTTAAAGTTT
GTTTAATTATTTGTTAAGATTGTCTAAGGCCAAAGGCAATTGCGAAATCAAGTCTGTCAA
GTACAATAACATTTTTTAAAAGAAAATGGATCCCACTGTTTCTTGGCCACAGAGAAAGC
ACCCAGACGCCACAGGCTCTGTGCGATTTCAAAACAAACCATGATGGAGTGGCGGCCAGT
CCAGCCTTTTAAAGAACGTGAGGTGGAGCAGCCAGGTGAAAGGCCTGGCGGGGAGGAAAG
TGAAACGCCTGAATCAAAAGCAGTTTTCTAATTTTGACTTTAAATTTTTTCATCCGCCGGA

272/392

GACACTGCTCCCATTTGTGGGGGGACATTAGCAACATCACTCAGAAGCCTGTGTTCTTCA
AGAGCAGGTGTTCTCAGCCTCACATGCCCTGCCGTGCTGGACTCAGGACTGAAGTGCTGT
AAAGCAAGGAGCTGCTGAGAAGGAGCACTCCACTGTGTGCCTGGAGAATGGCTCTCACTA
CTCACCTTGTCTTTCAGCTTCCAGTGTCTTGGGTTTTTTTATACTTTGACAGCTTTTTTTT
AATTGCATACATGAGACTGTGTTGACTTTTTTTTAGTTATGTGAAACACTTTGCCGCAGGC
CGCCTGGCAGAGGCAGGAAATGCTCCAGCAGTGGCTCAGTGCTCCCTGGTGTCTGCTGCA
TGGCATCCTGGATGCTTAGCATGCAAGTTCCCTCCATCATTGCCACCTTGGTAGAGAGGG
ATGGCTCCCCACCTCAGCGTTGGGGATTACAGCTCCAGCCTCCTTCTTGGTTGTCATAG
TGATAGGGTAGCCTTATTGCCCCCTCTTCTTATACCCTAAAACCTTCTACACTAGTGCCA
TGGGAACCAGGTCTGAAAAAGTAGAGAGAAGTGAAAGTAGAGTCTGGGAAGTAGCTGCCT
ATAACTGAGACTAGACGGAAAAGGAATACTCGTGTATTTTAAGATATGAATGTGACTCAA
GACTCGAGGCCGATACGAGGCTGTGATTCTGCCTTTGGATGGATGTTGCTGTACACAGAT
GCTACAGACTTGTACTAACACACCGTAATTTGGCATTGTGTTAACCTCATTATATAAAGC
TTCAAAAAAACCCA

273/392

FIGURE 262

MALRRPPRLRLCARLPDFLLLLFRGCLIGAVNLKSSNRTPVVQEFESVELSCIITDSQT
SDPRIEWKKIQDEQTTYVFFDNKIQGDLGRAEILGKTS�KIWNVTRRDSALYRCEVVAR
·NDRKEIDEIVIELTVQVKPVTVPVCRVPKAVPVGKMATLHCQESEGHPRPHYSWYRNDVPL
PTDSRANPRFRNSSFHLNSETGTLVFTAVHKDDSGQYYCIASNDAGSARCEEQEMEVDL
NIGGIIGGVLVVLAVLALITLGICCAYYRRGYFINNKQDGESYKNPGKPDGVNYIRTDEEG
DFRHKSSFVI

Important features of the protein:**Signal peptide:**

amino acids 1-30

Transmembrane domain:

amino acids 243-263

N-glycosylation sites:

amino acids 104-107, 192-195

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 107-110

Casein kinase II phosphorylation site:

amino acids 106-109, 296-299

Tyrosine kinase phosphorylation site:

amino acids 69-77

N-myristoylation sites:

amino acids 26-31, 215-220, 226-231, 243-248, 244-249, 262-267

274/392

FIGURE 263

CCAGGACCAGGGCGCACCGGCTCAGCCTCTCACTTGTGAGAGGCCGGGGAAGAGAAGCAA
AGCGCAACGGTGTGGTCCAAGCCGGGGCTTCTGCTTCGCCTCTAGGACATACACGGGACC
CCCTAACTTCAGTCCCCCAAACGCGCACCCCTCGAAGTCTTGAAGTCCAGCCCCGCACATC
CACGCGCGGCACAGGCGCGGCAGGCGGCAGGTCCCGGCCGAAGGCGATGCGCGCAGGGGG
TCGGGCAGCTGGGCTCGGGCGGGCGGGAGTAGGGCCCCGGCAGGAGGCAGGAGGCTGCAT
ATTCAGAGTCGCGGGCTGCGCCCTGGGCAGAGGCCGCCCTCGCTCCACGCAACACCTGCT
GCTGCCACCGCGCCGCGATGAGCCGCGTGGTCTCGCTGCTGCTGGGCGCCGCGCTGCTCT
GCGGCCACGGAGCCTTCTGCCGCCGCGTGGTCAAGCGCCAAAAGGTGTGTTTTGCTGACT
TCAAGCATCCCTGCTACAAAATGGCCTACTTCCATGAACTGTCCAGCCGAGTGAGCTTTC
AGGAGGCACGCTGGCTTGTGAGAGTGAGGGAGGAGTCCTCCTCAGCCTTGAGAAATGAAG
CAGAACAGAAGTTAATAGAGAGCATGTTGCAAAACCTGACAAAACCCGGGACAGGGATTT
CTGATGGTGATTTCTGGATAGGGCTTTGGAGGAATGGAGATGGGCAAACATCTGGTGCCT
GCCCAGATCTCTACCAGTGGTCTGATGGAAGCAATTCCCAGTACCGAAACTGGTACACAG
ATGAACCTTCCCTGCGGAAGTGAAAAGTGTGTTGTGATGTATCACCACCAACTGCCAATC
CTGGCCTTGGGGGTCCCTACCTTTACCAGTGGAAATGATGACAGGTGTAACATGAAGCACA
ATTATATTTGCAAGTATGAACCAGAGATTAATCCAACAGCCCCCTGTAGAAAAGCCTTATC
TTACAAATCAACCAGGAGACACCCATCAGAATGTGGTTGTTACTGAAGCAGGTATAATTC
CCAATCTAATTTATGTTGTTATACCAACAATACCCCTGCTCTTACTGATACTGGTTGCTT
TTGGAACCTGTTGTTTCCAGATGCTGCATAAAAGTAAAGGAAGAACAAAACTAGTCCAA
ACCAGTCTACACTGTGGATTTCAAAGAGTACCAGAAAAGAAAGTGGCATGGAAGTATAAAT
AACTCATTGACTTGGTTCCAGAATTTTGTAAATCTGGATCTGTATAAGGAATGGCATCAG
ACAATAGCTTGGAAATGGCTTGAAATCACAAAGGATCTGCAAGATGAACTGTAAGCTCCC
CCTTGAGGCAAATATTAAGTAATTTTATATGTCTATTATTTCATTTAAAGAATATGCT
GTGCTAATAATGGAGTGAGACATGCTTATTTTGCTAAAGGATGCACCCAACTTCAAAC
TCAAGCAAATGAAATGGACAATGCAGATAAAGTTGTTATCAACACGTGGGAGTATGTGT
GTTAGAAGCAATTCCTTTTATTTCTTTCACCTTTCATAAGTTGTTATCTAGTCAATGTAA
TGATATTTGTATTGAAATTTACAGTGTGCAAAAGTATTTTACCTTTGCATAAGTGTTTGA
TAAAAATGAACTGTTCTAATATTTATTTTTATGGCATCTCATTTTTCAATACATGCTCTT
TTGATTAAAGAACTTATTACTGTTGTCAACTGAATTCACACACACACAAATATAGTACC
ATAGAAAAAGTTTGTCTTCGAAATAATTCATCTTTCAGCTTCTCTGCTTTTGGTCAAT
GTCTAGGAAATCTCTTCAGAAATAAGAAGCTATTTCAATTAAGTGTGATATAAACCTCCTC
AAACATTTTACTTAGAGGCAAGGATTGTCTAATTTCAATTGTGCAAGACATGTGCCTTAT
AATTATTTTACTTAAATTAACAGATTTTGTAAATAATGTAACCTTTGTTAATAGGTGC
ATAAACACTAATGCAGTCAATTTGAACAAAAGAAGTGACATACACAATATAAATCATATG
TCTTCACACGTTGCCTATATAATGAGAAGCAGCTCTCTGAGGGTTCTGAAATCAATGTGG
TCCCTCTCTTGCCCACTAAACAAAGATGGTTGTTTCGGGGTTTGGGATTGACACTGGAGGC
AGATAGTTGCAAAGTTAGTCTAAGGTTTCCCTAGCTGTATTTAGCCTCTGACTATATTAG
TATACAAAGAGGTCATGTGGTTGAGACCAGGTGAATAGTCACTATCAGTGTGGAGACAAG
CACAGCACACAGACATTTTAGGAAGGAAAGGAACACGAAATCGTGTGAAAATGGGTGG
AACCCATCAGTGATCGCATATTCATTGATGAGGGTTTGGCTTGAGATAGAAAATGGTGGCT
CCTTCTGTCTTATCTCCTAGTTTCTTCAATGCTTACGCCTTGTTCTTCTCAAGAGAAAG
TTGTAACCTCTCTGGTCTTCATATGTCCCTGTGCTCCTTTTAAACCAATAAAGAGTTCTTG
TTTCTGGGGGAAA

275/392

FIGURE 264

MSRVVSLLLGAALLCGHGAFRRVVSQGQKVCFAFKHPCYKMAFHELSRVSFQEARLA
CESEGGVLLSLENEAEQKLIESMLQNLTKPGTGISDGDFWIGLWRNGDGQTSACPDLYQ
WSDGSNSQYRNWYTDEPSCGSEKCVVMYHQPTANPGLGGPYLYQWNDDRCNMKHNKY
EPEINPTAPVEKPYLTNQPQDTHQNVVTEAGIIPNLIYVVIPTIPLLLILVAFGTCCF
QMLHKSKGRKTSPNQSTLWISKSTRKESGMEV

Important features of the protein:**Signal peptide:**

amino acids 1-21

Transmembrane domain:

amino acids 214-235

N-glycosylation sites:

amino acids 86-89 and 255-258

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 266-269

N-myristoylation sites:

amino acids 27-32, 66-71, 91-96, 93-98, 102-107, 109-114, 140-145 and 212-217

276/392

FIGURE 265

GGAGAATGGAGAGAGCAGTGAGAGTGGAGTCCGGGGTCCTGGTCGGGGTGGTCTGTCTGC
TCCTGGCATGCCCTGCCACAGCCACTGGGCCCCGAAGTTGCTCAGCCTGAAGTAGACACCA
CCCTGGGTCGTGTGCGAGGCCGGCAGGTGGGCGTGAAGGGCACAGACCGCCTTGTGAATG
TCTTTCTGGGCATTCCATTGCCCAGCCGCCACTGGGCCCTGACCGGTTCTCAGCCCCAC
ACCCAGCACAGCCCTGGGAGGGTGTGCGGGATGCCAGCACTGCGCCCCCAATGTGCCTAC
AAGACGTGGAGAGCATGAACAGCAGCAGATTTGTCCTCAACGGAAAACAGCAGATCTTCT
CCGTTTCAGAGGACTGCCTGGTCCTCAACGTCTATAGCCCAGCTGAGGTCCCCGCAGGGT
CCGGTAGGCCGGTCATGGTATGGGTCCATGGAGGCGCTCTGATAACTGGCGCTGCCACCT
CCTACGATGGATCAGCTCTGGCTGCCTATGGGGATGTGGTTCGTGGTTACAGTCCAGTACC
GCCTTGGGGTCCTTGGCTTCTTCAGCACTGGAGATGAGCATGCACCTGGCAACCAGGGCT
TCCTAGATGTGGTAGCTGCTTTGCGCTGGGTGCAAGAAAACATCGCCCCCTTCGGGGGTG
ACCTCAACTGTGTCACTGTCTTTGGTGGATCTGCCGGTGGGAGCATCATCTCTGGCCTGG
TCCTGTCCCCAGTGGCTGCAGGGCTGTTCCACAGAGCCATCACACAGAGTGGGGTCATCA
CCACCCCAGGGATCATCGACTCTCACCTTGGCCCCTAGCTCAGAAAATCGAAACACCT
TGGCCTGCAGCTCCAGCTCCCCGGCTGAGATGGTGCAGTGCCTTCAGCAGAAAGAAGGAG
AAGAGCTGGTCCTTAGCAAGAAGCTGAAAAATACTATCTATCCTCTCACCGTTGATGGCA
CTGTCTTCCCCAAAAGCCCCAAGGAACTCCTGAAGGAGAAGCCCTTCCACTCTGTGCCCT
TCCTCATGGGTGTCAACAACCATGAGTTTCTGAGTGCCTCATCCCCAGGGGCTGGGGTCTCC
TGGATACAATGGAGCAGATGAGCCGGGAGGACATGCTGGCCATCTCAACACCCGTCTTGA
CCAGTCTGGATGTGCCCCCTGAGATGATGCCACCGTCATAGATGAATACCTAGGAAGCA
ACTCGGACGCACAAGCCAAATGCCAGGCGTTCCAGGAATTCATGGGTGACGTATTTCATCA
ATGTTCCACCGTCAGTTTTTCAAGATACCTTCGAGATTCTGGAAGCCCTGTCTTTTTCT
ATGAGTTCCAGCATCGACCCAGTTCTTTTGCAGAGATCAAACCTGCCTGGGTGAAGGCTG
ATCATGGGGCCGAGGGTGCTTTTTGTGTTCCGAGGTCCCTTCCTCATGGACGAGAGCTCCC
GCCTGGCCTTTCCAGAGGCCACAGAGGAGGAGAAGCAGCTAAGCCTCACCATGATGGCCC
AGTGGACCCACTTTGCCCCGACAGGGGACCCCAATAGCAAGGCTCTGCCTCCTTGGCCCC
AATTCAACCAGGCGGAACAATATCTGGAGATCAACCCAGTGCCACGGGCCGGACAGAAGT
TCAGGGAGGCCTGGATGCAGTTCTGGTCAGAGACGCTCCCCAGCAAGATACAACAGTGGC
ACCAGAAGCAGAAGAACAGGAAGGCCAGGAGGACCTCTGAGAGGCCAGGCCTGAACCTTCT
TGGCTGGGGCAAACCACTCTTCAAGTGGTGGCAGAGTCCCAGCACGGCAGCCCGCCTCTC
CCCCTGCTGAGACTTTAATCTCCACCAGCCCTTAAAGTGTGCGCCGCTCTGTGACTGGAG
TTATGCTCTTTTGAATGTCAAGGCCGCTCCACCTCTGGGGCATTGTACAAGTTCT
TCCCTCTCCCTGAAGTGCTTTTCTGCTTTCTTCGTGGTAGGTTCTAGCACATTCTCTA
GCTTCCTGGAGGACTCACTCCCCAGGAAGCCTTCCCTGCCTTCTCTGGGCTGTGCGGCCC
CGAGTCTGCGTCCATTAGAGCACAGTCCACCCGAGGCTAGCACCGTGTCTGTGTCTGTCT
CCCCCTCAGAGGAGCTCTCTCAAAATGGGGATTAGCCTAACCCCACTCTGTACCCACAC
CAGGATCGGGTGGGACCTGGAGCTAGGGGGTGTGTTGCTGAGTGAGTGAGTGAAACACAGA
ATATGGGAATGGCAGCTGCTGAACCTGAACCCAGAGCCTTCAGGTGCCAAAGCCATACTC
AGGCCCCCACCAGCATTTGTCCACCCTGGCCAGAAGGGTGCATGCCAATGGCAGAGACCTG
GGATGGGAGAAGTCTTGGGGCGCCAGGGGATCCAGCCTAGAGCAGACCTTAGCCCCCTGAC
TAAGGCCTCAGACTAGGGCGGGAGGGGTCTCCTCCTCTCTGCTGCCAGTCTTGGCCCCCT
GCACAAGACAACAGAATCCATCAGGGCCATGAGTGTACCCAGACCTGACCTCACCAAT
TCCAGCCCCCTGACCTCAGGACGCTGGATGCCAGCTCCCAGCCCCAGTGCCGGGTCTCTCC
CTCCCTTCTGCTTGGGGAGACCAGTTTCTGGGGAGCTTCCAAGAGCACCCACCAAGAC
ACAGCAGGACAGGCCAGGGGAGGGCATCTGGACCAGGGCATCCGTGGGGCTATTGTACA
GAGAAAAGAAGAGACCCACCACTCGGGCTGCAAAAGGTGAAAAGCACCAAGAGGTTTTT

277/392

AGATGGAAGTGAGAGGTGACAGTGTGCTGGCAGCCCTCACAGCCCTCGCTTGCTCTCCCT
GCCGCCTCTGCCTGGGCTCCCACTTTGGCAGCACTTGAGGAGCCCTTCAACCCGCCGCTG
CACTGTAGGAGCCCCCTTTCTGGGCTGGCCAAGGCCGGAGCCAGCTCCCTCAGCTTGCGGG
GAGGTGCGGAGGGAGAGGGGCGGGCAGGAACCGGGGCTGCGCGCAGCGCTTGCGGGCCAG
AGTGAGTTCGGGTGGGCGTGGGCTCGGCGGGGCCCCACTCAGAGCAGCTGGCCGGCCCC
AGGCAGTGAGGGCCTTAGCACCTGGGCCAGCAGCTGCTGTGCTCGATTTCTCGCTGGGCC
TTAGCTGCCTCCCCGCGGGGCAGGGCTCGGGACCTGCAGCCCTCCATGCCTGACCCTCCC
CCCACCCCCCGTGGGCTCCTGTGCGGCCGGAGCCTCCCCAAGGAGCGCCGCCCTTGCTC
CACAGCGCCCAGTCCCATCGACCACCCAAGGGCTGAGGAGTGCGGGTGACAGCGCGGGA
CTGGCAGGCAGCTCCACCTGCTGCCCCAGTGCTGGATCCACTGGGTGAAGCCAGCTGGGC
TCCTGAGTCTGGTGGGGACTTGGAGAACCCTTTATGTCTAGCTAAGGGATTGTAAATACAC
CGATGGGCACTCTGTATCTAGCTCAAGGTTTGTAACACACCAATCAGCACCTGTGTCT
AGCTCAGTGTTTGTGAATGCACCAATCCACACTCTGTATCTGGCTACTCTGGTGGGGACT
TGGAGAACCTTTGTGTCCACACTCTGTATCTAGCTAATCTAGTGGGGATGTGGAGAACCT
TTGTGTCTAGCTCAGGGATCGTAAACGCACCAATCAGCACCTGTCAAAACAGACCACTT
GACTCTCTGTAAAATGGACCAATCAGCAGGATGTGGGTGGGGCGAGACAAGAGAATAAAA
GCAGGCTGCCTGAGCCAGCAGTGACAACCCCCCTCGGGTCCCCTCCACGCCGTGGAAGC
TTTGTTCCTTCGCTCTTTGCAATAAATCTTGCTACTGCCCAAAA

278/392

FIGURE 266

MERAVRVESGVLVGVVCLLLACPATATGPEVAQPEVDTTLGRVRGRQVGVKGTDRLVNVF
LGIPFAQPPLGPDRFSAPHPAQPWEGVRDASTAPPMCLQDVESMNSSRFVLNGKQQIFSV
SEDCLVLNVYSPAIEVPAGSGRPVMVWVHGGALITGAATSYDGSALAAYGDVVVVTVQYRL
GVLGFFSTGDEHAPGNQGFLDVVAALRWVQENIAPFGDLNLCVTVFGGSAGGSIISGLVL
SPVAAGLFHRAITQSGVITTPGIIDSHPWPLAQKIANLACSSSSPAEMVQCLOQKEGEE
LVLSSKKLKNTIYPLTVDGTVPKSPKELLKEKPFHSVPFLMGVNNHEFSWLI PRGWGLLD
TMEQMSREDMLAISTPVLTSLDVPPPEMMPTVIDEYLGNSDAQAKCQAFQEFMGDVFINV
PTVSFSRYLRDSGSPVFFYEFQHRPSSF AKIKPAWVKADHGAEGAFVFGGPFLMDESSL
AFPEATEEEKQLSLTMMAQWTHFARTGDPNSKALPPWPQFNQAEQYLEINPVPRAGQKFR
EAWMQFWSETLPSKIQQWHQKQKNRKAQEDL

Important features of the protein:**Signal peptide:**

amino acids 1-27

Transmembrane domain:

amino acids 226-245

N-glycosylation site:

amino acids 105-109

N-myristoylation sites:amino acids 10-16, 49-55, 62-68, 86-92, 150-156, 155-161,
162-168, 217-223, 227-233, 228-234, 232-238, 262-268, 357-363,
461-467**Prokaryotic membrane lipoprotein lipid attachment site:**

amino acids 12-23

Carboxylesterases type-B serine active site:

amino acids 216-232

279/392

FIGURE 267

TGTCGCCTGGCCCTCGCCATGCAGACCCCGCGAGCGTCCCCTCCCCGCCCCGGCCCTCCTG
CTTCTGCTGCTGCTACTGGGGGGCGCCACGGCCTCTTTCCTGAGGAGCCGCCGCCGCTT
AGCGTGGCCCCCAGGGACTACCTGAACCACTATCCCGTGTTTGTGGGCAGCGGGCCCCGGA
CGCCTGACCCCCCGCAGAAGGTGCTGACGACCTCAACATCCAGCGAGTCCTGCGGGTCAAC
AGGACGCTGTTCAATTGGGGACAGGGACAACCTCTACCGCGTAGAGCTGGAGCCCCCCACG
TCCACGGAGCTGCGGTACCAGAGGAAGCTGACCTGGAGATCTAACCCCAGCGACATAAAC
GTGTGTTCGGATGAAGGGCAAACAGGAGGGCGAGTGTCGAAACTTCGTAAAGGTGCTGCTC
CTTCGGGACGAGTCCACGCTCTTTGTGTGCGGTTCCAACGCCTTCAACCCGGTGTCGCC
AACTACAGCATAGACACCCTGCAGCCCGTCGGAGACAACATCAGCGGTATGGCCCCGCTGC
CCGTACGACCCCAAGCACGCCAATGTTGCCCTCTTCTCTGACGGGATGCTCTTACAGCT
ACTGTTACCGACTTCTAGCCATTGATGCTGTCTATCTACCGCAGCCTCGGGGACAGGCCC
ACCTTGCGCACCGTGAAACATGACTCCAAGTGGTTCAAAGAGCCTTACTTTGTCCATGCG
GTGGAGTGGGGCAGCCATGTCTACTTCTTCTTCCGGGAGATTGCGATGGAGTTTAACTAC
CTGGAGAAGGTGGTGGTGTCCCGCGTGGCCCCGAGTGTCGCAAGAACGACGTGGGAGGCTCC
CCCCGCGTGCTGGAGAAGCAGTGGACGTCCTTCTGAAGGCGCGGCTCAACTGCTCTGTA
CCCCGAGACTCCCATTTCTACTTCAACGTGCTGCAGGCTGTACGGGCGTGCTCAGCCTC
GGGGGCGGGCCCGTGCTGCTGCGCCTTTTTCACGCCCAGCAACAGCATCCCTGGCTCG
GCTGTCTGCGCCTTTGACCTGACACAGGTGGCAGCTGTGTTTGAAGGCCGCTTCCGAGAG
CAGAAGTCCCCCGAGTCCATCTGGACGCCGGTGCCGGAGGATCAGGTGCCTCGACCCCGG
CCCCGGTGCTGCGCAGCCCCCGGGATGCAGTACAATGCCTCCAGCGCCTTGCCGGATGAC
ATCCTCAACTTTGTCAAGACCCACCTCTGATGGACGAGGCGGTGCCCTCGCTGGGCCAT
GCGCCCTGGATCCTGCGGACCCTGATGAGGCACCAGCTGACTCGAGTGGCTGTGGACGTG
GGAGCCGGCCCCCTGGGGCAACCAGACCGTTGTCTTCTGGGTTCTGAGGCGGGGACGGTC
CTCAAGTTCTCTGTCGGGCCCAATGCCAGCACCTCAGGGACGTCTGGGCTCAGTGTCTTC
CTGGAGGAGTTTGAGACCTACCGGCCGGACAGGTGTGGACGGCCCCGGCGGTGGCGAGACA
GGGCAGCGGCTGCTGAGCTTGAGCTGGACGCAGCTTCCGGGGGGCCTGCTGGCTGCCTTC
CCCCGCTGCGTGCTCCGAGTGCCTGTGGCTCGCTGCCAGCAGTACTCGGGGTGTATGAAG
AACTGTATCGGCAGTCAGGACCCCTACTGCGGGTGGGCCCCCGACGGCTCCTGCATCTTC
CTCAGCCCCGGGCACCAGAGCCGCCTTTGAGCAGGACGTGTCCGGGGCCAGCACCTCAGGC
TTAGGGGACTGCACAGGACTCCTGCGGGCCAGCCTCTCCGAGGACCGCGCGGGGCTGGTG
TCGGTGAACCTGCTGGTAACGTCGTCGGTGGCGGCCCTTCGTGGTGGGAGCCGTGGTGTCC
GGCTTCAGCGTGGGCTGGTTTCGTGGGCCTCCGTGAGCGGCGGGAGCTGGCCCCGGCGCAAG
GACAAGGAGGCCATCCTGGCGCACGGGGCGGGCGAGGCGGTGCTGAGCGTCAGCCGCCTG
GGCGAGCGCAGGGCGCAGGGTCCCGGGGGCGGGGCGGAGGCGGTGGCGGTGGCGCCGGG
GTTCCCCCGGAGGCCCTGCTGGCGCCCCCTGATGCAGAACGGCTGGGCCAAGGCCACGCTG
CTGCAGGGCGGGCCCCACGACCTGGACTCGGGGCTGCTGCCCACGCCCGAGCAGACGCCG
CTGCCGCGAGAAGCGCCTGCCACTCCGCACCCGCACCCCCACGCCCTGGGCCCCCGCGCC
TGGGACCACGGCCACCCCTGCTCCCGGCCTCCGCTTATCCTCCCTCCTGCTGCTGGCG
CCCCGCCGGGCCCCCGAGCAGCCCCCGCGCCTGGGGAGCCGACCCCCGACGGCCGCCTC
TATGCTGCCCCGGCCCGGCCGCGCCTCCACGGCGACTTCCCGCTACCCCCACGCCAGC
CCGGACCGCCGGCGGGTGCTGTCGCGCCACGGGCCCCCTTGACCCAGCCTCAGCCGCC
GATGGCCTCCCGCGGCCCTGGAGCCCGCCCCGACGGGCAGCCTGAGGAGGCCACTGGGC
CCCCACGCCCTCCGGCCGCCACCTGCGCCGCACCCACACGTTCAACAGCGGCGAGGCC
CGGCCTGGGGACCGCCACCGCGGCTGCCACGCCCCGGCGGGCACAGACTTGGCCACCTC
CTCCCCATATGGGGGGCGGACAGGACTGCGCCCCCGTGCCCTAGGCCGGGGCCCCCG
ATGCCTTGGCAGTGCCAGCCACGGGAACCAGGAGCGAGAGACGGTGCCAGAACGCCGGGG
CCCCGGGCAACTCCGAGTGGGTGCTCAAGTCCCCCCCCGCGACCCACCCGCGGAGTGGGG

280/392

GCCCCCTCCGCCACAAGGAAGCACAACCAGCTCGCCCTCCCCCTACCCGGGGCCGCAGGA
CGCTGAGACGGTTTTGGGGGTGGGTGGGCGGGAGGACTTTGCTATGGATTTGAGGTTGACC
TTATGCGCGTAGGTTTTGGTTTTTTTTTGCAGTTTTGGTTTCTTTTGC GGTTTTCTAACC
AATTGCACAACCTCCGTTCTCGGGGTGGCGGCAGGCAGGGGAGGCTTGGACGCCGGTGGGG
AATGGGGGGCCACAGCTGCAGACCTAAGCCCTCCCCACCCCTGGAAAGGTCCCTCCCCA
ACCCAGGCCCTGGCGTGTGTGGGTGTGCGTGCGTGTGCGTGCCGTGTTCTGTGTGCAAGG
GGCCGGGAGGTGGGCGTGTGTGTGCGTGCCAGCGAAGGCTGCTGTGGGCGTGTGTGTCA
AGTGGGCCACGCGTGCAGGGTGTGTGTCCACGAGCGACGATCGTGGTGGCCCCAGCGGCC
TGGGCGTTGGCTGAGCCGACGCTGGGGCTTCCAGAAGGCCCGGGGTCTCCGAGGTGCCG
GTTAGGAGTTTGAACCCCCCCCCACTCTGCAGAGGGAAGCGGGGACAATGCCGGGGTTTCA
GGCAGGAGACACGAGGAGGGCCTGCCCCGAAGTCACATCGGCAGCAGCTGTCTAAAGGGC
TTGGGGGCCTGGGGGGCGGCGAAGGTGGGTGGGGCCCCCTCTGTAAATACGGCCCCAGGGT
GGTGAGAGAGTCCCATGCCACCCGTCCCCTTGTGACCTCCCCCTATGACCTCCAGCTGA
CCATGCATGCCACGTGGCTGGCTGGGTCCCTCTGCCCTCTTTGGAGTTTGCTCCCCCAGC
CCCCTCCCCATCAATAAAACTCTGTTTACAACCAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

281/392

FIGURE 268

MQTPRASPPRPALLLLLLLLLLLGGAHGLFPPEPPPLSVAPRDYLNHYVPFVVGSGPGRLTPAE
GADDLNIQRLRVNRTLFIGDRDNLRYVELEPPTSTELRYQRKLTWRSNPSDINVCRMKG
KQEGECRNFKVLLLRDESTLFVCGSNAFNPCANYSIDTLQPVGDNISGMARCPYDPKH
ANVALFSDGMLFTATVTDFLAIDAVIYRSLGDRPTLRTVKHDSKWFKEPYFVHAVEWGSH
VYFFFREIAMEFNYLEKVVVSRVARVCKNDVGGSPRVLEKQWTSFLKARLNCSPVGD SHF
YFNVLQAVTGVS LGGRPVLAVFSTPSNSIPGSAVCAFDLTQVA AVFEGRFREQKSPES
IWTPVPEDQVPRPRPGCCAAPGMQYNASSALPDDILNFVKTHPLMDEAVPSLG HAPWILR
TLMRHQLTRVAVDVGAGPWGNQTVVFLGSEAGTVLKFLVRPNASTSGTSGLSVFLEEFET
YRPDRCGRPGGGETGQRLLSLELDAASGGLLA AFPRCVVRVPVARCQQYSGCMKNCIGSQ
DPYCGWAPDGSCIFLSPGTRAAFEQDVSGASTSGLGDCTGLLRASLSEDRAGLVSNNLLV
TSSVAAFVVGAVVSGFSVGWFVGLRERRELARRKDKEAILAHGAGEAVLSVSRLGERRAQ
GPGGRGGGGGGGAGVPPEALLAPLMQNGWAKATLLQGGPHDLDSGLLPTPEQTPLPQKRL
PTPHPHPHALGPRAWDHGHPLLPASASSSLLLLAPARAPEQPPAPGEPTPDGRLYAARPG
RASHGDFPLTPHASPD RRRVVSAPTGPLDPASAADGLPRPWSPPTGSLRRPLGPHAPPA
ATLRRTHTFNSGEARPGDRHRGCHARPGTDLAHL LPYGGADRTAPPVP

Important features of the protein:**Signal peptide:**

amino acids 1-25

Transmembrane domains:

amino acids 318-339, 598-617

N-glycosylation sites.amino acids 74-78, 155-159, 167-171, 291-295, 386-390,
441-445, 462-466**Glycosaminoglycan attachment sites:**

amino acids 51-55, 573-577

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 102-106

N-myristoylation sites:amino acids 21-27, 50-56, 189-195, 333-339, 382-388, 448-454,
490-496, 491-497, 508-514, 509-515, 531-537, 558-564, 569-575,
574-580, 580-586, 610-616, 643-649, 663-669, 666-672, 667-673,
668-674, 669-675, 670-676, 868-874, 879-885

282/392

FIGURE 269

ATCTGAGTGAGCTAACTGACACAATGAAACTGTCAGGCATGTTTCTGCTCCTCTCTCTGG
CTCTTTTCTGCTTTTAAACAGGTGTCTTCAGTCAGGGAGGACAGGTTGACTGTGGTGAGT
TCCAGGACCCCAAGGTCTACTGCACTCGGGAATCTAACCACACTGTGGCTCTGATGGCC
AGACATATGGCAATAAATGTGCCTTCTGTAAGGCCATAGTGAAAAGTGGTGGAAAGATTA
GCCTAAAGCATCCTGGAAAATGCTGAGTTAAAGCCAATGTTTCTTGGTGACTTGCCAGCT
TTTGCAGCCTTCTTTTCTCACTTCTGCTTATACTTTTGCTGGTGGATTCCCTTAATTCAT
AAAGACATACCTACTCTGCCTGGGTCTTGAGGAGTTCAATGTATGTCTATTTCTCTTGAT
TCACTTGTCAATAAAGTACATTCTGCAAAGCAAAA

283/392

FIGURE 270

MKLSGMFLLLSLALFCFLTGVFSQGGQVDCGEFQDPKVYCTRESNPFCGSDGQTYGNKCA
FCKAIVKSGGKISLKHPGKC

Important features of the protein:

Signal peptide:

amino acids 1-23

N-myristoylation sites:

amino acids 26-32, 52-58, 56-62, 69-75

Kazal serine protease inhibitors family signature:

amino acids 40-63

284/392

FIGURE 271

AACTTCTACATGGGCCTCCTGCTGCTGGTGCTCTTCCTCAGCCTCCTGCCGGTGGCCTAC
ACCATCATGTCCCTCCCACCCTCCTTTGACTGCGGGCCGTTCAGGTGCAGAGTCTCAGTT
GCCCCGGGAGCACCTCCCCTCCCGAGGCAGTCTGCTCAGAGGGCCTCGGCCCAGAATTCCA
GTTCTGGTTTTCATGCCAGCCTGTAAAAGGCCATGGAACTTTGGGTGAATCACCGATGCCA
TTTAAGAGGGTTTTCTGCCAGGATGGAAATGTTAGGTCGTTCTGTGTCTGCGCTGTTTCAT
TTCAGTAGCCACCAGCCACCTGTGGCCGTTGAGTGCTTGAAATGAGGAACTGAGAAAATT
AATTTCTCATGTATTTTTTCTCATTTATTTATTAATTTTTTAACTGATAGTTGTACATATTT
GGGGGTACATGTGATATTTGGATACATGTATACAATATATAATGATCAAATCAGGGTAAC
TGGGATATCCATCACATCAAACATTTATTTTTTTATTCTTTTTTAGACAGAGTCTCACTCTG
TCACCCAGGCTGGAGTGCAGTGGTGCCATCTCAGCTTACTGCAACCTCTGCCTGCCAGGT
TCAAGCGATTCTCATGCCTCCACCTCCCAAGTAGCTGGGACTACAGGCATGCACCACAAT
GCCCAACTAATTTTTTGATTTTTTAGTAGAGACGGGGTTTTTGCCATGTTGCCAGGCTGGC
CTTGAACCTCTGGCCTCAAACAATCCACTTGCCTCGGCCTCCCAAGTGTTATGATTACA
GGCGTGAGCCACCGTGCCTGGCCTAAACATTTATCTTTTCTTTGTGTTGGGAACTTTGAA
ATTATACAATGAATTATTGTTAACTGTCATCTCCCTGCTGTGCTATGGAACACTGGGACT
TCTTCCCTCTATCTAACTGTATATTTGTACCAGTTAACCAACCGTACTTCATCCCCACTC
CTCTCTATCCTTCCCAACCTCTGATCACCTCATCTACTCTCTACCTCCATGAGATCCAC
TTTTTTTAGCTCCACATGTGAGTAAGAAAATGCAATATTTGTCTTTCTGTGCCTGGCTTA
TTTCACTTAACATAATGACTTCCTGTTCCATCCATGTTGCTGCAAATGACAGGATTTTCGT
TCTTAATTTCAATTAAATAACCACACATGGCAAAA

285/392

FIGURE 272

MGLLLLVLFLSLLPVAYTIMSLPPSFDCGPFRCRVSVAREHLPSRGSLLRGPRPRI PVLV
SCQPVKGHGTLGESPMPPFKRVFCQDGNVRSFCVCAVHFSSHQPPVAVECLK

Important features of the protein:

Signal peptide:

amino acids 1-18

N-myristoylation site:

amino acids 86-92

Zinc carboxypeptidases, zinc-binding region 2 signature:

amino acids 68-79

286/392

FIGURE 273

TTCTGAAGTAACGGAAGCTACCTTGTATAAAGACCTCAACACTGCTGACCATGATCAGCG
CAGCCTGGAGCATCTTCCTCATCGGGACTAAAATTGGGCTGTTCTTCAAGTAGCACCTC
TATCAGTTATGGCTAAATCCTGTCCATCTGTGTGTCGCTGCGATGCGGGTTTCATTTACT
GTAATGATCGCTTTCTGACATCCATTCCAACAGGAATACCAGAGGATGCTACAACCTCTCT
ACCTTCAGAACAACCAAATAAATAATGCTGGGATTCTTCAGATTTGAAAACTTGCTGA
AAGTAGAAAGAATATACCTATACCACAACAGTTTAGATGAATTCCTACCAACCTCCCAA
AGTATGTAAAAGAGTTACATTTGCAAGAAAATAACATAAGGACTATCACTTATGATTAC
TTTCAAAAATTCCCTATCTGGAAGAATTACATTTAGATGACAACCTCTGTCTCTGCAGTTA
GCATAGAAGAGGGAGCATTCCGAGACAGCAACTATCTCCGACTGCTTTTCTGTCCCGTA
ATCACCTTAGCACAAATTCCTGGGGTTTGCCAGGACTATAGAAGAACTACGCTTGATG
ATAATCGCATATCCACTATTTTCATCACCATCTCTTCAAGGTCTCACTAGTCTAAAACGCC
TGGTTCTAGATGGAAACCTGTTGAACAATCATGGTTTAGGTGACAAAGTTTTCTTCAACC
TAGTTAATTTGACAGAGCTGTCCCTGGTGCGGAATTCCTGACTGCTGCACCAGTAAACC
TTCCAGGCACAAACCTGAGGAAGCTTTATCTTCAAGATAACCACATCAATCGGGTGCCCC
CAAATGCTTTTTTCTTATCTAAGGCAGCTCTATCGACTGGATATGTCCAATAATAACCTAA
GTAATTTACCTCAGGGTATCTTTGATGATTTGGACAATATAACACAACCTGATTCTTCGCA
ACAATCCCTGGTATTGCGGGTGCAAGATGAAATGGGTACGTGACTGGTTACAATCACTAC
CTGTGAAGGTCAACGTGCGTGGGCTCATGTGCCAAGCCCCAGAAAAGGTTCTGTTGGATGG
CTATTAAGGATCTCAATGCAGAACTGTTTGATTGTAAGGACAGTGGGATTGTAAGCACCA
TTCAGATAACCACTGCAATACCCAACACAGTGTATCCTGCCCAAGGACAGTGGCCAGCTC
CAGTGACCAACAGCCAGATATTAAGAACCCCAAGCTCACTAAGGATCAACAAACCACAG
GGAGTCCCTCAAGAAAAACAATTACAATTACTGTGAAGTCTGTACCTCTGATACCATT
ATATCTCTTGGAACCTTGCTCTACCTATGACTGCTTTGAGACTCAGCTGGCTTAACTGG
GCCATAGCCCGGCATTTGGATCTATAACAGAAACAATTGTAACAGGGGAACGCAGTGAGT
ACTTGGTCAACAGCCCTGGAGCCTGATTACCCCTATAAAGTATGCATGGTTCCCATGGAAA
CCAGCAACCTCTACCTATTTGATGAAACTCCTGTTTGTATTGAGACTGAACTGCACCCC
TTCGAATGTACAACCTACAACCACCTCAATCGAGAGCAAGAGAAAGAACCTTACAAAA
ACCCCAATTTACCTTTGGCTGCCATCATTTGGTGGGGCTGTGGCCCTGGTTACCATTGCCC
TTCTTGCTTTAGTGTGTTGGTATGTTTCATAGGAATGGATCGCTCTTCTCAAGGAACTGTG
CATATAGCAAAGGGAGGAGAAGAAAGGATGACTATGCAGAAGCTGGCACTAAGAAGGACA
ACTCTATCCTGGAAATCAGGGAACTTCTTTTCAGATGTTACCAATAAGCAATGAACCCA
TCTCGAAGGAGGAGTTTGTAAACACACCATATTTCTCCTAATGGAATGAATCTGTACA
AAAACAATCACAGTGAAAGCAGTAGTAACCGAAGCTACAGAGACAGTGGTATTCCAGACT
CAGATCACTCACACTCATGATGCTGAAGGACTCACAGCAGACTTGTGTTTTGGGTTTTTT
AAACCTAAGGGAGGTGATGGT

287/392

FIGURE 274

MISAAWSIFLIGTKIGLFLQVAPLSVMAKSCPSVCRC DAGFIYCND RFLTSIPTGIPEDA
TTLYLQNNQINNAGIPSDLKNLLKVERIYLYHNSLDEFPTNLPKYVKELHLQENNIRTTIT
YDSLSKIPYLEELHLLDDNSVSAVSIIEGAFRDSNYLRLLFLSRNHLSTIPWGLPRTIEEL
RLDDNRISTISSPSLQGLTSLKRLVLDGNLLNNHGLGDKVFFNLVNLTELSLVRNSLTAA
PVNLPGTNLRKLYLQDNHINRVPPNAFSYLRQLYRLDMSNNNLSNLPQGI FDDL DNITQL
ILRNNPWYCGCKMKWVRDWLQSLPVKVNVRGLMCQAPEKVRGMAIKDLNAELFDCKDSGI
VSTIQITTAIPNTVYPAQGQWPAPVTKQPDIKNPKLTKDQQTGSPSRKTITITVKS VTS
DTIHISWKLALPMTALRLSWLKLGHSPAFGSITETIVTGERSEYLVTALEPDSPYKVC MV
PMETSNLYLFDETPVCIETETAPLRMYNPTTTLNREQEKEPYKNPNLPLAAIIGGAVALV
TIALLLVCWYVHRNGSLFSRNCAYSKGRRRKDDYAEAGTKKDNSILEIRETSFQMLPIS
NEPISKEEFVIHTIFPPNGMNLKYNNHSESSNRSYRDSGIPDS DSHSHS

Important features of the protein:**Signal peptide:**

amino acids 1-28

Transmembrane domain:

amino acids 531-552

N-glycosylation sites:

amino acids 226-229, 282-285, 296-299, 555-558, 626-629, 633-636

Tyrosine kinase phosphorylation site:

amino acids 515-522

N-myristoylation sites:

amino acids 12-17, 172-177, 208-213, 359-364, 534-539, 556-561, 640-645

Amidation site:

amino acids 567-570

Leucine zipper pattern:

amino acids 159-180

Phospholipase A2 aspartic acid active site:

amino acids 34-44

288/392

FIGURE 275

AGGGCCCCGCGGGTGGAGAGAGCGACGCCCCGAGGGGATGGCGGCAGCGTCCCGGAGCGCCT
CTGGCTGGGCGCTACTGCTGCTGGTGGCACTTTGGCAGCAGCGCGCGGCCGGCTCCGGCG
TCTTCCAGCTGCAGCTGCAGGAGTTCATCAACGAGCGCGGCGTACTGGCCAGTGGGCGGC
CTTGCGAGCCCCGGCTGCCGACTTCTTCCGCGTCTGCCTTAAGCACTTCCAGGCGGTCTG
TCTCGCCCCGACCCCTGCACCTTCGGGACCGTCTCCACGCCGGTATTGGGCACCAACTCCT
TCGCTGTCCGGGACGACAGTAGCGGCGGGGGCGCAACCCTCTCCAAGTCCCTTCAATT
TCACCTGGCCGGGTACCTTCTCGCTCATCATCGAAGCTTGGCACGCGCCAGGAGACGACC
TGCGGCCAGAGGCCTTGCCACCAGATGCACTCATCAGCAAGATCGCCATCCAGGGCTCCC
TAGCTGTGGGTGAGAACTGGTTATTGGATGAGCAAACCAGCACCCCTCACAAGGCTGCGCT
ACTCTTACCGGGTCATCTGCAGTGACAACTACTATGGAGACAACTGCTCCCGCCTGTGCA
AGAAGCGCAATGACCACTTCGGCCACTATGTGTGCCAGCCAGATGGCAACTTGTCTGCC
TGCCCCGGTTGGACTGGGGAATATTGCCAACAGCCCTATCTGTCTTTCCGGGCTGTATGAAC
AGAATGGCTACTGCAGCAAGCCAGCAGAGTGCCTCTGCCGCCAGGCTGGCAGGGCCGGC
TGTGTAACGAATGCATCCCCACAATGGCTGTGCCACGGCACCTGCAGCACTCCCTGGC
AATGTACTTGTGATGAGGGCTGGGGAGGCCTGTTTTGTGACCAAGATCTCAACTACTGCA
CCCACCACTCCCCATGCAAGAATGGGGCAACGTGCTCCAACAGTGGGCAGCGAAGCTACA
CCTGCACCTGTGCGCCAGGCTACACTGGTGTGGACTGTGAGCTGGAGCTCAGCGAGTGTG
ACAGCAACCCCTGTGCGCAATGGAGGCAGCTGTAAGGACCAGGAGGATGGCTACCACTGCC
TGTGTCTCCGGGCTACTATGGCCTGCACTGTGAACACAGCACCTTGAGCTGCGCCGACT
CCCCCTGCTTCAATGGGGGCTCCTGCCGGGAGCGCAACCAGGGGGCCAACTATGCTTGTG
AATGTCCCCCAACTTCACCGGCTCCAAGTGCAGAGAAGAAAGTGGACAGGTGCACCAGCA
ACCCCTGTGCCAACGGGGGACAGTGCCTGAACCGAGGTCCAAGCCGCATGTGCCGCTGCC
GTCCTGGATTACGGGCACCTACTGTGAACCTCACGTCAGCGACTGTGCCCGTAACCCTT
GCGCCACGGTGGCACTTGCCATGACCTGGAGAATGGGCTCATGTGCACCTGCCCTGCCG
GCTTCTCTGGCCGACGCTGTGAGGTGCGGACATCCATCGATGCCTGTGCCCTCGAGTCCCT
GCTTCAACAGGGCCACCTGCTACACCGACCTCTCCACAGACACCTTTGTGTGCAACTGCC
CTTATGGCTTTGTGGGCAGCCGCTGCGAGTTCCCCGTGGGCTTGCCGCCAGCTTCCCCT
GGGTGGCCGTCTCGCTGGGTGTGGGGCTGGCAGTGTGCTGGTACTGCTGGGCATGGTGG
CAGTGGCTGTGCGGCAGCTGCGGCTTCGACGGCCGGACGACGGCAGCAGGGAAGCCATGA
ACAACTTGTGCGGACTTCCAGAAGGACAACCTGATTCTGCGCGCCAGCTTAAAAACACAA
ACCAGAAGAAGGAGCTGGAAGTGGACTGTGGCCTGGACAAGTCCAAGTGTGGCAAACAGC
AAAACCACACATTGGACTATAATCTGGCCCCAGGGCCCCCTGGGGCGGGGGACCATGCCAG
GAAAGTTTCCCACAGTGACAAGAGCTTAGGAGAGAAGGCGCCACTGCGGTTACACAGTG
AAAAGCCAGAGTGTGCGATATCAGCGATATGCTCCCCAGGGACTCCATGTACCAGTCTG
TGTGTTTGATATCAGAGGAGAGGAATGAATGTGTCAATTGCCACGGAGGTATAAGGCAGGA
GCCTACCTGGACATCCCTGCTCAGCCCCGCGCTGGACCTTCCTTCTGCATTGTTTACA

289/392

FIGURE 276

MAAASRSASGWALLLLVALWQQRAAGSGVFQLQLQEFINERGVLASGRPCPEPGCRTFFRV
CLKHFQAVVSPGPCTFGTVSTPVLGTNSFAVRDDSSGGGRNPLQLPFNFTWPGTFSLIIE
AWHAPGDDLRLPEALPPDALISKIAIQGSLAVGQNWLLDEQTSTLTRLRYSYRVICSDNYY
GDNC SRLCKKRNDHFGHYVCQPDGNLSCLPGWTGEYCQQPICLSGCHEQNGYCSKPAECL
CRPGWQGRLCNECIPHNGCRHGTCSTPWQCTCDEGWGGLFCDQDLNYCTHHSPCKNGATC
SNSGQRSYTCTCRPGYTGVDCLELSECDSNPCRNNGSCKDQEDGYHCLCPPGYGLHCE
HSTLSCADSPCFNGGSCRERNQGANYACECPPNFTGSNCEKKVDRCTSNPCANGGQCLNR
GPSRMCRCRPGFTGTYCELHVSDCARNPCAHHGGTCHDLENGLMCTCPAGFSGRRCEVRTS
IDACASSPCFN RATCYTDLSTDTFVCNCPYGFVGSRCFPVGLPPSFPWVAVSLGVGLAV
LLVLLGMVAVAVRQLRLRRPDDGSREAMNNLSDFQKDNLI PAAQLKNTNQKKELEVDCGL
DKSNCGKQQNHTLDYNLAPGPLGRGTM PGKFPHSDKSLGEKAPLRLHSEKPECRISAICS
PRDSMYQSVCLISEERNECVIATEV

Important features of the protein:**Signal peptide:**

amino acids 1-26

Transmembrane domain:

amino acids 530-552

N-glycosylation sites:amino acids 108-112, 183-187, 205-209, 393-397, 570-574,
610-614**Glycosaminoglycan attachment site:**

amino acids 96-100

Tyrosine kinase phosphorylation site:

amino acids 340-347

N-myristoylation sites:amino acids 42-48, 204-210, 258-264, 277-283, 297-303,
383-389, 415-421, 461-467, 522-528, 535-541, 563-569,
599-605, 625-631**Amidation site:**

amino acids 471-475

Aspartic acid and asparagine hydroxylation site:

amino acids 339-351

EGF-like domain cysteine pattern signature:amino acids 173-185, 206-218, 239-251, 270-282, 310-322,
348-360, 388-400, 426-438, 464-476, 506-518**Calcium-binding EGF-like:**amino acids 224-245, 255-276, 295-316, 333-354, 373-394,
411-432, 440-452

290/392

FIGURE 277

GGCAGTGCAGCCGCCTCACAGGTCGGCGGACGGGGCCAGGCGGGCGGCCTCCTGAACCGAA
CCGAATCGGCTCCTCGGGCCGTCGTCTCTCCCGCCCCCTCCTCGCCCCGCCCGGAGTTTTC
TTTCGGTTTCTTCCAAGATTCTTGGCCTTCCCTCGACGGAGCCGGGCCCAGTGCAGGGGGC
GCAGGGCGCGGGAGCTCCACCTCCTCGGCTTTCCCTGCGTCCAGAGGCTGGCATGGCGCG
GGCCGAGTACTGAGCGCACGGTCGGGGCACAGCAGGGGCCGGGGGGTGCAGCTGGCTCGCG
CCTCCTCTCCGGCCGCGTCTCCTCCGGTCCCTGGCGAAAGCCATTGAGACACCAGCTGG
ACGTACGCGCGCGGAGCATGTCTGGGAGTCAGAGCGAGGTGGCTCCATCCCCGCAGAGTC
CGCGGAGCCCCGAGATGGGACGGGACTTGCGGCCCGGGTCCCGCGTGCTCCTGCTCCTGC
TTCTGCTCCTGCTGGTGTACCTGACTCAGCCAGGCAATGGCAACGAGGGCAGCGTCACTG
GAAGTTGTTATTGTGGTAAAAGAATTTCTTCCGACTCCCCGCCATCGGTTCACTTCATGA
ATCGTCTCCGAAACACCTGAGAGCTTACCATCGGTGTCTATACTACACGAGGTTCCAGC
TCCTTTCTGGAGCGTGTGTGGGGGCAACAAGGACCCATGGGTTCAGGAATTGATGAGCT
GTCTTGATCTCAAAGAATGTGGACATGCTTACTCGGGGATTGTGGCCCACCAGAAGCATT
TACTTCTTACCAGCCCCCAATTTCTCAGGCCTCAGAGGGGGCATCTTCAGATATCCACA
CCCCTGCCCAGATGCTCCTGTCCACCTTGCACTCCACTCAGCGCCCCACCCTCCCAGTAG
GATCACTGTCTCGGACAAAGAGCTCACTCGTCCCAATGAAACCACCATTACACTGCGG
GCCACAGTCTGGCAGCTGGGCCTGAGGCTGGGGAGAACCAGAAGCAGCCGGAAAAAATG
CTGGTCCCACAGCCAGGACATCAGCCACAGTGCCAGTCCTGTGCCTCCTGGCCATCATCT
TCATCCTCACCGCAGCCCTTTCTTATGTGCTGTGCAAGAGGAGGAGGGGGCAGTCACCGC
AGTCCTCTCCAGATCTGCCGGTTCATTATATACCTGTGGCACCTGACTCTAATACCTGAG
CCAAGAATGGAAGCTTGTGAGGGTAACTGTGGCTTATTCTTACAAAAAGTGTAAATAAAG
GAGACTGACCCCTGACAACATGGTAGGCACTGTAAAAA

291/392

FIGURE 278

MGRDLRPGSRVLLLLLLLLLVYLTQPGNGNEGSVTGSCYCGKRISSDSPPSVQFMNRLRK
HLRAYHRCLYYTRFQLLSWSVCGGNKDPWVQELMSCLDLKECGHAYSGIVAHQKHLPTS
PPISQASEGASSDIHTPAQMLLSTLQSTQRPTLPVGSLSDDKELTRPNETTIHTAGHSLA
AGPEAGENQKQPEKNAGPTARTSATVPVLCLLAIIFILTAALSYVLCKRRRGQSPQSSPD
LPVHYIPVAPDSNT

Important features of the protein:

Signal peptide:

1-26

Transmembrane domain:

204-223

N-glycosylation site:

168-172

cAMP- and cGMP-dependent protein kinase phosphorylation site:

42-46

N-myristoylation site:

29-35, 32-38, 36-42, 156-162

Amidation site:

40-44

292/392

FIGURE 279

CGCGAGGCGCGGGGAGCCTGGGACCAGGAGCGAGAGCCGCTACCTGCAGCCGCCGCCCA
CGGCACGGCAGCCACCCATGGCGCTCCTGCTGTGCTTCGTGCTCCTGTGCGGAGTAGTGGA
TTTCGCCAGAAGTTTGTAGTATCACTACTCCTGAAGAGATGATTGAAAAAGCCAAAGGGGA
AACTGCCTATCTGCCATGCAAATTTACGCTTAGTCCCGAAGACCAGGGACCGCTGGACAT
CGAGTGGCTGATATCACCAGCTGATAATCAGAAGGTGGATCAAGTGATTATTTTATATTTC
TGGAGACAAAATTTATGATGACTACTATCCAGATCTGAAAGGCCGAGTACATTTTACGAG
TAATGATCTCAAATCTGGTGATGCATCAATAAATGTAACGAATTTACAACGTGCAGATAT
TGGCACATATCAGTGCAAAGTGAAAAAGCTCCTGGTGTTGCAAATAAGAAGATTCATCT
GGTAGTTCTTGTTAAGCCTTCAGGTGCGAGATGTTACGTTGATGGATCTGAAGAAATTGG
AAGTGACTTTAAGATAAAATGTGAACCAAAAGAAGGTTCACTTCCATTACAGTATGAGTG
GCAAAAATTGTCTGACTCACAGAAAATGCCACTTCATGGTTAGCAGAAATGACTTCATC
TGTTATATCTGTAAAAAATGCCTCTTCTGAGTACTCTGGGACATACAGCTGTACAGTCAG
AAACAGAGTGGGCTCTGATCAGTGCCCTGTTGCGTCTAAACGTTGTCCCTCCTTCAAATAA
AGCTGGACTAATTGCAGGAGCCATTATAGGAACTTTGCTTGCTCTAGCGCTCATTGGTCT
TATCATCTTTTGCTGTCTGTAAAAAGCGCAGAGAAGAAAAATATGAAAAGGAAGTTCATCA
CGATATCAGGGAAGATGTGCCACCTCCAAAGAGCCGTACGTCCACTGCCAGAAGCTACAT
CGGCAGTAATCATTCATCCCTGGGGTCCATGTCTCCTTCCAACATGGAAGGATATTCCAA
GACTCAGTATAACCAAGTACCAAGTGAAGACTTTGAACGCACTCCTCAGAGTCCGACTCT
CCCACCTGCTAAGTTCAAGTACCCTTACAAGACTGATGGAATTACAGTTGTATAAATATG
GACTACTGAAGAATCTGAAGTATTGTATTATTTGACTTTATTTTAGGCCTCTAGTAAAGA
CTTAAATGTTTTTTAAAAAAGCACAAAGGCACAGAGATTAGAGCAGCTGTAAGAACACAT
CTACTTTATGCAATGGCATTAGACATGTAAGTCAGATGTCATGTCAAAATTAGTACGAGC
CAAATTCCTTTGTTAAAAAACCTATGTATAGTGACACTGATAGTTAAAAGATGTTTTATT
ATATTTTCAATAACTACCACTAACAAATTTTTTAACCTTTTCATATGCATATTCCTGATATGT
GGTCTTTTAGGAAAAGTATGGTTAATAGTTGATTTTTTCAAAGGAAATTTTAAAATTCCTTA
CGTTCGTGTTAATGTTTTTGCTATTTAGTTAAATACATTGAAGGGAAATACCCGTTCTTT
TCCCCTTTTATGCACACAACAGAAACACGCGTTGTCATGCCTCAAACATTTTTTTATTTG
CAACTACATGATTTACACAATTCCTTAAACAACGACATAAAATAGATTTCCCTGTATA
TAAATAACTTACATACGCTCCATAAAGTAAATTCCTCAAAGGTGCTAGAACAAATCGTCCA
CTTCTACAGTGTTCTCGTATCCAACAGAGTTGATGCACAATATATAAATACTCAAGTCCA
ATATTA AAAA ACTTAGGCACCTTGACTA ACTTTAATAAAATTTCTCAAACATATCAATATC
TAAAGTGCATATATTTTTTAAGAAAGATTATTCTCAATAACTTCTATAAAAAATAAGTTTG
ATGGTTTGGCCCATCTAACTTCACTACTATTAGTAAGAACTTTTAACTTTTAAATGTGTAG
TAAGGTTTATTCTACCTTTTTCTCAACATGACACCAACACAATCAAAAACGAAGTTAGTG
AGGTGCTAACATGTGAGGATTAATCCAGTGATTCCGGTCACAATGCATTCCAGGAGGAGG
TACCCATGTCACTGGAATTGGGCGATATGGTTTATTTTTTCTTCCCTGATTTGGATAACC
AAATGGAACAGGAGGAGGATAGTGATTCTGATGGCCATTCCCTCGATACATTCTTGGCTT
TTTTCTGGGCAAAGGTGCCACATTGGAAGAGGTGGAAATATAAGTTCTGAAATCTGTAG
GGAAGAGAACACATTAAGTTAATTCAAAGGAAAAAATCATCATCTATGTTCCAGATTTCT
CATTAAAGACAAAGTTACCCACAACACTGAGATCACATCTAAGTGACACTCCTATTGTCA
GGTCTAAATACATTAAAAACCTCATGTGTAATAGGCGTATAATGTATAACAGGTGACCAA
TGTTTTCTGAATGCATAAAGAAATGAATAAACTCAAACACAGTACTTCCTAAACAACTTC
AACCAAAAAAGACCAAAACATGGAACGAATGGAAGCTTGTAAGGACATGCTTGTTTTAGT
CCAGTGGTTTCCACAGCTGGCTAAGCCAGGAGTCACTTGAGAGGCTTTTAAATACAAAACA
TTGGAGCTGGAGGCCATTATCCTTAGCAAACCTAATGCAGAAACAGAAAATCAACTACCGC
ATGTTCTCACTTATAAGTGGGAGGTAATGATAAGAACTTATGAACACAAAGAAGGAAACA
ATAGACATTGGAGTCTATTTGAGAGGGGAGGGTGGGAGAAGGAAAAGGAGCAGAAAAGAT
AACTATTGAGTACTGCCTTCACACCTGGGTGATGAAATAATATGTACAACAAATCCCTGT
GACACATGTTTACCTATGGAACAAACCTTCATGTGTATCCCTAAACCTAAAATAAAAGTT

293/392

FIGURE 280

MALLLCFVLLCGVVDFARSLSTTTPEEMIEKAKGETAYLPCKFTLSPEDQGPLDIEWLIS
PADNQKVDQVIILYSGDKIYDDYYPDLKGRVHFTSNDLKSGDASINVTNLQLSDIGTYQC
KVKKAPGVANKKIHLVVLVKPSGARCIVDGSSEEIGSDFKIKCEPKEGSLPQYEWQKLS
SQKMPTSWLAEMTSSVISVKNASSEYSGTYSCTVRNRVGSQCLLRNLNVPPSNKAGLIA
GAIIGTLLALALIGLIIFCCRKKRREEKYEKEVHHDIREDVPPPKSRTSTARSYIGSNHS
SLGSMSPSNMEGYSKTQYNQVPSEDFERTPQSPTLPPAKFKYPYKTDGITVV

Signal sequence.

amino acids 1-19

Transmembrane domain:

amino acids 236-257

N-glycosylation sites:

amino acids 106-110, 201-205, 298-302

Tyrosine kinase phosphorylation sites:

amino acids 31-39, 78-85, 262-270

N-myristoylation sites:amino acids 116-122, 208-214, 219-225, 237-243, 241-247,
245-251, 296-302**Myelin P0 protein:**

amino acids 96-125

294/392

FIGURE 281

TGCATCAGTGCCCAGGCAAGCCCAGGAGTTGACATTTCTCTGCCCAGCCATGGGCCTCAC
CCTGCTCTTGCTGCTGCTCCTGGGACTAGAAGGTCAGGGCATAGTTGGCAGCCTCCCTGA
GGTGCTGCAGGCACCCGTGGGAAGCTCCATTCTGGTGCCAGTGCCACTACAGGCTCCAGGA
TGTCAAAGCTCAGAAGGTGTGGTGCCGGTTCTTGCCGGAGGGGTGCCAGCCCCCTGGTGTC
CTCAGCTGTGGATCGCAGAGCTCCAGCGGGCAGGCGTACGTTTCTCACAGACCTGGGTGG
GGGCCTGCTGCAGGTGGAAATGGTTACCCTGCAGGAAGAGGATGCTGGCGAGTATGGCTG
CATGGTGATGGGGCCAGGGGGCCCCAGATTTTGACAGAGTCTCTCTGAACATACTGCC
CCCAGAGGAAGAAGAAGAGACCCATAAGATTGGCAGTCTGGCTGAGAACGCATTCTCAGA
CCCTGCAGGCAGTGCCAACCCCTTTGGAACCCAGCCAGGATGAGAAGAGCATCCCCCTTGAT
CTGGGGTGCTGTGCTCCTGGTAGGTCTGCTGGTGGCAGCGGTGGTGCTGTTTGCTGTGAT
GGCCAAGAGGAAACAAGAATCCCTCCTCAGTGGTCCACCACGTCAGTGACTCTGGACCGG
CTGCTGAATTGCCTTTGGATGTACCACACATTAGGCTTGACTCACCACCTTCATTTGACA
ATACCACCTACACCAGCCTACCTCTTGATTCCCCATCAGGAAAACCTTCACTCCCAGCTC
CATCCTCATTGCCCCCTCTACCTCCTAAGGTCCTGGTCTGCTCCAAGCCTGTGACATATG
CCACAGTAATCTTCCCGGGAGGGAACAAGGGTGGAGGGACCTCGTGTGGGCCAGCCCAGA
ATCCACCTAACAATCAGACTCCATCCAGCTAAGCTGCTCATCACACTTTAAACTCATGAG
GACCATCCCTAGGGGTTCTGTGCATCCATCCAGCCAGCTCATGCCCTAGGATCCTTAGGA
TATCTGAGCAACCAGGGACTTTAAGATCTAATCCAATGTCCTAACTTTACTAGGGAAAGT
GACGCTCAGACATGACTGAGATGTCTTGGGGAAGACCTCCCTGCACCCAACTCCCCACT
GGTTCTTCTACCATTACACACTGGGCTAAATAAACCTAATAATGATGTGCAAAAAAAAAA
AA

295/392

FIGURE 282

MGLTLLLLLLLLLGLGQGIVGSLPEVLQAPVGSSILVQCHYRLQDVKAQKVWCRFLPEGCQ
PLVSSAVDRRAPAGRRTFLTDLGGGLLQVEMVTLQEEDAGEYGCMVDGARGPQILHRVSL
NILPPEEEEEETHKIGSLAENAFSDPAGSANPLEPSQDEKSIPLIWGAVLLVGLLVAAVVL
FAVMAKRKQESLLSGPPRQ

Important features of the protein:

Signal peptide:

amino acids 1-15

Transmembrane domain:

amino acids 161-181

N-myristoylation sites:

amino acids 17-23, 172-178

Amidation site:

amino acids 73-79

296/392

FIGURE 283

GTAGCATAGTGTGCAGTTCACCTGGACCAAAAGCTTTGGCTGCACCTCTTCTGGAAAGCTG
GCCATGGGGCTCTTCATGATCATTGCAATTCTGCTGTTCCAGAAACCCACAGTAACCGAA
CAACTTAAGAAGTGCTGGAATAACTATGTACAAGGACATTGCAGGAAAATCTGCAGAGTA
AATGAAGTGCCTGAGGCACTATGTGAAAATGGGAGATACTGTTGCCTCAATATCAAGGAA
CTGGAAGCATGTAAAAAATTACAAAGCCACCTCGTCCAAAGCCAGCAACACTTGCCTG
ACTCTTCAAGACTATGTTACAATAATAGAAAATTTCCCAAGCCTGAAGACACAGTCTACA
TAAATCAAATACAATTTCTGTTTTCACTTGCTTCTCAACCTAGTCTAATAAACTAAGGTGA
TGAGATATACATCTTCTTCTTCTGGTTTCTTGATCCTTAAAATGACCTTCGAGCATATT
CTAATAAAGTGCATTGCCAGTTAAAAA

297/392

FIGURE 284

MGLFMIIAILLFQKPTVTEQLKKCWNNYVQGHCRKICRVNEVPEALCENGRYCCLNIKEL
EACKKITKPPRPKPATLALTLDYVTIIENFPSLKTQST

Important features of the protein:

Signal peptide:

None

Transmembrane domain:

None

cAMP- and cGMP-dependent protein kinase phosphorylation site:

64-68

298/392

FIGURE 285

GATGGCGCAGCCACAGCTTCTGTGAGATTTCGATTTCTCCCCAGTTCCCCTGTGGGTCTGA
GGGGACCAGAAGGGTGAGCTACGTTGGCTTTCTGGAAGGGGAGGCTATATGCGTCAATTC
CCCCAAAACAAGTTTGTGACATTTCCCCTGAAATGTCATTCTCTATCTATTCACTGCAAGTG
CCTGCTGTTCCAGGCCTTACCTGCTGGGCACTAACGGCGGAGCCAGGATGGGGACAGAAT
AAAGGAGCCACGACCTGTGCCACCAACTCGCACTCAGACTCTGAACTCAGACCTGAAATC
TTCTCTTCACGGGAGGCTTGGCAGTTTTTCTTACTCCTGTGGTCTCCAGATTTCAGGCCT
AAGATGAAAGCCTCTAGTCTTGCCCTTCAGCCTTCTCTCTGCTGCGTTTTATCTCCTATGG
ACTCCTTCCACTGGACTGAAGACACTCAATTTGGGAAGCTGTGTGATCGCCACAAACCTT
CAGGAAATACGAAATGGATTTTCTGAGATACGGGGCAGTGTGCAAGCCAAAGATGGAAAC
ATTGACATCAGAATCTTAAGGAGGACTGAGTCTTTGCAAGACACAAAGCCTGCGAATCGA
TGCTGCCTCCTGCGCCATTTGCTAAGACTCTATCTGGACAGGGTATTTAAAACTACCAG
ACCCCTGACCATTATACTCTCCGGAAGATCAGCAGCCTCGCCAATTCCTTTCTTACCATC
AAGAAGGACCTCCGGCTCTCTCATGCCCACATGACATGCCATTGTGGGGAGGAAGCAATG
AAGAAATACAGCCAGATTCTGAGTCACTTTGAAAAGCTGGAACCTCAGGCAGCAGTTGTG
AAGGCTTTGGGGGAAGTAGACATTCTTCTGCAATGGATGGAGGAGACAGAATAGGAGGAA
AGTGATGCTGCTGCTAAGAATATTCGAGGTCAAGAGCTCCAGTCTTCAATACCTGCAGAG
GAGGCATGACCCCAAACCACCATCTCTTTACTGTACTAGTCTTGTGCTGGTCACAGTGTA
TCTTATTTATGCATTACTTGCTTCCTTGCAATGATTGTCTTTATGCATCCCCAATCTTAAT
TGAGACCATACTTGTATAAGATTTTGTAAATATCTTTCTGCTATTGGATATATTTATTAG
TTAATATATTTATTTATTTTGTCTATTTAATGTATTTATTTTTTTACTTGGACATGAAA
CTTTAAAAAAATTCACAGATTATATTTATAACCTGACTAGAGCAGGTGATGTATTTTTAT
ACAGTAAAAAAAACCTTGTAATTCAGAAAGAGTGGCTAGGGGGGTATTTCATTTG
TATTCAACTAAGGACATATTTACTCATGCTGATGCTCTGTGAGATATTTGAAATTGAACC
AATGACTACTTAGGATGGGTGTGGAATAAGTTTTGATGTGGAATTGCACATCTACCTTA
CAATTACTGACCATCCCCAGTAGACTCCCCAGTCCCATAATTGTGTATCTTCCAGCCAGG
AATCCTACACGGCCAGCATGTATTTCTACAAATAAAGTTTTCTTGCATACCAAAAAAAA
AAAAAAA

299/392

FIGURE 286

MKASSLAFSLLSAAFYLLWTPSTGLKTLNLGSCVIATNLQEIRNGFSEIRGSVQAKDGN
DIRILRRTESLQDTKPANRCCLLRHLLRLYLDRVFKNYQTPDHYTLRKISSLANSLFTIK
KDLRLSHAHMTCHCGEEAMKKYSQILSHFEKLEPQAAVVKALGELDILLQWMEETE

300/392

FIGURE 287

AATGCCCCATGCGCACCCACAGCTCGCGCTCCTGCAAGTGTTCTTTCTGGTGTTCCTCG
ATGGCGTCCGGCCTCAGCCCTCTTCTCCCCATCAGGGGCAGTGCCACGTCTTTGGAGC
TGCAGCGAGGGACGGATGGCGGAACCTCCAGTCCCCTTCAGAGGCGACTGCAACTCGCC
CGGCCGTGCCTGGACTCCCTACAGTGGTCCCTACTCTCGTGACTCCCTCGGCCCCCTGGGA
ATAGGACTGTGGACCTCTTCCCAGTCTTACCGATCTGTGTCTGTGACTTGACTCCTGGAG
CCTGCGATATAAATTGCTGCTGCGACAGGGACTGCTATCTTCTCCATCCGAGGACAGTTT
TCTCCTTCTGCCTTCCAGGCAGCGTAAGGTCTTCAAGCTGGGTTTGTGTAGACAACTCTG
TTATCTTCAGGAGTAATTCCCCGTTTCCTTCAAGAGTTTTTCATGGATTCTAATGGAATCA
GGCAGTTTTGTGTCCATGTGAACAACTCAAACCTTAAACTATTTCCAGAAGCTTCAAAGG
TCAATGCAACCAACTTCCAGGCCCTGGCTGCAGAGTTTGGAGGCGAATCATTCACTTCAA
CATTCCAAACTCAATCACCACCATCTTTTTACAGGGCTGGGGACCCCATCTTACTTACT
TCCCCAAGTGGTCTGTAATAAGCTTGCTGAGACAACTGCAGGAGTTGGAGCTGGGGGAC
TCTGTGCTGAAAGCAATCCTGCAGGTTTCCTAGAGAGTAAAAGTACAACTTGCACTCGTT
TTTTCAAGAACCTGGCTAGTAGCTGTACCTTGGATTGAGCCCTCAATGCTGCCTCTTACT
ATAACTTCACAGTCTTAAAGGTTCCAAGAAGCATGACTGATCCACAGAATATGGAGTTCC
AGGTTCTGTAATACTTACCTCACAGGCTAATGCTCCTCTGTTGGCTGGAAACACTTGTC
AGAATGTAGTTTCTCAGGTCACCTATGAGATAGAGACCAATGGGACTTTTGGAAATCCAGA
AAGTTTCTGTGAGTTTGGGACAAACCAACCTGACTGTTGAGCCAGGCGCTTCCTTACAGC
AACACTTCATCCTTCGCTTCAGGGCTTTTCAACAGAGCACAGCTGCTTCTCTCACCAGTC
CTAGAAGTGGGAATCCTGGCTATATAGTTGGGAAGCCACTCTTGGCTCTGACTGATGATA
TAAGTTACTCAATGACCCTCTTACAGAGCCAGGGTAATGGAAGTTGCTCTGTTAAAAGAC
ATGAAGTGCAGTTTGGAGTGAATGCAATATCTGGATGCAAGCTCAGGTTGAAGAAGGCAG
ACTGCAGCCACTTGCAGCAGGAGATTTATCAGACTCTTCATGGAAGGCCAGACCAGAGT
ATGTTGCCATCTTTGGTAATGCTGACCCAGCCAGAAAGGAGGGTGGACCAGGATCCTCA
ACAGGCACTGCAGCATTTCAGCTATAAACTGTACTTCCTGCTGTCTCATAACAGTTTCCC
TGGAGATCCAGGTATTGTGGGCATATGTAGGTCTCCTGTCCAACCCGCAAGCTCATGTAT
CAGGAGTTCGATTCTTATACCAGTGCCAGTCTATACAGGATTCTCAGCAAGTTACAGAAG
TATCTTTGACAACTCTTGTGAACCTTTGTGGACATTACCCAGAAGCCACAGCCTCCAAGGG
GCCAACCCAAAATGGACTGGAAATGGCCATTGACTTCTTTCCCTTCAAAGTGGCATTCA
GCAGAGGAGTATTCTCTCAAAAATGCTCAGTCTCTCCCATCCTTATCCTGTGCCTCTTAC
TACTTGGAGTTCTCAACCTAGAGACTATGTGAAGAAAAGAAAATAATCAGATTTTCAGTTT
TCCCTATGAGAACTCTGAGGCAGCCACTTATCTTGGCTAAATAGAACCTCACCTGCTCA
TGACCAGAGAGCATTTAGGATAATAGATGACCTAACTGAAGGAATCCTTGTATATGAAAG
GAGTTATTTTAGAAAAGCAATAAAAATATTTTATTCATCNTAAAAAAAAA

301/392

FIGURE 288

MRTPQLALLQVFFLVFPDGVPRQPSSSSPSGAVPTSLELQRGTDGGTLQSPSEATATRP
PGLPTVVPTLVTPSAPGNRTVDLFPVLPICVCDLTPGACDINCCCDRDCYLLHPRTVFSF
CLPGSVRSSSWVCVDNSVIFRSNSPFPSPRVFMDSNQIRQFCVHVNNNSLNYFQKLQKVNA
TNFQALAAEFGGESFTSTFQTQSPPSFYRAGDPILTYFPKWSVISLLRQPAGVGAGGLCA
ESNPAGFLESKSTTCTRFFKNLASSCTLDSALNAASYNFTVLKVPRSM TDPQNMEFQVP
VILTSQANAPLLAGNTCQNVVSQVTYEIETNGTFGIQKVS VSLGQTNLTVEPGASLQQHF
ILRFRAFQQSTAASLTSPRSGNPGYIVGKPLLALTDDISYSMTLLQSQNGSGCSVKRHEV
QFGVNAISGCKLRLKKADCSHLQQEIYQTLHGRPRPEYVAIFGNADPAQKGGWTRILNRH
CSISAINCTSCCLIPVSLEIQVLWAYVGLLSNPQAHVSGVRFLYQCQSIQDSQQVTEVSL
TTLVNFVDITQKPQPPRGQPKMDWKWPFDFPFKVAFSRGVFSQKCSVSPILILCLLLLG
VLNLETM

Important features of the protein:**Signal peptide:**

amino acids 1-22

Transmembrane domains:

amino acids 484-505, 581-600

N-glycosylation sites:amino acids 78-82, 165-169, 179-185, 279-285, 331-337,
347-351, 410-414, 487-491**N-myristoylation sites:**

amino acids 30-36, 41-47, 124-130, 232-238, 236-242, 409-415

Prokaryotic membrane lipoprotein lipid attachment site:

amino acids 420-431

302/392

FIGURE 289

CGCGGAGCCCTGCGCTGGGAGGTGCACGGTGTGCACGCTGGACTGGACCCCCATGCAACC
CCGCGCCCTGCGCCTTAACCAGGACTGCTCCGCGCGCCCCTGAGCCTCGGGCTCCGGCCC
GGACCTGCAGCCTCCAGGTGGCTGGGAAGAACTCTCCAACAATAAATACATTTGATAAG
AAAGATGGCTTTAAAAGTGCTACTAGAACAAGAGAAAACGTTTTTCACTCTTTTAGTATT
ACTAGGCTATTTGTTCATGTAAAGTGACTTGTGAATCAGGAGACTGTAGACAGCAAGAATT
CAGGGATCGGTCTGGAACTGTGTTCCCTGCAACCAGTGTGGGCCAGGCATGGAGTTGTC
TAAGGAATGTGGCTTCGGCTATGGGGAGGATGCACAGTGTGTGACGTGCCGGCTGCACAG
GTTCAAGGAGGACTGGGGCTTCAGAAATGCAAGCCCTGTCTGGACTGCGCAGTGGTGAA
CCGCTTTTCAAGAGGCAAATTGTTTCAGCCACCAGTGATGCCATCTGCGGGGACTGCTTGCC
AGGATTTTATAGGAAGACGAACTTGTTCGGCTTTCAAGACATGGAGTGTGTGCCTTGTTG
AGACCTCTCTCTCTTACGAACCGCACTGTGCCAGCAAGGTCAACCTCGTGAAGATCGC
GTCCACGGCCTCCAGCCACGGGACACGGCGCTGGCTGCCGTTATCTGCAGCGCTCTGGC
CACCGTCTGTGGCCCTGCTCATCTCTGTGTCTATCTATTGTAAGAGACAGTTTATGGA
GAAGAAACCAGCTGGTCTCTGCGGTGCGAGGACATTCAGTACAACGGCTCTGAGCTGTC
GTGTTTTGACAGACCTCAGCTCCACGAATATGCCACAGAGCCTGCTGCCAGTGCCGCCG
TGACTCAGTGCAGACCTGCGGGCCGGTGCCTTGTCTCCATCCATGTGCTGTGAGGAGGC
CTGCAGCCCCAACCCGGCGACTCTTGTTGTGGGGTGCATTCTGCAGCCAGTCTTCAGGC
AAGAAACGCAGGCCCAGCCGGGAGATGGTGCCGACTTCTTCGGATCCCTCACGCAGTC
CATCTGTGGCGAGTTTTTCAGATGCCTGGCCTCTGATGCAGAATCCCATGGGTGGTGACAA
CATCTCTTTTTGTGACTCTTATCTGAACCTCACTGGAGAAGACATTCATTCTCTCAATCC
AGAACTTGAAAGCTCAACGTCTTTTGATTCAAATAGCAGTCAAGATTTGTTGGTGGGGC
TGTTCCAGTCCAGTCTCATTCTGAAAACCTTACAGCAGCTACTGATTTATCTAGATATAA
CAACACACTGGTAGAATCAGCATCAACTCAGGATGCACTAACTATGAGAAGCCAGCTAGA
TCAGGAGAGTGGCGCTGTCTATCCACCCAGCCACTCAGACGTCCCTCCAGGAAGCTTAAAG
AACCTGCTTCTTTCTGCAGTAGAAGCGTGTGCTGGAACCCAAAGAGTACTCCTTTGTTAG
GCTTATGGAAGTGCAGTCTGGACCTTGCATGGCTTCTGGGGCAAAAATAAATCTGAACC
AACTGACGGCATTGTAAGCCTTTTCAGCCAGTTGCTTCTGAGCCAGACCAGCTGTAAGCT
GAAACCTCAATGAATAACAAGAAAAGACTCCAGGCCGACTCATGATACTCTGCATCTTTC
CTACATGAGAAGCTTCTCTGCCACAAAAGTGACTTCAAAGACTGATGGGTGAGCTGGCA
GCCTATGAGATTGTGGACATATAACAAGAAACAGAAATGCCCTCATGCTTATTTTCATGG
TGATTGTGGTTTTACAAGACTGAAGACCCAGAGTATACTTTTTCTTCCAGAAATAATTT
CATACCGCTATGAAATATCAGATAAATTACCTTAGCTTTTATGTAGAATGGGTTCAAA
GTGAGTGTCTTATTTGAGAAGGACACTTTTTTCATCATCTAACTGATTCGCATAGGTGG
TTAGAATGGCCCTCATATTGCCTGCCTAAATCTTGGGTTTATTAGATGAAGTTTACTGAA
TCAGAGGAATCAGACAGAGGAGGATAGCTCTTCCAGAATCCACACTTCTGACCTCAGCC
TCGGTCTCATGAACACCCGCTGATCTCAGGAGAACACCTGGGCTAGGGAATGTGGTCGAG
AAAGGGCAGCCCATGCCCAGAAATTAACACATATTGTAGAGACTTGTATGCAAAGGTTGG
CATATTTATATGAAAATTAGTTGCTATAGAAACATTTGTTGCATCTGTCCCTCTGCCTGA
GCTTAGAAGGTTATAGAAAAGGGTATTTATAAACATAAATGACCTTTTACTTGCATTGT
ATCTTATACTAAAGGCTTTAGAAATTACAACATATCAGGTTCCCTACTACTGAAGTAGC
CTTCCGTGAGAACACACCACATGTTAGGACTAGAAGAAAATGCACAATTTGTAGGGGTTT
GGATGAAGCAGCTGTAACCTGCCCTAGTGTAGTTTGACCAGGACATTGTCTGTCTCCTTCC
AATTGTGTAAGATTAGTTAGCACATCATCTCCTACTTTAGCCATCCGGTGTGGATTAA
GAGGACGGTGCTTCTTTCTATTAAAGTGCTCCATCCCTACCATCTACACATTAGCATTG
TCTCTAGAGCTAAGACAGAAATTAACCCCGTTCAGTCAAAAGCAGGGAATGGTTCAATTT
ACTCTTAATCTTTATGCCCTGGAGAAGACCTACTTGAACAGGGCATATTTTTTTAGACTTC
TGAACATCAGTATGTTTCGAGGGTACTATGATATTTTGGTTTGAATTGCCCTGCCAAGT
CACTGTCTTTTAACTTTTAACTGAATATTAAATGTATCTGTCTTCT

303/392

FIGURE 290

MALKVLLLEQEKTFFTLLVLLGYLSCKVTCESGDCRQQEFRDRSGNCVPCNQCGPGMELSK
ECGFGYGEDAQCVTCRLHRFKEDWGFQKCKPCLDCAVVNRFOKANCSATSDAICGDCLPG
FYRKTKLVGFQDMECVPCGDPPPPYEPHCASKVNLVKIASTASSPRDTALAAVICSALAT
VLLALLILCVIYCKRQFMKKPSWSLSQDIQYNGSELSCFDRPQLHEYAHRACCQCRRD
SVQTCGPVRLLLPSMCCEEACSPNPATLGCGVHSAASLQARNAGPAGEMVPTFFGSLTQSI
CGEFSDAWPLMQNPMGGDNISFCDSYPELTGEDIHSLNPELESSTSLDSNSSQDLVGGAV
PVQSHSENFTAATDLRYNNTLVESASTQDALTMRSQLDQESGAVIHPATQTSLQEA

Important features of the protein:**Signal peptide:**

Amino acids 1-25

Transmembrane domain:

Amino acids 169-192

N-glycosylation sites:

Amino acids 105-109; 214-218; 319-323; 350-354; 368-372; 379-383

cAMP- and cGMP-dependent protein kinase phosphorylation sites:

Amino acids 200-204; 238-242

Tyrosine kinase phosphorylation site:

Amino acids 207-214

N-myristoylation sites:

Amino acids 55-61; 215-221; 270-276

Prokaryotic membrane lipoprotein lipid attachment site:

Amino acids 259-270

TNFR/NGFR family cysteine-rich region proteins:

Amino acids 89-96

304/392

FIGURE 291

CCTGGAGCCGGAAGCGCGGCTGCAGCAGGGCGAGGCTCCAGGTGGGGTTCGGTTCGCGATC
CAGCCTAGCGTGTCACGATGCGGCTGGGCTCCGGGACTTTTCGCTACCTGTTGCGTAGCG
ATCGAGGTGCTAGGGATCGCGGTCTTCCTTCGGGGATTCTTCCCGGCTCCCGTTCGTTCC
TCTGCCAGAGCGGAACACGGAGCGGAGCCCCAGCGCCCGAACCTCGGCTGGAGCCAGT
TCTAACTGGACCACGCTGCCACCACCTCTCTTCAGTAAAGTTGTTATTGTTCTGATAGAT
GCCTTGAGAGATGATTTTGTGTTTGGGTCAAAGGGTGTGAAATTTATGCCCTACACAAC
TACCTTGTTGAAAAAGGAGCATCTCACAGTTTGTGGCTGAAGCAAAGCCACCTACAGTT
ACTATGCCTCGAATCAAGGCATTGATGACGGGGAGCCTTCCTGGCTTGTGCGAGTCATC
AGGAACCTCAATTCTCCTGCACTGCTGGAAGACAGTGTGATAAGACAAGCAAAAGCAGCT
GGAAAAAGAATAGTCTTTTATGGAGATGAAACCTGGGTAAATTATTCCCAAAGCATTTT
GTGGAATATGATGGAACAACCTCATTTTTCTGTGTCAGATTACACAGAGGTGGATAATAAT
GTCACGAGGCATTTGGATAAAGTATTAAAAAGAGGAGATTGGGACATATTAATCCTCCAC
TACCTGGGGCTGGACCACATTGGCCACATTTAGGGCCCAACAGCCCCCTGATTGGGCAG
AAGCTGAGCGAGATGGACAGCGTGCTGATGAAGATCCACACCTCACTGCAGTCGAAGGAG
AGAGAGACGCCTTTACCCAATTTGCTGGTTCTTTGTGGTGACCATGGCATGTCTGAAACA
GGAAGTCACGGGGCTCCTCCACCGAGGAGGTGAATACACCTCTGATTTTAATCAGTTCT
GCGTTTGAAAGGAAACCCGGTGATATCCGACATCCAAAGCACGTCCAATAGACGGATGTG
GCTGCGACACTGGCGATAGCACTTGCTTACCGATTCCAAAAGACAGTGTAGGGAGCCTC
CTATTCCAGTTGTGGAAGGAAGACCAATGAGAGAGCAGTTGAGATTTTACATTTGAAT
ACAGTGCAGCTTAGTAACTGTTGCAAGAGAATGTGCCGTATATGAAAAAGATCCTGGG
TTTGAGCAGTTTAAATGTCAGAAAGATTGCATGGGAACTGGATCAGACTGTACTTGGAG
GAAAAGCATTCAGAAAGTCTTATTCAACCTGGGCTCCAAGGTTCTCAGGCAGTACCTGGAT
GCTCTGAAGACGCTGAGCTTGTCCCTGAGTGCACAAGTGGCCCAGTTCTCACCCCTGCTCC
TGCTCAGCGTCCCACAGGCACTGCACAGAAAGGCTGAGCTGGAAGTCCCACTGTCTCTC
CTGGGTTTCTCTGCTCTTTTATTGTTGATCCTGGTTCTTTCGGCCGTTACGTCATTG
TGTGCACCTCAGCTGAAAGTTCTGTCTACTTCTGTGGCCTCTCGTGGCTGGCGGCAGGCT
GCCTTTCGTTTACCAGACTCTGGTTGAACACCTGGTGTGTGCCAAGTGCTGGCAGTGCCC
TGGACAGGGGGCCTCAGGGAAGGACGTGGAGCAGCCTTATCCAGGCCTCTGGGTGTCCC
GACACAGGTGTTACATCTGTGCTGTGAGTGCAGATGCCTCAGTTCTTGGAAGCTAGGT
TCCTGCGACTGTTACCAAGGTGATTGTAAAGAGCTGGCGGTACACAGAGGAACAAGCCCC
CAGCTGAGGGGGTGTGTGAATCGGACAGCCTCCAGCAGAGGTGTGGGAGCTGCAGCTGA
GGGAAGAAGAGACAATCGGCCTGGACACTCAGGAGGGTCAAAGGAGACTTGGTGCACAC
ACTCATCCTGCCACCCCCAGAATGCATCCTGCCTCATCAGGTCCAGATTTCTTCCAAGG
CGGACGTTTCTGTTGGAATTCTTAGTCCTTGGCCTCGGACACCTTCATTCGTTAGCTGG
GGAGTGGTGGTGAGGCAGTGAAGAAGAGGCGGATGGTCACTCAGATCCACAGAGCCCA
GGATCAAGGGACCCACTGCAGTGGCAGCAGGACTGTTGGGCCCCCAACCCCTGCAC
AGCCCTCATCCCCTCTTGGCTTGAGCCGTCAGAGGCCCTGTGCTGAGTGTCTGACCGAGA
CACTCACAGCTTTGTCTATCAGGGCACAGGCTTCCTCGGAGCCAGGATGATCTGTGCCACG
CTTGCACCTCGGGCCCATCTGGGCTCATGCTCTCTCCTGCTATTGAATTAGTACCTAG
CTGCACACAGTATGTAGTTACCAAAAGAATAAACGGCAATAATTGAGAAAAAAA

305/392

FIGURE 292

MRLGSGTFATCCVAIEVLGIAVFLRGFFPAPVRSSARAEGHGAEPPEPSAGASSNWTTL
PPPLFSKVIVLIDALRDDFVFGSKGVKFMPYTTYLVEKGASHSFVAEAKPPTVTMPRIK
ALMTGSLPGFVDVIRNLNSPALLEDVIRQAKAAGKRIVFYGDETWVKLFPHFVEYDGT
TSFFVSDYTEVDNNVTRHLDKVLKRGDWDILILHYLGLDGHIGHISGPNSPLIGQKLSEMD
SVLMKIHTSLQSKERETPLPNLLVLCGDHGMSETGSHGASSTEEVNTPLILISSAFERKP
GDIRHPKHVQ

Important features of the protein:**Signal peptide:**

amino acids 1-34

Transmembrane domain:

amino acids 58-76

N-glycosylation sites:

amino acids 56-60, 194-198

N-myristoylation sites:amino acids 6-12, 52-58, 100-106, 125-131, 233-239, 270-276,
275-281, 278-284**Amidation site:**

amino acids 154-158

Cell attachment sequence:

amino acids 205-208

306/392

FIGURE 293

AGCCAGGCAGCACATCACAGCGGGAGGAGCTGTCCCAGGTGGCCCAGCTCAGCAATGGCA
ATGGGGGTCCCCAGAGTCATTCTGCTCTGCCTCTTTGGGGCTGCGCTCTGCCTGACAGGG
TCCCAAGCCCTGCAGTGCTACAGCTTTGAGCACACCTACTTTGGCCCCTTTGACCTCAGG
GCCATGAAGCTGCCCAGCATCTCCTGTCTCATGAGTGCTTTGAGGCTATCCTGTCTCTG
GACACCGGGTATCGCGCGCCGGTGACCCTGGTGCGGAAGGGCTGCTGGACCGGGCCTCCT
GCGGGCCAGACGCAATCGAACCCGGACGCGCTGCCGCCAGACTACTCGGTGGTGCGCGGC
TGCACAACTGACAAATGCAACGCCCACCTCATGACTCATGACGCCCTCCCCAACCTGAGC
CAAGCACCCGACCCGCCGACGCTCAGCGGCGCCGAGTGCTACGCCTGTATCGGGGTCCAC
CAGGATGACTGCGCTATCGGCAGGTCCCGACGAGTCCAGTGTCACCAGGACCAGACCGCC
TGCTTCCAGGGCAGTGGCAGAATGACAGTTGGCAATTTCTCAGTCCCTGTGTACATCAGA
ACCTGCCACCGGCCCTCCTGCACCACCGAGGGCACCACCAGCCCCTGGACAGCCATCGAC
CTCCAGGGCTCCTGCTGTGAGGGGTACCTCTGCAACAGGAAATCCATGACCCAGCCCTTC
ACCAGTGCTTCAGCCACCACCCCTCCCCGAGCACTACAGGTCCTGGCCCTGCTCCTCCCA
GTCCTCCTGCTGGTGGGGCTCTCAGCATAGACCGCCCCTCCAGGATGCTGGGGACAGGGC
TCACACACCTCATTCTTGCTGCTTCAGCCCCTATCACATAGCTCACTGGAAAATGATGTT
AAAGTAAGAATTGCAAAA

307/392

FIGURE 294

MAMGVPRVILLCLFGAALCLTGSQALQCYSFEHTYFGPFDLRAMKLPSISCPHECFEAIL
SLDTGYRAPVTLVKGCWTGPPAGQTQSNPDALPPDYSVVRGCTTDKCNAHLMTHDALPN
LSQAPDPPTLSGAECYACIGVHQDDCAIGRSRRVQCHQDQTACFQSGRMTVGNFSVPVY
IRTCHRPSCCTTEGTTSPWTAIDLQSCCEGYLCNRKSMTQPFTSASATTPPRALQVLALL
LPVLLLLVGLSA

Important features of the protein:

Signal peptide:

amino acids 1-19

Transmembrane domain:

amino acids 233-251

N-glycosylation sites:

amino acids 120-124, 174-178

N-myristoylation sites:

amino acids 15-21, 84-90

308/392

FIGURE 295

AATCGGCTGATTCTGCATCTGGAACTGCCTTCATCTTGAAAGAAAAGCTCCAGGTCCCT
TCTCCAGCCACCCAGCCCCAAGATGGTGATGCTGCTGCTGCTGCTTTCCGCACTGGCTGG
CCTCTTCGGTGCGGCAGAGGGACAAGCATTTCATCTTGGAAGTGCCCCAATCCTCCGGT
GCAGGAGAATTTTGACGTGAATAAGTATCTCGGAAGATGGTACGAAATTGAGAAGATCCC
AACAACCTTTGAGAATGGACGCTGCATCCAGGCCAACTACTACTAATGGAACCGGAAA
GATCAAAGTGTTAAACCAGGAGTTGAGAGCTGATGGAAGTGAATCAAATCGAAGGTGA
AGCCACCCAGTTAACCTCACAGAGCCTGCCAAGCTGGAAGTTAAGTTTTCCTGGTTTAT
GCCATCGGCACCGTACTGGATCCTGGCCACCGACTATGAGAACTATGCCCTCGTGATTTC
CTGTACCTGCATCATCCAACTTTTTTCACGTGGATTTTGCTTGGATCTTGGCAAGAAACCC
TAATCTCCCTCCAGAAACAGTGGACTCTCTAAAAAATATCCTGACTTCTAATAACATTGA
TGTCAGAAAATGACGGTCACAGACCAGGTGAACTGCCCCAAGCTCTCGTAACCAGGTTTC
TACAGGGAGGCTGCACCCACTCCATGTTACTTCTGCTTCGCTTTCCCCTACCCACCCCC
CCCCATAAAGACAAACCAATCAACCACGACAAAGGAAGTTGACCTGAACATGTAACCAT
GCCCTACCCTGTTACCTTGCTAGCTGCAAAATAAACTTGTTGCTGACCTGCTGTGCTCGC
AAAAAA

309/392

FIGURE 296

MVMLLLLLLSALAGLFGAAEGQAFHLGKCPNPPVQENFDVNKYLGRWYEIEKIPTTFENG
RCIQANYSLMENGKIKVLNQELRADGTVNQIEGEATPVNLTEPAKLEVKFSWFMPSPAY
WILATDYENYALVYSCTCIIQLFHVDFAWILARNPNLPPETVDSLKNILTSNNIDVKKM
TVTDQVNCPKLS

Signal sequence:

1-16

N-glycosylation site:

65-68

98-101

cAMP- and cGMP-dependent protein kinase phosphorylation site:

175-178

N-myristoylation site:

13-18

16-21

Lipocalin proteins:

36-47

120-130

Lipocalin / cytosolic fatty-acid binding proteins:

41-185

310/392

FIGURE 297

GGGTGATTGAACTAAACCTTCGCCGCACCGAGTTTGCAGTACGGCCGTCACCCGCACCGC
TGCCTGCTTGCGGTTGGAGAAATCAAGGCCCTACCGGGCCTCCGTAGTCACCTCTCTATA
GTGGGCGTGGCCGAGGCCGGGGTGACCCTGCCGGAGCCTCCGCTGCCAGCGACATGTTCA
AGGTAATTCAGAGGTCCGTGGGGCCAGCCAGCCTGAGCTTGCTCACCTTCAAAGTCTATG
CAGCACCAAAAAAGGACTCACCTCCCAAAAATTCCGTGAAGGTTGATGAGCTTTCACCTCT
ACTCAGTTCCCTGAGGGTCAATCGAAGTATGTGGAGGAGGCAAGGAGCCAGCTTGAAGAAA
GCATCTCACAGCTCCGACACTATTGCGAGCCATACACAACCTGGTGTCAGGAAACGTACT
CCCAAACCTAAGCCCAAGATGCAAAGTTTGGTTCAATGGGGGTTAGACAGCTATGACTATC
TCCAAAATGCACCTCCTGGATTTTTTCCGAGACTTGGTGTTATTGGTTTTGCTGGCCTTA
TTGGACTCCTTTTGGCTAGAGGTTCAAAAATAAAGAAGCTAGTGTATCCGCCTGGTTTTCA
TGGGATTAGCTGCCTCCCTCTATTATCCACAACAAGCCATCGTGTTTGCCAGGTCAGTG
GGGAGAGATTATATGACTGGGGTTTACGAGGATATATAGTCATAGAAGATTTGTGGAAGG
AGAACTTTCAAAAGCCAGGAAATGTGAAGAATTACCTGGAACCTAAGTAGAAAACTCCAT
GCTCTGCCATCTTAATCAGTTATAGGTAAACATTGGAACTCCATAGAATAAAATCAGTAT
TTCTACAGAAAAATGGCATAGAAGTCAGTATTGAATGTATTAAATTGGCTTTCTTCTTCA
GGAAAACTAGACCAGACCTCTGTTATCTTCTGTGAAATCATCCTACAAGCAAACTAACC
TGGAATCCCTTCACCTAGAGATAATGTACAAGCCTTAGAACTCCTCATTTCTCATGTTGCT
ATTTATGTACCTAATTAAAACCCAAGTTTAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAA

311/392

FIGURE 298

MFKVIQRSVGPASLSLLTFKVYAAPKSDSPPKNSVKVDELSTLYSVPEGQSKYVEEARSQ
EESISQLRHYCEPYTTWCQETYSQTKPKMQSLVQWGLDSYDYLQNAPPGFFPRLGVIGFA
GLIGLLLARGSKIKKLVYPPGFMGLAASLYYPQQAIVFAQVSGERLYDWGLRGYIVIEDL
WKENFQKPGNVKNSPGTK

Important features:

Signal peptide:

Amino acids 1-23

Transmembrane domain:

Amino acids 111-130

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 26-30

Tyrosine kinase phosphorylation site:

Amino acids 36-44

N-myristoylation sites:

Amino acids 124-130;144-150;189-195

312/392

FIGURE 299

CCGCTGAGATGTACGAACTTCCGGTTCTCCGGGCAGCTGCCACTGCTGTAGCTTCTGCCA
CCTGCCACGACCGGGCCTCTCCCTGGCGTTTGGTCACCTCTGCTTCATTCTCCACCGCGC
CTATGGTCCCTCTTGGAGCCAGCGTGGCGGGCCTGGCGGGCTCCCGGGTGGTGAGAGAGCG
GTCCGGGAACGATGAAGGCCTCGCAGTGCTGCTGCTGTCTCAGCCACCTCTTGGCTTCCG
TCCTCCTCCTGCTGTTGCTGCCTGAACTAAGCGGGCCCCCTGGCAGTCCTGCTGCAGGCAG
CCGAGGCCGCGCCAGGTCTTGGGCCTCCTGACCCTAGACCACGGACATTACCGCCGCTGC
CACCGGGCCCTACCCCTGCCCAGCAGCCGGGCGGTGGTCTGGCTGAAGCTGCGGGGCCGC
GGGGCTCCGAGGGAGGCAATGGCAGCAACCCTGTGGCCGGGCTTGAGACGGACGATCACG
GAGGGAAGGCCGGGGAAGGCTCGGTGGGTGGCGGCCTTGCTGTGAGCCCCAACCTGGCG
ACAAGCCCATGACCCAGCGGGCCCTGACCGTGTGATGGTGGTGAGCGGCGCGGTGCTGG
TGTACTTCGTGGTCAGGACGGTCAGGATGAGAAGAAGAAACCGAAAGACTAGGAGATATG
GAGTTTTGGACACTAACATAGAAAATATGGAATTGACACCTTTAGAACAGGATGATGAGG
ATGATGACAACACGTTGTTTGATGCCAATCATCCTCGAAGATAAGAATGTGCCTTTTGAT
GAAAGAACTTTATCTTTCTACAATGAAGAGTGGAATTTCTATGTTTAAGGAATAAGAAGC
CACTATATCAATGTTGGGGGGGTATTTAAGTTACATATATTTTAACAACCTTTAATTTGC
TGTTGCAATAAATAACCGTATCCTTTTTATTATATCTTTATATGTATAGAAGTACTCTATTA
ATGGGCTCAGAGATGTTGGGGATAAAGTATACTGTAATAATTTATCTGTTTGAAAATTAC
TATAAAACGGTGTTTTCTGGTCCGTTTTTGTTCCTGCTTACCATATGATTGTAAATTGT
TTTATGTATTAATCAGTTAATGCTAATTATTTTTGCTGATGTCATATGTTAAAGAGCTAT
AAATTCCAACAACCAACTGGTGTGTAAAAATAATTTAAATTTCTTTACTGAAAGGTAT
TTCCCATTTTTGTGGGGAAAAGAAGCCAAATTTATTACTTTGTGTTGGGGTTTTTAAAT
ATTAAGAAATGTCTAAGTTATTGTTTGCAAAACAATAAATATGATTTTAAATTCCTTAA
AAAAAAA

313/392

FIGURE 300

MKASQCCCCLSHLLASVLLLLLLPELSGPLAVLLQAAEAAPGLGPPDPRPRTLPPPLPPGP
TPAQQPGRGLAEAAGPRGSEGGNGSNPVAGLETDDHGGKAGEGSVGGGLAVSPNPGDKPM
TQRALTVLMVVSGAVLVYFVVRTVRMRRNRKTRRYGVLDTNIE NMELTPLEQDDEDDDN
TLFDANHPRR

Signal peptide:
amino acids 1-28

Transmembrane domain:
amino acids 124-140

N-glycosylation site:
amino acids 83-87

N-myristoylation sites:
amino acids 69-75, 78-84, 81-87, 97-103, 103-109, 106-112,
157-160

314/392

FIGURE 301

CTCGGCTGGATTTAAGGTTGCCGCTAGCCGCTGGGAATTTAAGGGACCCACACTACCTT
CCCGAAGTTGAAGGCAAGCGGTGATTGTTTGTAGACGGCGCTTTGTCATGGGACCTGTGC
GGTTGGGAATATTGCTTTTCCTTTTGGCCGTGCACGAGGCTTGGGCTGGGATGTTGA
AGGAGGAGGACGATGACACAGAACGCTTGCCCAGCAAATGCGAAGTGTGTAAGCTGCTGA
GCACAGAGCTACAGGCGGAAGTGAAGTGCACCGGTGATCTCGAGAGGTGCTGGAGCTGG
GGCAGGTGCTGGATACAGGCAAGAGGAAGAGACACGTGCCTTACAGCGTTTCAGAGACAA
GGCTGGAAGAGGCTTAGAGAATTTATGTGAGCGGATCCTGGACTATAGTGTTACGCTG
AGCGCAAGGGCTCACTGAGATATGCCAAGGGTCAGAGTCAGACCATGGCAACACTGAAAG
GCCTAGTGCAGAAGGGGGTGAAGGTGGATCTGGGGATCCCTCTGGAGCTTTGGGATGAGC
CCAGCGTGGAGGTCACATACCTCAAGAAGCAGTGTGAGACCATGTTGGAGGAGTTTGAAG
ACATTGTGGGAGACTGGTACTTCCACCATCAGGAGCAGCCCCACAAAATTTCTCTGTG
AAGGTCATGTGCTCCAGCTGCTGAAACTGCATGTCTACAGGAACTTGGACTGGAAAGG
AGATCACAGATGGGGAAGAGAAAACAGAAGGGGAGGAAGAGCAGGAGGAGGAGGAGGAAG
AGGAGGAAGAGGAAGGGGGAGACAAGATGACCAAGACAGGAAGCCACCCAACTTGACC
GAGAAGATCTTTTGACCCTTGCCTTTGAGCCCCCAGGAGGGGAAGGGATCATGGAGAGCCC
TCTAAAGCCTGCACTCTCCCTGCTCCACAGCTTTCAGGGTGTGTTTATGAGTGACTCCAC
CCAAGCTTGTAGCTGTTCTCTCCCATCTAACCTCAGGCAAGATCCTGGTGAAACAGCATG
ACATGGCTTCTGGGGTGGAGGGTGGGGGTGGAGGTCCTGCTCCTAGAGATGAACTCTATC
CAGCCCCCTTAATTGGCAGGTGTATGTGCTGACAGTACTGAAAGCTTTCCTCTTTAACTGA
TCCCACCCCCACCCAAAAGTCAGCAGTGGCACTGGAGCTGTGGGCTTTGGGGAAGTCACT
TAGCTCCTTAAGGTCTGTTTTTAGACCCTTCCAAGGAAGAGGCCAGAACGGACATTCTCT
GCGATCTATATACATTGCCTGTATCCAGGAGGCTACACACCAGCAAACCGTGAAGGAGAA
TGGGACACTGGGTCATGGCCTGGAGTTGCTGATAATTTAGGTGGGATAGATACTTGGTCT
ACTTAAGCTCAATGTAACCCAGAGCCCACCATATAGTTTTATAGGTGCTCAACTTTCTAT
ATCGCTATTAACTTTTTTCTTTTTTTCTA

315/392

FIGURE 302

MGPVRLGILLFLFLAVHEAWAGMLKEEDDDTERLPSKCEVCKLLSTELQAELSRTGRSRE
VLELGQVLDTGKRKRHPYVSSETRLEEALENLCERILDYSVHAERKGSRLRYAKGQSQTM
ATLKGLVQKGVKVDLGIPLELWDEPSVEVTYLKKQCETMLEEFEDIVGDWYFHHQEQLQ
NFLCEGHVLPAAETACLQETWTGKEITDGEEKTEGEEEQEEEEEEEEEGGDKMTKTGSH
PKLDREDL

Important features of the protein:

Signal peptide:

amino acids 1-21

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 106-110

N-myristoylation site:

amino acids 115-121

Amidation site:

amino acids 70-74

316/392

FIGURE 303

CTCCTGCACTAGGCTCTCAGCCAGGGATGATGCGCTGCTGCCGCCGCCGCTGCTGCTGCC
GGCAACCACCCCATGCCCTGAGGCCGTTGCTGTTGCTGCCCCCTCGTCCTTTTACCTCCCC
TGGCAGCAGCTGCAGCGGGCCCAAACCGATGTGACACCATATAACCAGGGCTTCGCCGAGT
GTCTCATCCGCTTGGGGGACAGCATGGGCCGCGGAGGCGAGCTGGAGACCATCTGCAGGT
CTTGGAATGACTTCCATGCCTGTGCCTCTCAGGTCTGTGTCAGGCTGTCCGGAGGAGGCAG
CTGCAGTGTGGGAATCACTACAGCAAGAAGCTCGCCAGGCCCCCGTCCGAATAACTTGC
ACACTCTGTGCGGTGCCCCGGTGCATGTTCTGGGAGCGCGGCACAGGCTCCGAAACCAACC
AGGAGACGCTGCGGGCTACAGCGCCTGCACTCCCCATGGCCCCCTGCGCCCCCACTGCTGG
CGGCTGCTCTGGCTCTGGCCTACCTCCTGAGGCCTCTGGCCTAGCTTGTTGGGTTGGGTA
GCAGCGCCCGTACCTCCAGCCCTGCTCTGGCGGTGGTTGTCCAGGCTCTGCAGAGCGCAG
CAGGGCTTTTCATTAAAGGTATTTATATTTGTA

317/392

FIGURE 304

MMRCCRRRCCCRQPPHALRPLLLLPLVLLPPLAAAAAGPNRCDTIYQGFAECLIRLGDSM
GRGGELETICRSWNDFHACASQVLSGCPPEEAAVWESLQQEARQAPRPNNLHTLCGAPVH
VRERGTGSETNQETLRATAPALPMAPAPPLLAALALAYLLRPLA

Signal peptide:

Amino acids 1-35

Transmembrane domain:

Amino acids 141-157

N-myristoylation site:

Amino acids 127-133

Prokaryotic membrane lipoprotein lipid attachment site:

Amino acids 77-88

318/392

FIGURE 305

AAGTACTTGTGTCCGGGTGGTGGACTGGATTAGCTGCGGAGCCCTGGAAGCTGCCTGTCC
TTCTCCCTGTGCTTAACCAGAGGTGCCCATGGGTGGACAATGAGGCTGGTCACAGCAGC
ACTGTTACTGGGTCTCATGATGGTGGTCACTGGAGACGAGGATGAGAACAGCCCGTGTGC
CCATGAGGCCCTCTTGGACGAGGACACCCTCTTTTGCCAGGGCCTTGAAGTTTTCTACCC
AGAGTTGGGGAACATTGGCTGCAAGGTTGTTCCCTGATTGTAACAACCTACAGACAGAAGAT
CACCTCCTGGATGGAGCCGATAGTCAAGTTCCCGGGGGCCGTGGACGGCGCAACCTATAT
CCTGGTGATGGTGGATCCAGATGCCCCTAGCAGAGCAGAACCAGACAGAGATTCTGGAG
ACATTGGCTGGTAACAGATATCAAGGGCGCCGACCTGAAGAAAGGGAAGATTCAAGGGCCA
GGAGTTATCAGCCTACCAGGCTCCCTCCCCACCGGCACACAGTGGCTTCCATCGCTACCA
GTTCTTTGTCTATCTTCAGGAAGGAAAAGTCATCTCTCTCCTTCCCAAGGAAAACAAAAC
TCGAGGCTCTTGAAAAATGGACAGATTTCTGAACCGCTTCCACCTGGGCGAACCTGAAGC
AAGCACCCAGTTCATGACCCAGAACTACCAGGACTCACCAACCCTCCAGGCTCCAGAGG
AAGGGCCAGCGAGCCCAAGCACAAAACCAGGCAGAGATAGCTGCCTGCTAGATAGCCGGC
TTTGCCATCCGGGCATGTGGCCCACTGCTCACCACCGACGATGTGGGTATGGAACCCCC
TCTGGATACAGAACCCTTCTTTTCCAAATTAAAAAAAAAAATCATCAA

319/392

FIGURE 306

MGWTMRLVTAALLLGLMMVVTGDEDENSPCAHEALLDEDTLFCQGLEVFYPELGNI GCKV
VPDCNNYRQKITSWMEPIVKFPGAVDGATYILVMVDPDAPSRAEPRQRFWRHWLVTDIKG
ADLKKGKIQGQELSAYQAPSPPAHSGFHR YQFFVYLQEGKVISLLPKENKTRGSWKMDRF
LNR FHLGEPEASTQFMTQNYQDSPTLQAPRGRASEPKHKTRQR

Important features of the protein:

Signal peptide:

amino acids 1-22

N-glycosylation site:

amino acids 169-173

Tyrosine kinase phosphorylation site:

amino acids 59-68

N-myristoylation sites:

amino acids 54-60, 83-89, 130-136

Phosphatidylethanolamine signature:

amino acids 113-157

320/392

FIGURE 307

AAGGAGCAGCCCGCAAGCACCAAGTGAGAGGCATGAAGTTACAGTGTGTTTCCCTTTGGC
TCCTGGGTACAATACTGATATTGTGCTCAGTAGACAACCACGGTCTCAGGAGATGTCTGA
TTTCCACAGACATGCACCATATAGAAGAGAGTTTCCAAGAAATCAAAAGAGCCATCCAAG
CTAAGGACACCTTCCCAAATGTCACTATCCTGTCCACATTGGAGACTCTGCAGATCATTA
AGCCCTTAGATGTGTGCTGCGTGACCAAGAACCTCCTGGCGTTCTACGTGGACAGGGTGT
TCAAGGATCATCAGGAGCCAAACCCAAAATCTTGAGAAAAATCAGCAGCATTGCCAACT
CTTTCCTCTACATGCAGAAAACCTCTGCGGCAATGTCAGGAACAGAGGCAGTGTCACTGCA
GGCAGGAAGCCACCAATGCCACCAGAGTCATCCATGACAACCTATGATCAGCTGGAGGTCC
ACGCTGCTGCCATTAAATCCCTGGGAGAGCTCGACGTCTTCTAGCCTGGATTAATAAGA
ATCATGAAGTAATGTTCTCAGCTTGATGACAAGGAACCTGTATAGTGATCCAGGGATGAA
CACCCCCTGTGCGGTTTACTGTGGGAGACAGCCCACCTTGAAGGGGAAGGAGATGGGGAA
GGCCCCCTTGACAGCTGAAAGTCCCACTGGCTGGCCTCAGGCTGTCTTATTCCGCTTGAAAA
TAGGCAAAAAGTCTACTGTGGTATTTGTAATAAACTCTATCTGCTGAAAGGGCCTGCAGG
CCATCCTGGGAGTAAAGGGCTGCCTTCCCATCTAATTTATTGTAAAGTCATATAGTCCAT
GTCTGTGATGTGAGCCAAGTGATATCCTGTAGTACACATTGTACTGAGTGGTTTTTCTGA
ATAAATTCATATTTTACCTATGA

321/392

FIGURE 308

MKLQCVSLWLLGTILILCSVDNHGLRRCLISTDMHHIEESFQEIKRAIQAKDTFPNVTIL
STLETLQIIKPLDVCCVTKNLLAFYVDRVFKDHQEPNPKILRKISSIANSFLYMQKTLRQ
CQEQRQCHCRQEATNATRVIHNDYDQLEVHAAAIKSLGELDVFLAWINKNHEVMFSA

Signal sequence:
amino acids 1-18

N-glycosylation sites:
amino acids 56-60, 135-139

cAMP- and cGMP-dependent protein kinase phosphorylation site:
amino acids 102-106

N-myristoylation site:
amino acids 24-30

Actinin-type actin-binding domain signature 1:
amino acids 159-169

322/392

FIGURE 309

GTCGACCCACGCGTCCGAAGCTGCTGGAGCCACGATTTCAGTCCCCTGGACTGTAGATAAA
GACCCTTTCCTTGCCAGGTGCTGAGACAACCACACTATGAGAGGCACTCCAGGAGACGCTG
ATGGTGGAGGAAGGGCCGCTCTATCAATCAATCACTGTTGCTGTTATCACATGCAAGTATC
CAGAGGCTCTTGAGCAAGGCAGAGGGGATCCCATTTATTTGGGAATCCAGAATCCAGAAA
TGTGTTTGTATTGTGAGAAGGTTGGAGAACAGCCACATTGCAGCTAAAAGAGCAGAAGA
TCATGGATCTGTATGGCCAACCCGAGCCCGTGAAACCCTTCCTTTTCTACCGTGCCAAGA
CTGGTAGGACCTCCACCCTTGAGTCTGTGGCCTTCCCGGACTGGTTTCATTGCCTCCTCCA
AGAGAGACCAGCCCATCATTCTGACTTCAGAACTTGGGAAGTCATACAACACTGCCTTTG
AATTAAATATAAATGACTGAACTCAGCCTAGAGGTGGCAGCTTGGTCTTTGCTTTAAAGT
TTCTGGTTCCCAATGTGTTTTTCGTCTACATTTTCTTAGTGTCAATTTTCACGCTGGTGCTG
AGACAGGAGCAAGGCTGCTGTTATCATCTCATTTTATAATGAAGAAGAAGCAATTACTTC
ATAGCAACTGAAGAACAGGATGTGGCCTCAGAAGCAGGAGAGCTGGGTGGTATAAGGCTG
TCCTCTCAAGCTGGTGCTGTGTAGGCCACAAGGCATCTGCATGAGTGACTTTAAGACTCA
AAGACCAAACACTGAGCTTTCTTCTAGGGGTGGGTATGAAGATGCTTCAGAGCTCATGCG
CGTTACCCACGATGGCATGACTAGCACAGAGCTGATCTCTGTTTTCTGTTTTGCTTTATTC
CCTCTTGGGATGATATCATCCAGTCTTTATATGTTGCCAATATACCTCATTGTGTGTAAT
AGAACCCTTCTTAGCATTAAGACCTTGTAACAAAAATAATTCTTGGGGTGGGTATGAAGA
TGCTTCAGAGCTCATGCGGTTACCCACGATGGCATGACTAGCACAGAGCTGATCTCTGT
TTCTGTTTTGCTTTATTCCCTCTTGGGATGATATCATCCAGTCTTTATATGTTGCCAATA
TACCTCATTGTGTGTAATAGAACCCTTCTTAGCATTAAGACCTTGTAACAAAAATAATTC
TTGTGTTAAGTTAAATCATTTTTGTCCTAATTGTAATGTGTAATCTTAAAGTTAAATAAA
CTTTGTGTATTTATATAATAATAAAGCTAAACTGATATAAAATAAAGAAAGAGTAAACTG

323/392

FIGURE 310

MRGTPGDADGGGRAVYQSITVAVITCKYPEALEQGRGDPIYLGIONPEMCLYCEKVGEQP
TLQLKEQKIMDLYGQPEPVKPFLFYRAKTGRTSTLESVAFPDWFIASSKRDQPIILTSEL
GKSYNTAFELNIND

Signal sequence:
amino acids 1-17

N-myristoylation site:
amino acids 10-16

Cell attachment sequence:
amino acids 36-39

324/392

FIGURE 311

GCGAGGCTGCACCAGCGCCTGGCACCATGAGGACGCCTGGGCCTCTGCCCCGTGCTGCTGC
TGCTCCTGGCGGGAGCCCCGCGCGCGGCCACTCCCCGACCTGCTACTCCCGCATGC
GGGCCCTGAGCCAGGAGATCACCCGCGACTTCAACCTCCTGCAGGTCTCGGAGCCCTCGG
AGCCATGTGTGAGATACCTGCCCAGGCTGTACCTGGACATACACAATTACTGTGTGCTGG
ACAAGCTGCGGGACTTTGTGGCCTCGCCCCCGTGTTGGAAAGTGGCCCAGGTAGATTCTT
TGAAGGACAAAGCACGGAAGCTGTACACCATCATGAACTCGTTCTGCAGGAGAGATTTGG
TATTCCTGTTGGATGACTGCAATGCCTTGGAATACCCAATCCCAGTGACTACGGTCCTGC
CAGATCGTCAGCGCTTAAGGGAAGTGAAGACCAGAGAAAGAACCCAAGAGAACTAAAGTTAT
GTCAGCTACCCAGACTTAATGGGCCAGAGCCATGACCCTCACAGGTCTTGTTAGTTGT
ATCTGAAACTGTTATGTATCTCTACCTTCTGGAAAACAGGGCTGGTATTCCTACCCAG
GAACCTCCTTTGAGCATAGAGTTAGCAACCATGCTTCTCATTCCCTTGACTCATGTCTTG
CCAGGATGGTTAGATACACAGCATGTTGATTGGTCACTAAAAAGAAGAAAAGGACTAAC
AAGCTTCACTTTTATGAACAACTATTTTGAGAACATGCACAATAGTATGTTTTTATTACT
GGTTTAATGGAGTAATGGTACTTTTATTCTTTCTTGATAGAAACCTGCTTACATTTAACC
AAGCTTCTATTATGCCTTTTTCTAACACAGACTTTCTTCACTGTCTTTCAATTTAAAAAGA
AATTAATGCTCTTAAGATATATATTTTACGTAGTGCTGACAGGACCCACTCTTTCATTGA
AAGGTGATGAAAATCAAATAAAGAATCTCTTCACATGGA

325/392

FIGURE 312

MRTPGPLPVLLLLLAGAPAARPTPPTCYSRMRALSQEITRDFNLLQVSEPSEPCVRYLPR
LYLDIHNYCVLDKLRDFVASPPCWKVAQVDSLKDKARKLYTIMNSFCRRDLVFLDDCNA
LEYPIPVTTVLPRQR

Important features of the protein:

Signal peptide:

amino acids 1-19

Tyrosine kinase phosphorylation site:

amino acids 60-69

N-myristoylation site:

amino acids 16-22

326/392

FIGURE 313

GAGCGACGCTGTCTCTAGTCGCTGATCCCAAATGCACCGGCTCATCTTTGTCTACACTCT
AATCTGCGCAAACCTTTTGCAGCTGTCGGGACACTTCTGCAACCCCGCAGAGCGCATCCAT
CAAAGCTTTGCGCAACGCCAACCTCAGGCGAGATGACTTGTACCGAAGAGATGAGACCAT
CCAGGTGAAAGGAAACGGCTACGTGCAGAGTCCTAGATTCCCGAACAGCTACCCAGGAA
CCTGCTCCTGACATGGCGGCTTCACTCTCAGGAGAATACACGGATACAGCTAGTGTGTTGA
CAATCAGTTTGGATTAGAGGAAGCAGAAAATGATATCTGTAGGTATGATTTTGTGGAAGT
TGAAGATATATCCGAAACCAGTACCATTATTAGAGGACGATGGTGTGGACACAAGGAAGT
TCCTCCAAGGATAAAATCAAGAACGAACCAAATTAAAATCACATTCAAGTCCGATGACTA
CTTTGTGGCTAAACCTGGATTCAAGATTTATTATTCTTTGCTGGAAGATTCCAACCCGC
AGCAGCTTCAGAGACCAACTGGGAATCTGTACAAGCTCTATTTCAAGGGTATCCTATAA
CTCTCCATCAGTAACGGATCCCACTCTGATTGCGGATGCTCTGGACAAAAAATTGCAGA
ATTTGATACAGTGGAAGATCTGCTCAAGTACTTCAATCCAGAGTCATGGCAAGAAGATCT
TGAGAATATGTATCTGGACACCCCTCGGTATCGAGGCAGGTCATACCATGACCGGAAGTC
AAAAGTTGACCTGGATAGGCTCAATGATGATGCCAAGCGTTACAGTTGCACTCCAGGAA
TTACTCGGTCAATATAAGAGAAGAGCTGAAGTTGGCCAATGTGGTCTTCTTTCCACGTTG
CCTCCTCGTGCAGCGCTGTGGAGGAAATTGTGGCTGTGGAAGTGTCAACTGGAGGTCCTG
CACATGCAATTCAAGGAAAACCGTGAAAAAGTATCATGAGGTATTACAGTTTGAGCCTGG
CCACATCAAGAGGAGGGGTAGAGCTAAGACCATGGCTCTAGTTGACATCCAGTTGGATCA
CCATGAACGATGCGATTGTATCTGCAGCTCAAGACCACCTCGATTAAGAGAATGTGCACAT
CCTTACATTAAGCCTGAGAGAA

327/392

FIGURE 314

MHRLIFVYTLICANFCSCRDTSATPQSASIKALRNANLRRDDLYRRDETIQVKGNGYVQS
PRFPNSYPRNLLLTWRLHSQENTRIQLVFDNQFGLLEEAENDICRYDFVEVEDISETSTII
RGRWCGHKEVPPRIKSRTNQIKITFKSDDYFVAKPGFKIYYSLLEDFQPAAASETNWESV
TSSISGVSYNPSVTDPTLIADALDKKIAEFDTVEDLLKYFNPESWQEDLENMYLDTPRY
RGRSYHDRKSKVDLRLNDDAKRYSCTPRNYSVNIREELKLANVVFPRCLLVQRCGGNC
GCGTVNWRSCTCNSGKTVKKYHEVLQFEPGHIKRRGRAKTMALVDIQLDHHHERCDCICSS
RPPR

Signal peptide:

amino acids 1-18

N-glycosylation site:

amino acids 270-274

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 262-266

Tyrosine kinase phosphorylation site:

amino acids 256-265

N-myristoylation sites:

amino acids 94-100, 186-192, 297-303, 298-304

TonB-dependent receptor proteins signature 1:

amino acids 1-56

328/392

FIGURE 315

CGGCTCGAGGCTCCCGCCAGGAGAAAGGAACATTCTGAGGGGAGTCTACACCCTGTGGAG
CTCAAGATGGTCCTGAGTGGGGCGCTGTGCTTCCGAATGAAGGACTCGGCATTGAAGGTG
CTTTATCTGCATAATAACCAGCTTCTAGCTGGAGGGCTGCATGCAGGGAAGGTCATTAAA
GGTGAAGAGATCAGCGTGGTCCCAATCGGTGGCTGGATGCCAGCCTGTCCCCCGTCATC
CTGGGTGTCCAGGGTGAAGCCAGTGCCTGTCTGTGGGGTGGGGCAGGAGCCGACTCTA
ACACTAGAGCCAGTGAACATCATGGAGCTCTATCTTGGTGCCAAGGAATCCAAGAGCTTC
ACCTTCTACCGGCGGGACATGGGGCTCACCTCCAGCTTCGAGTCGGCTGCCTACCCGGGC
TGGTTCCTGTGCACGGTGCCTGAAGCCGATCAGCCTGTCTCAGACTCACCCAGCTTCCCGAG
AATGGTGGCTGGAATGCCCCCATCACAGACTTCTACTTCCAGCAGTGTGACTAGGGCAAC
GTGCCCCCAGAACTCCCTGGGCAGAGCCAGCTCGGGTGAGGGGTGAGTGGAGGAGACCC
ATGGCGGACAATCACTCTCTCTGCTCTCAGGACCCCCACGTCTGACTTAGTGGGCACCTG
ACCACTTTGTCTTCTGGTTCCAGTTTGGATAAAATTCTGAGATTTGGAGCTCAGTCCACG
GTCCTCCCCCACTGGATGGTGCTACTGCTGTGGAACCTTGTA AAAACCATGTGGGGTAAA
CTGGGAATAACATGAAAAGATTTCTGTGGGGGTGGGGTGGGGGAGTGGTGGGAATCATTC
CTGCTTAATGGTAACTGACAAGTGTTACCCTGAGCCCCGCAGGCCAACCCATCCCCAGTT
GAGCCTTATAGGGTCACTAGCTCTCCACATGAAGTCCTGTCACTCACCCTGTGCAGGAG
AGGGAGGTGGTCATAGAGTCAGGGATCTATGGCCCTTGGCCCAGCCCCACCCCTTCCCT
TTAATCCTGCCACTGTCTATATGCTACCTTTCCTATCTCTTCCCTCATCATCTTGTGTGG
GCATGAGGAGGTGGTGATGTCTAGAAGAAATGGCTCGAGCTCAGAAGATAAAAGATAAGTA
GGGTATGCTGATCCTCTTTTAAAAACCCAAGATACAATCAAATCCCAGATGCTGGTCTC
TATTCCTCATGAAAAGTGCTCATGACATATTGAGAAGACCTACTTACAAAGTGGCATATA
TTGCAATTTATTTTAAATTAAGATACCTATTTATATATTTCTTTATAGAAAAAAGTCTG
GAAGAGTTTACTTCAATTGTAGCAATGTCTAGGGTGGTGGCAGTATAGGTGATTTTCTTT
TAATTCCTGTAAATTTATCTGTATTTCTTAATTTTCTACAATGAAGATGAATTCCTGTGA
TAAAAATAAGAAAAGAAATTAATCTTGAGGTAAGCAGAGCAGACATCATCTCTGATTGTC
CTCAGCCTCCACTTCCCCAGAGTAAATTCAAATTGAATCGAGCTCTGCTGCTCTGGTTGG
TTGTAGTAGTGATCAGGAAACAGATCTCAGCAAAGCCACTGAGGAGGAGGCTGTGCTGAG
TTTGTGTGGCTGGAATCTCTGGGTAAGGAACCTTAAAGAACAAAAATCATCTGGTAATTCT
TTCCTAGAAGGATCACAGCCCCCTGGGATTCCAAGGCATTGGATCCAGTCTCTAAGAAGGC
TGCTGTACTGGTTGAATTGTGTCCCCCTCAAATTCACATCCTTCTTGGAATCTCAGTCTG
TGAGTTTATTTGGAGATAAGGTCTCTGCAGATGTAGTTAGTTAAGACAAGGTCATGCTGG
ATGAAGGTAGACCTAAATTCATATGACTGGTTTCTTGTATGAAAAGGAGAGGACACAG
AGACAGAGGAGACGCGGGGAAGACTATGTAAAGATGAAGGCAGAGATCGGAGTTTTCAG
CCACAAGCTAAGAAACACCAAGGATTGTGGCAACCATCAGAAGCTTGGAAGAGGCAAGA
AGAATTCTTCCCTAGAGGCTTTAGAGGGATAACGGCTCTGCTGAAACCTTAATCTCAGAC
TTCCAGCCTCCTGAACGAAGAAAGATAAAATTTCCGGCTGTTTTAAAGCCACCAAGGATAAT
TGGTTACAGCAGCTCTAGGAAACTAATACAGCTGCTAAAATGATCCCTGTCTCCTCGTGT
TTACATTCTGTGTGTGTCCCCCTCCACAATGTACCAAAGTTGTCTTTGTGACCAATAGAA
TATGGCAGAAGTGATGGCATGCCACTTCCAAGATTAGGTTATAAAAGACACTGCAGCTTC
TACTTGAGCCCTCTCTCTCTGCCACCCACCGCCCCCAATCTATCTTGGCTCACTCGCTCT
GGGGGAAGCTAGCTGCCATGCTATGAGCAGGCCTATAAAGAGACTTACGTGGTAAAAAAT
GAAGTCTCCTGCCACAGCCACATTAGTGAACCTAGAAGCAGAGACTCTGTGAGATAATC
GATGTTTGTGTTGTTTAAAGTTGCTCAGTTTTGGTCTAACTTGTTATGCAGCAATAGATAAA
TAATATGCAGAGAAAGAG

329/392

FIGURE 316

MVLSGALCFRMKDSALKVLYLHNNQLLAGGLHAGKVIKGEESVVPNRWLDASLSPVILG
VQGGSQCLSCGVGQEPTLTLEPVNIMELYLGAKESKSFTFYRRDMGLTSSFESAAYPGWF
LCTVPEADQPVRLTQLPENGGWNAPIITDFYFQQCD

N-myristoylation sites:

amino acids 29-34, 30-35, 60-65, 63-68, 73-78, 91-96, 106-111

Interleukin-1 signature:

amino acids 111-131

Interleukin-1 proteins:

amino acids 8-29, 83-120, 95-134, 64-103

330/392

FIGURE 317

ATGGAACTTGGACTTGGAGGCCTCTCCACGCTGTCCCACTGCCCCTGGCCTAGGCGGCAG
CCTGCCCCTGTGGCCCACCCTGGCCGCTCTGGCTCTGCTGAGCAGCGTCGCAGAGGCCTCC
CTGGGCTCCGCGCCCCGCAGCCCTGCCCCCGCGAAGGCCCCCGCCTGTCCTGGCGTCC
CCCGCCGGCCACCTGCCGGGGGGACGCACGGCCCGCTGGTGCAAGAGAGCCCGGCGG
CCGCGCCCGCAGCCTTCTCGGCCCCGCGCCCCCGCCGCTGCACCCCCATCTGCTCTTCCC
CGCGGGGGCCGCGCGGCGCGGGCTGGGGGCCCGGGCAGCCGCGCTCGGGCAGCGGGGGCG
CGGGGCTGCCGCTGCGCTCGCAGCTGGTGCCGGTGCGCGCGCTCGGCCTGGGCCACCGC
TCCGACGAGCTGGTGCGTTTCCGCTTCTGCAGCGGCTCCTGCCGCCGCGCGCGCTCTCCA
CACGACCTCAGCCTGGCCAGCCTACTGGGCGCCGGGGCCCTGCGACCGCCCCCGGGCTCC
CGGCCCCTCAGCCAGCCCTGCTGCCGACCCACGCGCTACGAAGCGGTCTCCTTCATGGAC
GTCAACAGCACCTGGAGAACCGTGGACCGCCTCTCCGCCACCGCCTGCGGCTGCCTGGGC
TGA

331/392

FIGURE 318

MELGLGGLSTLSHCPWPRRQPALWPTLAALALLSSVAEASLGSA PRSPAPREGPPPVLAS
PAGHLPGGRTARWC SGRRARRPPPQPSRPAPPPPAPPSALPRGGRAARAGGP GSRARAAGA
RGCRLRSQ LVPVRALGLGHRSD ELVRFRFCSGSCRRARSPHDLSLASLLGAGALRPPPGS
RPVSQPCCRPTRYEAVSFMDVNSTWRTVDRLSATA CGCLG

signal sequence:

Amino acids 1-39

N-glycosylation site:

Amino acids 202-206

N-myristoylation sites:

Amino acids 6-12;67-73;102-108;109-115;119-125

332/392

FIGURE 319

GTTGCTATGTTGCCCAGGCTGGTCTTGAAGTGCCTTGACCTCCTAAAGTGTTGGAACCAC
AGACGTGAGCCACTCCACCCAGCCTAAACTTCATCTTCTTTGGATGAGATGAACACTTT
TAACAAGAGAACAGGACTCTATATAAATCGCTGTGGGCTCACCACCTCTAAGGAGGAGCA
CTGACTGAAGACAGAAAAATTGATGAACTGAAGAAGACATGGTCCATTATGCCTTACAAA
CTTACACAGTGCTTTGGGAATTCCAAAGTACTCAGTGGAGAGAGGTGTTTCAGGAGCCGT
AGAGCCAGATCGTCATCATGTCTGCATTGTGGCTGCTGCTGGGCCTCCTTGCCCTGATGG
ACTTGTCTGAAAGCAGCAACTGGGGATGCTATGGAAACATCCAAAGCCTGGACACCCCTG
GAGCATCTTGTGGGATTGGAAGACGTCACGGCCTGAACTACTGTGGAGTTCGTGCTTCTG
AAAGGCTGGCTGAAATAGACATGCCATACCTCCTGAAATATCAACCCATGATGCAAACCA
TTGGCCAAAAGTACTGCATGGATCCTGCCGTGATCGCTGGTGTCTTGTCCAGGAAGTCTC
CCGGTGACAAAATTCTGGTCAACATGGGCGATAGGACTAGCATGGTGCAGGACCCTGGCT
CTCAAGCTCCACATCCTGGATTAGTGAGTCTCAGGTTTCCCAGACAACTGAAGTTCTGA
CTACTAGAATCAAAGAAATCCAGAGGAGGTTTCCAACCTGGACCCCTGACCAGTACCTGA
GAGGTGGACTCTGTGCCTACAGTGGGGGTGCTGGCTATGTCCGAAGCAGCCAGGACCTGA
GCTGTGACTTCTGCAATGATGTCTTGCACGAGCCAAGTACCTCAAGAGACATGGCTTCT
AACATCTCAGATGAAACCCAAGACCATGATCACATATGCAGCCTCAAATGTTACACAGAT
AAAAGTAGCCAAGGGCACCTGTAACTGGGAATCTGAGTTTGACCTAAAAGTCATTAAAAT
AACATGAATCCCATTAATAAAAAAAAAAAAAA

333/392

FIGURE 320

MSALWLLLGLLALMDLSESSNWGCYGNIQSLDTPGASCGIGRRHGLNYCGVRASERLAEI
DMPYLLKYQPMQTIGQKYCMDPAVIAGVLSRKSPGDKILVNMGDRTSMVQDPGSQAPTS
WISESQVSQTTEVLTTRIKEIQRRFPTWTPDQYLRGGLCAYSGGAGYVRSSQDLSCDFCN
DVLARAKYLKRHGF

Important features of the protein:**Signal peptide:**

amino acids 1-19

N-myristoylation sites:

amino acids 23-29, 26-32, 35-41, 45-51, 50-56, 76-82, 156-162

Amidation site:

amino acids 40-44

334/392

FIGURE 321

GCCTTATAAAGTAGCCTCTGCATCTGCCTGCCTCGGGCAGAGGAGGGCTACCCTGGGGCT
GAGAGTTCACCTGTCTCAGGAACCACCTGAGCCCACAGATCCTGTGGGCAGCGGCCAGGG
CAGCCATGGCTTGGGCAAGTAGGCTGGGCCTGCTGCTGGCACTGCTGCTGCCCCGTGGTCTG
GTGCCTCCACGCCAGGCACCGTGGTCCGACTCAACAAGGCAGCATTGAGCTACGTGTCTG
AAATTGGGAAAGCCCCCTCTCCAGCGGGCCCTGCAGGTCACCTGTCCCTCATTTCTGGACT
GGAGTGGAGAGGCGCTTCAGCCCACCAGGATCCGGATTCTGAATGTCCATGTGCCCCGCC
TCCACCTGAAATTCATTGCTGGTTTCGGAGTGCGCCTGCTGGCAGCAGCTAATTTTACTT
TCAAGGTCTTTCGCGCCCCAGAGCCCCCTGGAGCTGACGCTGCCTGTGGAACCTGCTGGCTG
ACACCCGCGTGACCCAGAGCTCCATCAGGACCCCTGTGGTCAGCATCTCTGCCTGCTCTT
TATTCTCGGGCCACGCCAACGAGTTTGATGGCAGTAACAGCACCTCCACGCGCTGCTGG
TCCTGGTGCAGAAGCACATTAAAGCTGTCTTGAGTAACAAGCTGTGCCTGAGCATCTCCA
ACCTGGTGCAGGGTGTCAATGTCCACCTGGGCACCTTAATTGGCCTCAACCCCGTGGGTCT
CTGAGTCCCAGATCCGCTATTCCATGGTCAGTGTGCCCCTGTCACCAGTGACTACATTT
CCCTGGAAGTCAATGCTGTTCTCTTCTGCTGGGCAACCCCATCATCCTGCCCACGGATG
CCACCCCTTTTGTGTTGCCAAGGCATGTGGGTACCGAGGGCTCCATGGCCACCGTGGGCC
TCTCCAGCAGCTGTTTGACTCTGCGCTCCTGCTGCTGCAGAAGGCCGGTGCCCTCAACC
TGGACATCACAGGGCAGCTGAGGTCGGATGACAACCTGCTGAACACCTCTGCTCTGGGCC
GGCTCATCCCGAGGTGGCCCGCCAGTTTCCCGAGCCCATGCCTGTGGTGCTCAAGGTGC
GGCTGGGTGCCACACCTGTGGCCATGCTCCACACAAACAACGCCACCCTGCGGCTGCAGC
CCTTCGTGGAGGTCTGGCCACAGCCTCCAACTCGGCTTTCCAGTCCCTCTTCTCCCTGG
ATGTGGTAGTGAACCTTGAGACTCCAGCTCTCTGTGTCCAAGGTGAAGCTTCAGGGGACCA
CGTCTGTGCTGGGGGATGTCCAGCTCACGGTGGCCTCCTCCAACGTGGGCTTCATTGATA
CAGATCAGGTGCGCACACTGATGGGCACCGTTTTTTGAGAAGCCCCCTGCTGGACCATCTCA
ATGCTCTCTTGCCATGGGAATTGCCCTCCCTGGTGTGGTCAACCTCCACTATGTTGCCC
CTGAGATCTTTGTCTATGAGGGCTACGTGGTGATATCCAGTGGACTCTTCTACCAGAGCT
GAGGCAAGACCACTGGGAGGCCTGAGAGTGGGCCAGCTCGCTGCTCAGGCGAATTTCTCA
TTTCAAGCCACTGGGGAACTGAGGCAAAACCATACTTAGTCATCACCAACAAGCTGGAC
TGCTTAGCTGGGCTGTTTTATCTTCCCTGAGTGCCTGGGTCTCCCTCCCTCACTTCTGCC
CTTTCCCTTCTCCTCCTCTTCTCCTCCCTCTTCCCTCATCTCCCCCTCCTTCTCTGC
CCCACCCAGGGGGGAGCAGACTGCTCCTCCAGGCTGTATAGACCTGCCCTCTTGCATTA
AACAACCTCTCTTGAGCTGC

335/392

FIGURE 322

MAWASRLGLLLALLLPVVGASTPGTVVRLNKAALSYVSEIGKAPLQRALQVTVPHFLDWS
GEALQPTRIRILNVHVPRLHLKFIAGFGVRLLAANFTFKVFRAPPELELTLPVELLADT
RVTQSSIRTPVVSISACSLFSGHANEFDGSNSTSHALLVLVQKHIKAVLSNKLCLSISNL
VQGVNVHLGTLIGLNPVGPESQIRYSMVSVPVTSTSDYISLEVNAVLFLLGNPIILPTDAT
PFVLPRHVGTEGSMATVGLSQQLFDSALLLLQKAGALNLDITGQLRSDDNLLNTSALGRL
IPEVARQFPEPMPVVLKVR LGATPVAMLHTNNATLRLQPFVEVLATASNSAFQSLFSLDV
VVNLRLQLSVSKVKLQGTTSVLGDVQLTVASSNVGFIDTDQVRTLMGTVFEEKPLLDHLNA
LLAMGIALPGVVNLHYVAPEIFVYEGYVVISSGLFYQS

Important features of the protein:**Signal peptide:**

Amino acids 1-20

Transmembrane domain:

Amino acids 217-236

N-glycosylation sites:

Amino acids 96-100;151-155;293-297;332-336

N-myristoylation sites:

Amino acids 8-14;149-155;189-195;249-255;252-258;283-289

LBP / BPI / CETP family proteins:

Amino acids 22-50; 251-287

336/392

FIGURE 323

TTGAAAATCTACTCTATCAGCTGCTGTGGTTGCCACCATTCTCAGGACCCTCGCCATGAA
AGCCCTTATGCTGCTCACCCTGTCTGTTCTGCTCTGCTGGGTCTCAGCTGACATTGCTG
TCACTCCTGCTACAAGGTCCCTGTGCTGGGCTGTGTGGACCGGCAGTCCTGCCGCCTGGA
GCCAGGACAGCAATGCCTGACAACACATGCATACCTTGGTAAGATGTGGGTTTTCTCCAA
TCTGCGCTGTGGCACACCAGAAGAGCCCTGTCAGGAGGCCTTCAACCAAACCAACCGCAA
GCTGGGTCTGACATATAACACCACCTGCTGCAACAAGGACAACCTGCAACAGCGCAGGACC
CCGGCCCCACTCCAGCCCTGGGCCTTGTCTTCCTTACCTCCTTGGCTGGCCTTGGCCTCTG
GCTGCTGCACTGAGACTCATTCCATTGGCTGCCCCCTCCTCCCACCTGCCTTGGCCTGAGC
CTCTCTCCCTGTGTCTCTGTATCCCCTGGCTTTACAGAATCGTCTCTCCCTAGCTCCCAT
TTCTTTAATTAAACACTGTTCCGAGTGGTCTCCTCATCCATCCTTCCCACCTCACACCCT
TCACTCTCCTTTTTCTGGGTCCCTTCCCCTTCCCTTCCAGGACCTCCATTGGCTCCTAGA
AGGGCTCCCCACTTTGCTTCCTATACTCTGCTGTCCCCTACTTGAGGAGGGATTGGGATC
TGGGCCTGAAATGGGGCTTCTGTGTTGTCCCCAGTGAAGGCTCCCACAAGGACCTGATGA
CCTCACTGTACAGAGCTGACTCCCCAAACCCAGGCTCCCATATGTACCCCATCCCCCATA
CTCACCTCTTCCATTTTGAGTAATAAATGTCTGAGTCTGGAAAAAAAAAAAAAAAAAAAA

337/392

FIGURE 324

MKALMLLTLSVLLCWVSADIRCHSCYKVPVLGCVDRQSCRLEPGQQCLTTHAYLGKMWVF
SNLRCGTPEEPCQEAFNQTNRKLGITYNTTCCNKDNCNSAGPRPTPALGLVFLTSLAGLG
LWLLH

Important features of the protein:

Signal peptide:

amino acids 1-18

N-glycosylation sites:

amino acids 77-81, 88-92

N-myristoylation site:

amino acids 84-90

Ly-6 / u-PAR domain protein signature:

amino acids 85-98

338/392

FIGURE 325

ACGGGCCGCGACGGCAGTGACGTAGGGTTGGCGCACGGATCCGTTGCGGCTGCAGCTCTG
CAGTCGGGCCGTTTCCTTCGCCGCCGCCAGGGGTAGCGGTGTAGCTGCGCAGCGTCGCGCG
CGCTACCGCACCCAGGTTTCGGCCCCGTAGGCGTCTGGCAGCCCGGCGCCATCTTCATCGAG
CGCCATGGCCGCGAGCCTGCGGGCCGGGAGCGGCCGGGTACTGCTTGCTCCTCGGCTTGCA
TTTGTTTCTGCTGACCGCGGGCCCTGCCCTGGGCTGGAACGACCCTGACAGAATGTTGCT
GCGGGATGTAAAAGCTCTTACCCTCCACTATGACCGCTATACCACCTCCCGCAGGCTGGA
TCCCATCCACAGTTGAAATGTGTTGGAGGCACAGCTGGTTGTGATTCTTATACCCCAA
AGTCATACAGTGTCAGAACAAAGGCTGGGATGGGTATGATGTACAGTGGGAATGTAAGAC
GGACTTAGATATTGCATACAAATTTGGAAAACTGTGGTGAGCTGTGAAGGCTATGAGTC
CTCTGAAGACCAGTATGTACTAAGAGGTTCTTGTTGGCTTGAGTATAATTTAGATTATAC
AGAACTTGGCCTGCAGAACTGAAGGAGTCTGGAAAGCAGCACGGCTTTGCCTCTTCTC
TGATTATTATTATAAGTGTCCTCGGCGGATTCTGTAAACATGAGTGGATTGATTACCAT
CGTGGTACTCCTTGGGATCGCCTTTGTAGTCTATAAGCTGTTCTGAGTGACGGGCAGTA
TTCTCCTCCACCGTACTCTGAGTATCCTCCATTTTCCCACCGTTACCAGAGATTCACCAA
CTCAGCAGGACCTCCTCCCCCAGGCTTTAAGTCTGAGTTCACAGGACCACAGAATACTGG
CCATGGTGCAACTTCTGGTTTTTGGCAGTGCTTTTACAGGACAACAAGGATATGAAAATTC
AGGACCAGGGTTCTGGACAGGCTTGGGAACTGGTGGAATACTAGGATATTTGTTTGGCAG
CAATAGAGCGGCAACACCCTTCTCAGACTCGTGGTACTACCGTCTTATCCTCCCTCCTA
CCCTGGCACGTGGAATAGGGCTTACTCACCCCTTCATGGAGGCTCGGGCAGCTATTCCGT
ATGTTCAAACCTCAGACACGAAAACCAGAACTGCATCAGGATATGGTGGTACCAGGAGACG
ATAAAGTAGAAAAGTTGGAGTCAAACACTGGATGCAGAAATTTTGGATTTTTCATCACTTT
CTCTTTAGAAAAAAGTACTACCTGTAAACAATTGGGAAAAGGGGATATTCAAAGTTCT
GTGGTGTTATGTCCAGTGTTAGCTTTTTGTATTCTATTATTTGAGGCTAAAAGTTGATGTG
TGACAAAATACTTATGTGTTGTATGTCAGTGTAAACATGCAGATGTATATTGCAGTTTTTG
AAAGTGATCATTACTGTGGAATGCTAAAAATACATTAATTTCTAAAACCTGTGATGCCCT
AAGAAGCATTAAAGAATGAAGGTGTTGTACTAATAGAACTAAGTACAGAAAATTTAGTT
TTAGGTGGTTGTAGCTGATGAGTTATTACCTCATAGAGACTATAATATTCTATTTGGTAT
TATATTATTTGATGTTTGCTGTTCTTCAAACATTTAAATCAAGCTTTGGACTAATTATGC
TAATTTGTGAGTTCCTGATCACTTTTGAGCTCTGAAGCTTTGAATCATTCAGTGGTGGAGA
TGGCCTTCTGGTAACTGAATATTACCTTCTGTAGGAAAAGGTGGAAAATAAGCATCTAGA
AGGTTGTTGTGAATGACTCTGTGCTGGCAAAAATGCTTGAAACCTCTATATTTCTTTCGT
TCATAAGAGGTAAAGGTCAAATTTTTCAACAAAAGTCTTTTAATAACAAAAGCATGCAGT
TCTCTGTGAAATCTCAAATATTGTTGTAATAGTCTGTTTCAATCTTAAAAAGAATCA

339/392

FIGURE 326

MAAACGPGAAGYCLLLGLHLFLLTAGPALGWNDPDRMLLRDVKALTLHYDRYTTSRRLDP
IPQLKCVGGTAGCDSYTPKVIQCQNKGDYDVQWECKTDLDIAYKFGKTVVSCEGYESS
EDQYVLRGSCGLEYNLDYTELGLQKLKESGKQHGFAFSFYKSSADSCNMSGLITIV
VLLGIAFVVYKFLSDGQYSPPPYSEYPPFSHRYQRFTNSAGPPPPGFKSEFTGPQNTGH
GATSGFGSAFTGQQGYENSGPGFWTGLGTGGILGYLFGSNRAATPFSDSWYYPSYPPSY
GTWNRAYSPLHGGSGSYSVCSNSDTKTRTASGYGGTRRR

Signal peptide:
amino acids 1-30

Transmembrane domain:
amino acids 171-190

N-glycosylation site:
amino acids 172-176

Glycosaminoglycan attachment sites:
amino acids 244-248, 259-263, 331-335

Tyrosine kinase phosphorylation site:
amino acids 98-106

N-myristoylation sites:
amino acids 68-74, 69-75, 131-137, 241-247, 247-253, 266-272,
270-276, 278-284, 312-318

340/392

FIGURE 327

GGCACGAGGTGGAAGGGCTTTTACAAACAGATTGCTGGCCCCACCCCCCAGAATTTCTCA
TCAGGAGTGGGCAAGACCAATCATTTGCATTTCTGACAAGTTCCCAGGAGCTGCAGCTGC
TGGCCCTGGAACCACACTTTGAGAACCACTGCTTTAGACCAAACACCAAAGGAAGATGCA
GCCACCCTCCTTTACATGTCACAACGCTCAGGGTCCATGAGTACCTCAGGCTGTCCAGCT
GAGCTCCACCTGCAGCAGCCGAGATTCCCGACTCGCTCCACCATTGGGGGCTAGGAGTGA
AGCGTGTACCAATGGTCAGCTCATGGCCAGCCAGGAAAGCCTCTCTGCTGTGCGTCTGTG
CAGTTCTTGTTCTTCCCTGGAGGACTCTTGATCGCCTGTGATCTTGCCAGGAGACCAG
GTGCCTGGGTCCCTTCCTGGAAGGGGACAAGTTACACACCCAGCCCCATTTTCCCACCA
ACTTCTACATGCCTTGGGAGAACCTTCTACATGTTGGCTGCCCCCTTCCCCTATTTTCAGC
AGTGCCCAAGTCCTGCTTATAAACCTGAGGCCTGCTCCCCATACCTTCCCTGTGCAAGTGC
CAGCCGTTATTCAGGCAGCCCAATGTTGTTGAGGCCAGATGGATTCTTGAAGCAGCTG
GCCCATGGATGTGAGTCATCACAGTATTCTAGAAACAGAGAAGAGGTCTTAACCTAATGC
GCATAGAGAAATTGTTCTCATTGTAAACATACCCCTGTCCTTAGCTGATCTAGGTGGAAG
CCCAGCTTCATGTGCTAGGGGGCATGATAATGATAATAAAGGAATTGTATCTAGGACTAA

341/392

FIGURE 328

MVSSWPARKASLLCVC AVL VLPWRTL GSPVILARRPGAWVPSWKGTSYTPQPHFPTNFYM
PWENLLHVGCP LPLFQQCPVLLINLRPAPHTFPVQVPAVIPGSPMLLRPDGFLEAAGPWM

Signal peptide:
amino acids 1-27

cAMP- and cGMP-dependent protein kinase phosphorylation site:
amino acids 8-12

342/392

FIGURE 329

CAAAGAGTAGTCAGTCCCTTCTTGGCTCTGCTGACACTCGAGCCCACATTCCATCACCTG
CTCCCAATCATG CAGGTCTCCACTGCTGCCCTTGCCGTCCTCCTCTGCACCATGGCTCTC
TGCAACCAGGTCTCTCTGCACCACTTGCTGCTGACACGCCGACCGCTGCTGCTTCAGC
TACACCTCCCGACAGATTCCACAGAATTTTCATAGCTGACTACTTTGAGACGAGCAGCCAG
TGCTCCAAGCCCAGTGTTCATCTTCCTAACCAAGAGAGGCCGGCAGGTCTGTGCTGACCCC
AGTGAGGAGTGGGTCCAGAAATACGTCAGTGACCTGGAGCTGAGTGCCCTGAGGGGTCCAG
AAGCTTCGAGGCCCGAGCGACCTCAGTGGGCCAGTGGGGAGGAGCAGGAGCCTGAGCCTT
GGGAACATGCGTGTGACCTCTACAGCTACCTCTTCTATGGACTGGTTATTGCCAAACAGC
CACACTGTGGGACTCTTCTTAACTTAAATTTTAAATTTATTTATACTATTTAGTTTTTATA
ATTTATTTTTGATTTACAGTGTGTTTGTGATTGTTTGCTCTGAGAGTCCCCCTGTCCC
CTCCCCCTTCCCTCACAGTGTGTCTGGTGACAACCGAGTGGCTGTTCATCGGCCTGTGTAG
GCAGTCATGGCACCAAAGCCACCAGACTGACAAATGTGTATCAAATGCTTTTGTTCAGGG
CTGTGATCGGCCTGGGGAAATAATAAAGATGTTCTTTTAAACGGTAAAAAA

343/392

FIGURE 330

MQVSTAALAVLLCTMALCNQVLSAPLAADTPTACCFSTSRQIPQNFIADYFETSSQCSK
PSVIFLTKRGRQVCADPSEEWVQKYVSDLELSA

Signal sequence:

1-23

**Small cytokines (intercrine/chemokine) C-C subfamily
signature:**

1-35, 2-36, 10-44, 34-74, 50-90

Small cytokines (intecrine/chemokine):

24-89

344/392

FIGURE 331

GGCACGAGGTGAGACTTTAAATGAAATGTCTCACAAGCTAGGTGATCCAGGTTTTGTGGT
CTTTGCAACCCTTGTGGTCATTGTGGCCTTGATATTAATCTTCGTGGTGGGTCCTCGCCA
TGGACAGACAAACATTCTTGTGTACATAACAATCTGCTCTGTAATCGGCGCGTTTTTCAGT
CTCCTGTGTGAAGGGCCTGGGCATTGCTATCAAGGAGCTGTTTGCAGGGAAGCCTGTGCT
GCGGCATCCCCTGGCTTGGATTCTGCTGCTGAGCCTCATCGTCTGTGTGAGCACACAGAT
TAATTACCTAAATAGGGCCCTGGATATATTCAACACTTCCATTGTGACTCCAATATATTA
TGTATTCTTTACAACATCAGTTTTAACTTGTTGAGCTATTCTTTTTAAGGAGTGGCAAGA
TATGCCTGTTGACGATGTCATTGGTACTTTGAGTGGCTTCTTTACAATCATTGTGGGGAT
ATTCTTGTGTGCATGCCTTTAAAGACGTGAGCTTTAGTCTAGCAAGTCTGCCTGTGTCTTT
TCGAAAAGACGAGAAAGCAATGAATGGCAATCTCTCTAATATGTATGAAGTTCTTAATAA
TAATGAAGAAAGCTTAACCTGTGGAATCGAACAACACACTGGTGAAGTGTCTCCCGAAG
AAATGGAAATCTGACAGCTTTTTTAAGAAAGGTGTAATTAAAGGTTAATCTGTGATTGTTA
TGAAGTGAATTTGAATATCATCAGAATGTGTCTGAAAAAACATTTGTCTCAAATAATGTT
CTTTAAAGGCAATCTTTTTAAAGATTTCACTAATTTGGACCAAGAAATTACTTTTCTTGT
ATTTAAACAAACAATGGTAGCTCACTAAAATGACCTCAGCACATGACGATTTCTATTAAC
ATTTTATTGTTGTAGAAGTATTTTACATTTTCATCCCTTCTCCAAAAGCCGAATGCACTA
ATGACAGTTTTAAGTCTATGAAAATGCTTTATTTTTTCATTGGTGATGAAAGTCTGAAAT
GTGCATTTGTGATCCCCACTCCATCAATCCCTGACCATGTAAGGCTTTTTTATTTTAAAA
AAACAGAGTTATCCCAATACATTATCCTGTGATTTACCTTACCTACAAAAGTGGCTCCTG
TTTGTTTGTGATGATGATTGGTTTTATTTTTGAAATATTTATTAAGGGAAAATAAGTTACT
GAATGAAGGAACCTCTTTCTTACAAAACAAAAAAAGGGCAGAAATCACCCCAAGGAACG
ATTTCTCAGGTTGAGATGATCACCGTGAATCCGGCTTCCTCTGAGCATTGATGGCCTTA
GCACCTCATCAAGCCAGCACATCCTGCCTGCTGTTGCAGCCTGGCTGGGTTTATTCTTCA
GTTACCCCTAATCCCATGATGCCTGGAACCTTGATTACCGTTTTACATCAGCTCTTGTA
TTTCAGTATATTTTCATAATGAGTTATATTGTCATTTAGACTTTGAACAGCTCTGGGAAA
TAGAAGACTAGGTTGTTTCTTAAATTTAGCTCATGTTATAATAAAAAGTTGAAATG

345/392

FIGURE 332

MSHKLGDPGFVVFATLVVIVALILIFVVGPRHGQTNILVYITICSVIGAFSVSCVKGLGI
AIKELFAGKPVL RHPLAWILLLSLIVCVSTQINYNLRALDIFNTSIVTPIYYVFFTTSVL
TCSAILFKEWQDMPVDDVIGTSLSGFFTIIIVGIFLLHAFKDVSFSLASLPVSFRKDEKAMN
GNLSNMYEVLNNNEESLTCGIEQHTGENVSRNGNLTAF

Signal sequence:

1-33

Transmembrane domain:

40-60, 70-90, 103-123, 139-159

N-glycosylation site:

103-106, 182-185, 208-211, 215-218

N-myristoylation site:

57-62, 140-145, 181-186, 214-219

346/392

FIGURE 333

[illegible]

347/392

FIGURE 334

MAAGDGDVKLGTLGSGSESSNDGGSESPGDAGAAAEGGGWAAAALALLTGGGEMLLNVAL
VALVLLGAYRLWVRWGRRGLGAGAGAGEESPATSLPRMKKRDFSLEQLRQYDGSRNPRIL
LAVNGKVFDVTKGSKFYGPAGPYGIFAGR DASRGLATFCLDKDALRDEYDDLSDLNAVQM
ESVREWEMQFKEKYDYVGRLLKPGEEPSEYTDDEEDTKDHNKQD

Signal sequence:
None

Transmembrane domain:
45-65

Tyrosine kinase phosphorylation site:
202-210

N-myristoylation site:
11-16, 16-21, 37-42, 38-43, 79-84, 81-86, 83-88, 144-149

Amidation site:
75-78

349/392

FIGURE 336

TGRGYCGDHESSFGAMEEPGATPQPYLGLLLEELRRVVAALPEGMRPDSNLYGFPWELVI
CAAVVGFFAVLFFLWRSFRSVRSRLYVGREKKLALMLSGLIEEKSLLLEKFSLVQKEYEG
YEVESLKDASFEKEATEAQSLEATCEKLNRSNSELEDEILCLEKELKEEKSXHSEQDEL
MADISKRIQSLEDESKSLKSQVAEAKMTFQIFQMNEERLKIAIKDALNENSQQLQESQKQL
LQEAENVWKEQVSELNKQKVTFFEDSKVHAEQVLNDKESHIKTLTERLLKMKDWAAMLGEDI
TDDDNLELEMNSESENGAYLDNPPKGALKKLIHAAKLNASLKTLEGERNQIYIQLSEVDK
TKEELTEHIKNLQTQQASLQSENFHFNENQKLQQKLKVMTELYQENEMKLHRKLTVEEN
YRLEKEEKLKSVDEKISHATEELETYRKRAKDLEEEELERTIHSYQGQIISHEKKAHDNWL
AARNAERNLNDLRKENAHNRQKLTETELKFELLEKDPYALDVPNTAFGRGSRGPGNPDLH
QITNERGESSCDRLTDPHRAPSDTGSLSPWDQDRMMFPPPGQSYPDSSALPPQRQDRFC
SNSGRLSGPAELRSFNMPSLDKMDGSMPSSEMRNDTKDDLGNLNPVDSLSLPAENEATG
PGFVPPPLAPIRGPLFPVDARGPFLRRGPPFPPPPPGAMFGASRDYFPFRDFPGPPAPF
AMRNVYPPRGFPYLPFRPGFFPPPHSEGRSEFPSPGLIPPSNEPATEHPEPQOET

Signal sequence:

None

Transmembrane domain:

54-74

N-glycosylation site:

150-153, 338-341, 636-639

cAMP- and cGMP-dependent protein kinase phosphorylation site:

413-416

Tyrosine kinase phosphorylation site:

414-421

N-myristoylation site:

466-417, 625-630, 697-702

Leucine zipper pattern:

142-163

350/392

FIGURE 337

GGACTGCGGTCTCGGGCAGCAATGGCCGAGAAGCGCGACACACGGGACTCCGAAGCCCAG
CGGCTCCCCGACTCCTTCAAGGACAGCCCCAGTAAGGGCCTTGGACCTTGGCGATGGATT
TTGGTGCGGTTCTCATTCTTATTCACCGTTATAACTTTCCCAATCTCAATATGGATGTGC
ATAAAGATTATAAAAGAGTATGAAAGAGCCATCATCTTTAGATTGGGTGCGATTTTACAA
GGAGGAGCCAAAGGACCTGGTTTGTATTTTATTCTGCCATGCACTGACAGCTTCATCAAA
GTGGACATGAGAACTATTTCAATTTGATATTCTCTCCTCAGGAGATCCTGACAAAGGATTCA
GTGACAATTAGCGTGGATGGTGTGGTCTATTACCGCGTTTCTGAGAATGCAACCCTGGCTGTG
GCAAAATATCACCACGCTGACTCAGCAACCCGTCTTTTGGCACAACACTACTCTGAGGAAT
GTTCTGGGCACCAAGAATCTTTCTCAGATCCTCTCTGACAGAGAAGAAATTGCACACAAC
ATGCAGTCTACTCTGGATGATGCCACTGATGCCTGGGGAATAAAGGTGGAGCGTGTGGAA
ATTAAGGATGTGAAACTACCTGTGCAGCTCCAGAGAGCTATGGCTGCAGAAGCAGAAGCG
TCCCGCGAGGCCCGCGCCAAGGTTATTGCAGCCGAAGGAGAAATGAATGCATCCAGGGCT
CTGAAAGAAGCCTCCATGGTCATCACTGAATCTCCTGCAGCCCTTCAGCTCCGATACCTG
CAGACACTGACCACCATTGCTGCTGAGAAAACTCAACAATTGTCTTCCCTCTGCCATA
GATATGCTGCAAGGAATCATAGGGGCAAAACACAGCCATCTAGGCTAGTGTAGAGATGAG
CGCTAGCCTTCCAAGCATGAAGTCGGGGACCAATTAGCCTTTAACTCATAAAGAGAGGG
TAGGGCTTTTCTTTTCCATATGTCAATTGTGGTGTTCCCAGAATGTATAGCAGTTATAA
AAATAGGTGAAAGAATTGTTAGCTTGTAATACTGAGAGATTGGTGATTTATATAAGGTA
ATCTGTTAGTCTTAAATAGTTAAAGTTTGTATTTTATAGATTATTATGTAGTAGGTTAG
ATCCCTCTTGTTTGGACTTCCACTGACTCATTCTGAACCCCTAAGCACCCAGGCCACAG
GCAAGAACCTGGGCTGTAAGTCCACCTGACACCGCTGACTGGCTAAATGCTTTGCAGAA
AGTGATGACCTTACACCACAACCAGCTTCTCCAGGTCATATGTGCCTTACCTCCAGAAGT
CTTTTTTTTTTTTTTTTTTCTGAGATGGAGTTTCACTCTTGTTGCCAGGCTGGAGTGCAA
TAGCATGATCTCGGCTCACTGCAACCTCCGCCTCCTGGGTTCAAGAGATTCTCCTGCCTC
AGCCTCCCCAGTAGCTGGGATTACAGGCTCATGCCACCATGCCAGCTAATTTTTGTATT
ATTATTATTGTTTTTTAGTAGAGACGGGGTTTACCATGTTGGCCAGGCTAGTCACGAAC
TCCTAACCTCAGGTGATCCACCCACCTCTGCCTCCAAAGTGCTGGATTACAGGCTGAGCT
ACCACCTGGTTTGGAGAGTCTTAATTAATTGAAATTTCCCTAATGTTCAATTTATTTTCT
AAATCCAGCCGTGTTTTCAGAATAATCCTTACTTGAGAGTAGCCATTTTCTTGTGTACTTG
TCAGAACTAGAGGAAATAGCCAAGACTAATGAAAAACATTACTCTAACCTTAAAAGACT
TTTAAATTCACTACTAGAGTGGTCATTTTAAAATACATCCATGTTTAACTTATTTTGA
GCCTTCTTTTATGAGTAAATGATTCCTCCTGTTCTGTCTTCAAACCAGCTAAATATT
TGTCACAAAAGTGACTTTTTTCTCACTGTTGCCTATTTTCATATATCAGGTTTTAAATAG
TTTTAATTTTTTAATAAAATTTTTCTCTACGTTCTATATGCAATTGTTATATATCTATTT
GAATAGCTGAAGGACTAAAATACTTTTTTAAGAGATAACTTCAGGAAACCATTATATTTT
ACTATCTGCATGCTGTAACTGTGGTACACTGTGAAATATGTTGATTACAAACCCATTCA
TTACATAGTATAAGGAATTCACAGTATATTGACTATATAGTGTCTAATGACTGGGCAGAT
ACTGTCAACTTACAATATCTATATAGAGAGGCTTTAACTTACCTTACTCATTCTCTATG
ATGTATGACTTGATGCTGAAAGAGGAAGCTGGTCAGCTCCTCATGGACAACAAATCTTA
GTCTATAATATTAGGAGACATCTCTAGTTTTGCAATGTCTGTGAATCTGAGCAACCTGG
ACTTCTGCTTACTGGCCAGAAAGCTGGCGGGTGACATTTGTAACATTTCTCTTTGAGAC
TCTGAGTTTCACTAGAGAAGTCTAAGCATAACAGCTTTCTTTCCAGCACGAGCCTTTAT
AGCTCTCTTTAGCTCAACCACTCTGTCCATCCAGCCAATGGATGTCTTCCCTGTACCCA
ATTCAGCTTATTTTAGGGAAGCCTTGAACTACCATGTATCTGGCTCTAGCTGAGTTAT
TGAGGATTGAGCCAGTGCAACGTTAACTCAGTGCATTACATTTGATTTAAATGATGGT
TTTATCTGTTGTGTGAAGTGGTTCAACCTTGAGGACCAGGAGCCTCCATATCCTGACTGA
AAACCTTTTCTGAGACTTAGAGTAACAGTACTTTTGGTTCCTTGAGTTCTCCTGTCTCCA
GATACCTAAATGACCTTGACTTTTTCTGCCTTGTTGAATTCGTAGTCCAATCAGCTGAAATT
AAATCACTTGGGAGGGACGCATAGAAGGAGCTCTAGGAACACAGTGCCAGTGCAGAAGTT
TCTCCAGGTGGCCTCCCTTTCCAACAATGTACATAATAAAGTGATGCACCTTCACT

351/392

FIGURE 338

MAEKRDTRDSEAQRLPDSFKDSPSKGLGPCGWILVAFSFLFTVITFPISIWMCIKIKEY
ERAIIFRLGRILQGGAKGPGLFFILPCTDSFIKVD MRTISFDIPPQEILTKDSVTISVDG
VYYRVQ NATLAVANITNADSATRLLAQTTLRNVLTGNLSQILSDREEIAHNMQSTLDD
ATDAWGIKVERVEIKDVKL PVQLQRAMAAEAEASREARAKVIAAEGEMNASRALKEASMV
ITESPAALQLRYLQTLTTIAAEKNSTIVFPLPIDMLQGIIGAKHSHLG

Signal sequence:

1-45

Transmembrane domain:

None

N-glycosylation site:

128-131, 135-138, 159-162, 229-232, 264-267

**cAMP- and cGMP-dependent protein kinase phosphorylation
site:**

4-7

N-myristoylation site:

26-31, 278-283, 281-286

SPFH domain/Band 7 family:

39-230

352/392

FIGURE 339

TCTAGAGCCCTCTCCCAACATGGCGGCCTCAGCAAAAAAGAAGAATAAGAAGGGGAAGAC
TATCTCCCTAACAGACTTTCTGGCTGAGGATGGGGGTACTGGTGGAGGAAGCACCTATGT
TTCCAAACCAGTCAGCTGGGCTGATGAAACGGATGACCTGGAAGGAGATGTTTCGACCAC
TTGGCACAGTAACGATGACGATGTGTATAGGGCGCCTCCAATTGACCGTTCCATCCTTCC
CACTGCTCCACGGGCTGCTCGGGAACCCAATATCGACCGGAGCCGTCTTCCCAATCGCC
ACCCTACACTGCTTTTCTAGGAAACCTACCCTATGATGTTACAGAAGAGTCAATTAAGGA
ATTCTTTCGAGGATTAAATATCAGTGCAGTGCCTTTACCACGTGAACCCAGCAATCCAGA
GAGGTTGAAAGGTTTTGGTTATGCTGAATTTGAGGACCTGGATTCCCTGCTCAGTGCCCT
GAGTCTCAATGAAGAGTCTCTAGGTAACAGGAGAATTCGAGTGGACGTTGCTGATCAAGC
ACAGGATAAAGACAGGGATGATCGTTCCTTTGGCCGTGATAGAAATCGGGATTCTGACAA
AACAGATACAGACTGGAGGGCTCGTCCTGCTACAGACAGCTTTGATGACTACCCACCTAG
AAGAGGTGATGATAGCTTTGGAGACAAGTATCGAGATCGTTATGATTACAGACCGGTATCG
GGATGGGTATCGGGATGGGTATCGGGATGGCCACGCCGGGATATGGATCGATATGGTGG
CCGGGATCGCTATGATGACCGAGGCAGCAGAGACTATGATAGAGGCTATGATTCCCGGAT
AGGCAGTGGCAGAAGAGCATTGGCAGTGGGTATCGCAGGGATGATGACTACAGAGGAGG
CGGGGACCGCTATGAAGACCGATATGACAGACGGGATGATCGGTCTGGAGCTCCAGAGA
TGATTACTCTCGGGATGATTATAGGCGTGATGATAGAGGTCCCCCCCCAAAGACCCAACT
GAATCTAAAGCCTCGGAGTACTCCTGAAGAAGATGATTCCTCTGCTAGTACCTCCAGTC
CACTCGAGCTGCTTCTATCTTTGGAGGGGCAAAGCCTGTTGACACAGCTGCTAGAGAAAG
AGAAGTAGAAGAACGGCTACAGAAGGAACAAGAGAAGTTGCAGCGTCAGTGGAAATGAGCC
AAAAGTAGAACGACGGCCTCGGGAGAGACACCCAAGCTGGCGAAGTGAAGAACTCAGGA
ACGGGAACGGTCGAGGACAGGAAGTGAGTCATCACAAGCTGGGACCTCCACCACATCTAG
CAGAAATGCACGAAGGAGAGAGAGTGAGAAGTCTCTAGAAAATGAAACACTCAATAAGGA
GGAAGATTGCCACTCTCCAACCTCTAAACCTCCCAAACCTGATCAGCCCCATAAGGTAAT
GCCAGCCCCCTCCACCAAAGGAGAATGCTTGGGTGAAGCGAAGTTCTAACCCCTCCTGCTCG
ATCTCAGAGCTCAGACACAGAGCAGCAGTCCCCTACAAGTGGTGGGGGAAAAGTAGCTCC
AGCTCAACCATCTGAGGAAGGACCAGGAAGGAAAGATGAAAATAAAGTAGATGGGATGAA
TGCCCCAAAAGGCCAAACTGGGAACTCTAGCCGTGGTCCAGGAGACGGAGGGAACAGAGA
CCACTGGAAGGAGTCAGATAGGAAAGATGGCAAAAAGGATCAAGACTCCAGATCTGCACC
TGAGCCAAAAGAAACCTGAGGAAAATCCAGCTTCTAAGTTCAGTTCGCAAGCAAGTATGC
TGCTCTCTCTGTTGATGGTGAAGATGAAAATGAGGGAGAAGATTATGCCGAATAGACCTC
TACATCCTGTGCTTTTCTCCTAGTTTCTCTCCACCCTGGAACATTTCGAGAGCAAATCAA
ACCTCTATCCAGACAAGACAAAATAAACTCAACATCTCCTGAAGACCTTTCTTACCTTT
TTTTAAAAACAAAAXTGAAATTATTTTGCATGCTGCTGCAGCCTTTAAAGTATTGAAGT
AACTGGAGAATTGCCAATACAGCCAGAGAGAAAGGGACTACAGCTTTTTAGAGGAAAAGT
TGTGGTGCGTTATGTACCATGCAGTTGCCAGTGTGATTAGTGCCTAGGGGTCTCATTTA
GCAGAAATGGTAATGACAGTGATATAATGCCTGGAACCTGGTTGGGCAGTAGGGGAGGGA
GGTAGAAGGAAAAGTGTGAGATTTCTACCTTTTAGTTTTTCATCCTATTGTGGCATATATG
AATTCTCAAACATTATCTGAATAAAATTTTCCACTCTTGGAAGGTAGATTTAGCCTCAAG
TTGTTCTAGTCTCCAGGAGGCTGCCAGCCCCCTCCTCTTATTTAATTCTGAGTTTTGGGGG
CCAGCCTAGAGGGAATTCTTTTTTTTTTTTTTTTAAACCCCCAGGGGGTAGTTGGGAGT
GAGACTATAGGCCATAAAGAATGGGACTGCATTGGACCAAATAAATGGGAAAATCGTGG
TTTGAAAAGAAGCTTTTTGGGAAGTGATGAGTCATTTTGCACCAGGTAATAGGGGAAAATT
GTGTGACCTCCAGCAAACACATGAATGGTTATTTCTGGAGCCGGAAGCACTTGGGGGTC
GTGGTAATTCCCAGTGTTTTCTGTGTCCTAGTTTTTACCCTTTCTAAACACTGTCCTTTTTT
GAAAGTTTGAATATATCCACATTCTATTGAAACCTTGAACTAAAAATTTAGACTCTTA
TCGTATCTTAAGTTCTTCATGCTACTCTTAACCTCCAAAAAGCAGTATCTAAGTCACA
TACATGATGTCTTGGGCATTTTCTGAGCCATGGAGAACTCTGAAAGGAAGAATCGCTGCT
TTTCTCAAGCAAATCGGTTTCTTGATGTCTTTTGGTTCTCCTTGCTGCTCCTGATGCTT

354/392

FIGURE 340

MAASAKKKNKKGKTISLTDFLAEDGGTGGGSTYVSKPVSWADETDDLEGDVSTTWHSNDD
DVYRAPPIDRSILPTAPRAAREPNIDRSRLPKSPPYTAFLGNLPYDVTEESIKEFFRGLN
ISAVRLPREPSNPERLKGFYAEFEDLDSLLSALSLSNEESLGNRRIRVDVADQAQDKDRD
DRSFGRDRNRDSDKTDTDWRRARPATDSFDDYPPRRGDDSFQDKYRDRYDSRYRDGYRDG
YRDGPRRDMDRYGGDRYDDRGSRDYDRGYDSRIGSGRRAFGSGYRRDDDYRGGGDRYED
RYDRRDDRSWSSRDDYSRDDYRRDDRGPQRPKLNKPRSTPEEDDSSASTSQSTRAASI
FGGAKPVDTAAREREVEERLQKEQEKLRQWNEPKLERRPRERHPSWRSEETQERERSRT
GSESSQTGTSTTSSRNARRRESEKSLNETLNKEEDCHSPTSKPPKPDQPLKVMPPPPK
ENAVVKRSSNPPARSQSSDTEQQSPTSGGGKVAPAQPSEEGPGRKDENKVDGMNAPKGQT
GNSSRGPGDGGNRDHWKESDRKDGKKDQDSRSAPEPKPEENPASKFSSASKYAALSVDG
EDENEGEDYAE

Signal Sequence:

None

Transmembrane domain:

None

N-glycosylation site:

120-123, 448-451, 542-545

Glycosaminoglycan attachment site:

507-510

cAMP- and cGMP-dependent protein kinase phosphorylation site:

439-442, 486-489

Tyrosine kinase phosphorylation site:

225-233, 264-270

N-myristoylation site:

25-30, 26-31, 28-33, 118-123, 421-426, 428-433, 538-543

Amidation site:

276-279, 522-525, 563-566

Cell attachment sequence:

215-217

Eukaryotic putative RNA-binding region RNP-1 signature:

137-144

RNA recognition motif:

98-168

355/392

FIGURE 341

GCGTGGACACCACCTCAGCCCACTGAGCAGGAGTCACAGCACGAAGACCAAGCGCAAAGC
GACCCCTGCCCTCCATCCTGACTGCTCCTCCTAAGAGAGATGGCACCGGCCAGAGCAGGA
TTCTGCCCCCTTCTGCTGCTTCTGCTGCTGGGGTGTGGGTGGCAGAGATCCCAGTCAGT
GCCAAGCCCAAGGGCATGACCTCATCACAGTGGTTTAAAATTTCAGCACATGCAGCCCAGC
CCTCAAGCATGCAACTCAGCCATGAAAAACATTAACAAGCACACAAAACGGTGCAAAGAC
CTCAACACCTTCCTGCACGAGCCTTTCTCCAGTGTGGCCGCCACCTGCCAGACCCCCAAA
ATAGCCTGCAAGAATGGCGATAAAAACTGCCACCAGAGCCACGGGCCCCGTGTCCCTGACC
ATGTGTAAGCTCACCTCAGGGAAGTATCCGAAGTGCAGGTACAAAGAGAAGCGACAGAAC
AAGTCTTACGTAGTGGCCTGTAAGCCTCCCCAGAAAAAGGACTCTCAGCAATTCCACCTG
GTTCTGTACACTTGGACAGAGTCCTTTAGGTTTCCAGACTGGCTTGCTCTTTGGCTGAC
CTTCAATTCCCTCTCCAGGACTCCGCACCACTCCCCTACACCCAGAGCATTCTCTTCCCC
TCATCTCTTGGGGCTGTTCTGGTTCAGCCTCTGCTGGGAGGCTGAAGCTGACACTCTGG
TGAGCTGAGCTCTAGAGGGATGGCTTTTTCATCTTTTGTGCTGTTTTCCAGATGCTTA
TCCCCAAGAAACAGCAAGCTCAGGTCTGTGGGTTCCTGGTCTATGCCATTGCACATGTC
TCCCCTGCCCCCTGGCATTAGGGCAGCATGACAAGGAGAGGAAATAAATGGAAAGGGGGC
AA
AAA

356/392

FIGURE 342

MAPARAGFCPLLLLLLLGLWVAEIPVSAKPKGMTSSQWFKIQHMQPSPQACNSAMKNINK
HTKRCKDLNTFLHEPFSSVAATCQTPKIACKNGDKNCHQSHGPVSLTMCKLTSGKYPNCR
YKEKRQNKSYVVACKPPQKKDSQQFHLVPVHLDRVL

Important features of the protein**Signal peptide:**

1-22

Transmembrane domain:

none

N-glycosylation site:

127-131

cAMP- and cGMP-dependent protein kinase phosphorylation site:

139-143

N-myristoylation site:

18-24, 32-38

Pancreatic ribonuclease family signature:

65-72

Pancreatic ribonuclease family proteins:

49-93

357/392

FIGURE 343

GCATTTGCCACTGGTTGCAGATCAGGCGGACGAGGAGCCGGGAAGGCAGAGCCATGTGGC
TGCCCCCTGCTCTGCTCCTTCTCAGCCTCTCAGGCTGTTTCTCCATCCAAGGCCAGAGT
CTGTGAGAGCCCCAGAGCAGGGGTCCCTGACGGTTCATGCCACTATAAGCAAGGATGGG
AGACCTACATTAAGTGGTGGTGCCGAGGGGTGCGCTGGGATACATGCAAGATCCTCATTTG
AAACCAGAGGGTTCGGAGCAAGGAGAGAAGAGTGACCGTGTGTCCATCAAGGACAATCAGA
AAGACCGCACGTTCACTGTGACCATGGAGGGGCTCAGGCGAGATGACGCAGATGTTTACT
GGTGTGGGATTGAAAGAAGAGGACCTGACCTTGGGACTCAAGTGAAAGTGATCGTTGACC
CAGAGGGAGCGGCTTCCACAACAGCAAGCTCACCTACCAACAGCAATATGGCAGTGTTCA
TCGGCTCCCAAGAGGAACCACTACATGCTCCTGGTATTTGTGAAGGTGCCCATCTTGC
TCATCTTGGTCACTGCCATCCTCTGGTTGAAGGGGTCTCAGAGGGTCCCTGAGGAGCCAG
GGGAACAGCCTATCTACATGAACTTCTCCGAACCTCTGACTAAAGACATGGCCACTTAGA
GAGATGGATCTGCAGAGCCTTCCCTGCCCTGGCCACGTTTCCAGAAGAGACTCGGGCTGTG
GAAGGAACATCTACGAGTCCTCGGGATGCAGTGACTGAGATAGGGGGCCCTGGGCCTCCGC
CCTGGCCTTGGAGCTGGTGGGCACCTCCCTGTTCTGCACAGCTCAGGGACTTAGCCAGGT
CCTCTCCTGAGCCACCATCACCTCCTGGGGTGCCAGCACCTGTTCTCTTGGTCAGGAGCT
GTAGAGATGGAGCTCAAGCACTGGACGACTCTGTCCCCACTGCTGGAATAACTCGGGCAC
AGAGCATGGGACCAAGTACAGAAAGAGGTTGGGGGAGACCCCCCAGCCCTAGACTTCC
ATCATTCGGAGACCAACTCAACACCGTCTTTGCTGAGAACCTGATATATCCGTGTTTTT
TAAATTTTTTTTTTTCTAGCAAAGTTGGGTTTTAATGACTTATGTTTCATAGGAAACCTCT
CTGATCCACACACAAGGAGGGTGATCTGGGATGAGTTCCTGCTTCTAGGGCATGAGGG
GCTGGATGGACCCTGTCCCCAGGGAGGACATGGCTCTGAGTCCACAGGGCTGAGGAGCA
ATGGGAACCTCCCTGGCCCCGGCCCGGTGCTTGTCTCCCCCTCCACCTCTTCTCCTCC
TAGCTCCCCAAGCTCCCTGCCTATTCCTCCCAAGAGCCTGTGAGGCCTCAAGAACCACCT
CTCAGCATGACAGCTTGGGTCTCCTCCCCAAAAGAGCCTGTGAGGCCTCAAGAACCACCT
CCAGGTGGGGAGGGCAGTAACGAAAACCATCGCAGGAAATGGCACCCCTCCCTTTTCGGTG
ATGTTGAAATCATGTTACTAATGAAAACCTGTCTTAGGGAAGTGGTCTGTCTCCTCACAG
GCTTCACCCACGGCGATGAGGCCCTTGAATGTGGTCACTTTGTGCTGTATGGTTGAGGGA
CCCTCACACCAAAGGGACCTTCCCATGTGAGATGTGCTCCCGCCCCACCTGCCCCACAAG
CAAACACACCACACATGTTTCGGCATGTTGCCCTTTGAACACCCATGAGGACGCCTCCAAC
CTGCTCTTGGTTCTAATAGGGAGTACTGACTGTGAGCAGTGGATAAAGGAGAGGGGACCC
TCTGGTCCCTAGCATGGCACCCAGAGCCTCCCTCTTCTTGTCTTCTCAGCCAAAGAGAAA
CTTTCTCTGACTTTGAACTGAATTTAGGTCTCTGGCCAATGATGGGCCTGAAAATTCAT
AATGGCCAGAGAGGAGAGTTCGAGCCCGGCTAAGATCCCCTGAGTCATTCTGTGAGGGAC
CAAGACCCACAGTCCACCAGCCCCAGGGCCCTACCTCCTGGAATGCTTTCCTGGATCCAG
CTTCCCGAAGATCCGACCAGACCCAGGGAGGACGGCACCGCTCCGCGGGAGGGAAAGCCA
AAGCATGGTGCTTCACCAGCTGGACTCAGGGGCGAGGGGACATGGGCGCTTGTCAACGTG
ATGTCATTCTTTTCCACCGTTTCTTCTGTTGATATTCAATGAATCCGTCAATCTCTCT
GGGAAA

358/392

FIGURE 344

MWLPPALLLLSLSGCFSSIQGPESVRAPEQGS�TVQCHYKQGWETIYIKWWCRGVRWDTCKI
LIETRGSEQGEKSDRVSIKDNQKDRTFTVTMEGLRRDDADVWCGIERRGPDLTQVKVI
VDPEGAASTTASSPTNSNMAVFIGSHKRNHYMLLVFVKVPILLILVTAILWLKGSQRVPE
EPGEQPIYMNFSPLTKDMAT

Important features of the protein:

Signal peptide:

Amino acids 1-17

Transmembrane domain:

Amino acids 151-170

N-glycosylation site:

Amino acids 190-194

Tyrosine kinase phosphorylation site:

Amino acids 95-103

N-myristoylation sites:

Amino acids 66-72; 125-131

Prokaryotic membrane lipoprotein lipid attachment site:

Amino acids 5-16

359/392

FIGURE 345

CTGAGCTCCCGGGCTCCGGCAGCGCGCTGGCGGGGCGCCGCATTGCACACTCTGGGGGCG
CCGCAGTGTTCGTGGGATGGGGCAGCGGGCTGCAGCTGGCGGCCGGAATCCGCGCGCAGC
CCGGGTGCAAGTTCTCTCCTGTTGCCCTGAGTGCCCACTCCAGGCCCTCTGTATGAGTG
ACACTTCAGTCTGCCATGGAACCTGGCCCTGCTCTGGCCTGGCTCCTGCTCCTGAGCCTG
CTGGCGGATTGTCTGAAAGCTGCTCAGTCCCGAGACTTCACAGTGAAAGACATTATCTAC
CTCCATCCTTCAACCACACCATATCCTGGTGGATTAAATGTTTACCTGTGAAAAGGCA
GCAGACAATTATGAGTGCAACCGATGGGCTCCAGACATCTACTGCCCTCGAGAGACCAGA
TACTGCTACACTCAGCACACAATGGAAGTCACAGGAAACAGTATCTCAGTCACCAAACGC
TGTGTCCCCTGGAAGAGTGCTTATCCACTGGCTGCAGAGACTCCGAGCATGAAGGCCAC
AAGGTCTGCACTTCTTGTGTGAAGGAAATATCTGTAACCTGCCACTGCCCCGAAATGAA
ACTGATGCCACATTTGCCACGACGTACCTATAAATCAGACAAATGGGCACCCACGCTGT
ATGTCAGTGATAGTGTCTGCTTGTGGTTGTGGTTAGGGCTCATGTTATAGTGGCTCAGT
GGCTCCATGTGTAAATAGCGATCCATGGGGATCTCGATGGTCCACAGACCTGCATGAGTC
ATTGGCCTGACAGTAATTACACATGTGAGACACAACACTCTTGGAGGTCAACAGCCAA
GCATTGCCACTTACCATGAGGAATAAATGTTGCTTCATTGTAGCCATTTTGTGCTAACC
GAGACTCATCAAAGCCTTCTGTCAGTACAGCCCAAGTTCCATACCATAAACGTTTGT
CATTCCAAGAAGTAGTTCTGCATTTATCGAGATCTGGGGTCTTAATTTGGAAGAATACA
TGCATGAGATGCAGTAGGTCTGAGACTGTAAGATATTAGGAGTATGTTATAGGGGCATG
TATAGATGTGGGCTTTTTCAGGAGAAAAGTAACCATTGGTTTAAATATAATCATGAGTTCA
TTTGTAGCTTTAGAAATTTTAAAACATTGACTCCAACTGAATGGACTATTTCTTGGAAA
TTCTGACTGAGTCCCTGGAAGAGTAGTAATTCCAACAATTCAGCCATTTGTTCATTA
TTTTCCCAACATTCCTCTCCAGTGCTGGGAATCACATTTCTCTGTTCTGTGCAGAAGA
CAAAAAGGCAATCATAAAAGTTTGTATATTTGTGGGGGTGCCTGGAGGAGGATTTTCT
CAACTTAATGGAGCCACTGTCCATAAAGTGGCTGTTATCCCTTCATATAATTGGTGAGAT
CAGCCTTCTCCTTGACTTGGCACCTAATTATGCTTCATGAGATCCTAGATTCCACCTGAG
TCAATTGTGTCCAGAGCCCCAAACCAGGATGGAGTTGTTTCCCAGATATGGGGTTCTA
TTCAGCCATAGATAATCTAGACAGAGGATTTCAGAATGAAAGGAAAAATGTGTGGAGATT
AGTCCTAGTTTCACTCTGAGGGCCGACTAAGTGGCTCAGCCAGCTTCTTACTCCATCTGCA
GTTTCACTACTGCCAAAGAGCTCCCACTTCCAAATCCCAGTGACTTTATGGAGAAGATTCT
GCATTAAATTGTCTTTCGAATGATGGGGAAGCAAGGCATAATATGCGATGATGAGGAGAA
AGTAGACCAGTGAGGTGATTGCAAGACTAACAAGGAGACTCAATGGGAAGTTTTTCTTTC
TTTTAGATATTGCTTTTGAAGTAGATGGTAAAATTTTTGTCATCCTTCTTGTATTTTTTG
TACCCCAAGTTACAATTTTTCTTCTTCTTGTAAATAATTTAAACAGTATTTATTTTTGT
AAGGCATAACTAGAACTAAAATATATTCTAAAAAATTCATTATTCTGAACAAAGTGATC
AAATTAGAATACATATTTTTCAACAGTGGTAGAGCTTTTAATATATGTTTATTGAAAGTT
ATCTATAATACTTGCACCAAGTGTGAAAAAGTTAACATGTAGGCAAGAGCAATATGTTT
GTCTCAAGGATTTTTCCATGGTTTCTCAGTGATGGTGTCTGGAATTATTCAGGTGGTG
ACCATCACTGGTCTAAGTTTGTGTGCAGGGTTTTTCAGACGTGTTTTTGTGAACTTGGTA
GAACCATGGCTAATAAAGAGGACAGTGTGTGAGGGTCCATCTGCCCTCCATAGAAAAAT
GTCTCTGGCTCATAAAATGAGACTCCCTCAGGGACTAAATATGAACTGACAGCAGTAACT
CTGATACAGAATAATCTAAATTGCATCAAATGGCCTTAATTCAGAGTTTGTAGGCTTAT
CAGTATGTTGCTTTTAATTGGGGTGGGAAAGTAGAGGGAGAGAAAGCAAGACATTTATTA
AGCACCTCGTATGTGCCAGGCACTATGCTAAGCACTTTACATAAGTTAGGATTAATCCCT
GCAAGAACTCTATAAAGAATGTTACTAGCATTTACACTTCCCAAATGAAGGTACCAAAGC
TCAAACGCAATGTTGTGAAGCTGTTTCTTTCAGATTTAGGTTATGTGGGATGATGTGGGA
TTGAAGAGGAAAGAAAGGTGGGATTATCCCCCTAGGAAGACTTTCAGGCCTGACTTCATA
GGAATTCATCCATCTTATCATGTGGAGTTTATCTCACCCCTGCTGTTGCAGGATGCTATTT
GCATGTGTCCCAGGTGATGTTTTTCTTTGGGGAGTAGGGGTTTGGCTTCTCATTTCAT

360/392

CCCTCTTGCTAAAGAGGAGATAGTTGATGTTGCATCTAAAGATGCTATAAGACAATGAAAGTTTGATGTTGTAC
ATACCTACAAGTACCATTTTTTGTGCATGATTACACTCCACTGACATCTTCCAAGTACTAC
ATGTGATTGAATAAGAAACAAGAAAGTGACCACACCAAAGCCTCCCTGGCTGGTGTACAG
GGATCAGGTCCACAGTGGTGCAGATTCAACCACCACCCAGGGAGTGCTTGCAGACTCTGC
ATAGATGTTGCTGCATGCGTCCCATGTGCCTGTCAGAATGGCAGTGTTTAATTCTCTTGA
AAGAAAGTTATTTGCTCACTATCCCCAGCCTCAAGGAGCCAAGGAAGAGTCATTCACATG
GAAGGTCCGGGACTGGTCAGCCACTCTGACTTTTCTACCACATTAAATTCTCCATTACAT
CTCACTATTGGTAATGGCTTAAGTGTAAGAGCCATGATGTGTATATTAAGCTATGTGCC
ACATATTTATTTTTAGACTCTCCACAGCATTGATGTCAATATGGGATTAATGCCTAAACT
TTGTAAATATTGTACAGTTTGTAATCAATGAATAAAGGTTTTGAGTGTAATAAAAAAAAAA
AAAAAA

361/392

FIGURE 346

MEPGPALAWLLLLLLADCLKAAQSRDFTVKDIIYLHPSTTPYPGGFKCFTCEKAADNYE
CNRWAPDIYCPRETRYCYTQHTMEVTGNSISVTKRCVPLEECLSTGCRDSEHEGHKVCTS
CCEGNICNLPLPRNETDATFATTSPINQTNHPRCMSVIVSCLWLWLGLML

Important features of the protein:

Signal peptide:

1-22

Transmembrane domain:

None

N-glycosylation site:

134-138, 147-151

N-myristoylation site:

45-51, 87-93, 106-112, 124-130

Ly-6 / u-PAR domain protein:

115-128

362/392

FIGURE 347

GATCAAGCGCCTTCCTTTCCCTTCCTCTCCCTACTTGGCCTTTGCCCTAAGCCAAGACCT
GGCCATCAGCCTGGCTGCAGGGGCTGCAGAGCCAGCTGCACCTTTTTCAGGTATGGGGGA
GGGCCAGGCACCATGAAGCCAGTGTGGGTGCGCCACCCTTCTGTGGATGCTACTGCTGGTG
CCCAGGCTGGGGGCCGCCCCGGAAGGGGTCCCAGAAGAGGCCTCCTTCTACTATGGAACC
TTCCCTCCTGGCTTCTCCTGGGGCGTGGGCAGTTCCTGCCTACCAGACGGAGGGCGCCTGG
GACCAGGACGGGAAAGGGCCTAGCATCTGGGACGTCCTTACACACAGTGGGAAGGGGAAA
GTGCTTGGGAATGAGACGGCAGATGTAGCCTGTGACGGCTACTACAAGGTCCAGGAGGAC
ATCATTCTGCTGAGGGAAGTGCACGTCAACCACTACCGATTCTCCCTGTCTTGGCCCCGG
CTCCTGCCCACAGGCATCCGAGCCGAGCAGGTGAACAAGAAGGGAATCGAATTCTACAGT
GATCTTATCGATGCCCTTCTGAGCAGCAACATCACTCCCATCGTGACCTTGCACTACTGG
GATCTGCCACAGCTGCTCCAGGTCAAATACGGTGGGTGGCAGAATGTGAGCATGGCCAAC
TACTTCAGAGACTACGCCAACCTGTGCTTTGAGGCCTTTGGGGACCGTGTGAAGCACTGG
ATCACGTTCAGTGATCCTCGGGCAATGGCAGAAAAAGGCTATGAGACGGGGCCACCATGCG
CCGGGCCTGAAGCTCCGCGGCACCGGCCTGTACAAGGCAGCACACCACATCATTAAGGCC
CACGCCAAAACCTGGCATTCTTATAACACCACGTGGCGCAGCAAGCAGCAAGGTCTGGTG
GGAATTTCACTGAAGTGTGACTGGGGGGAACCTGTGGACATTAGTAACCCCAAGGACCTA
GAGGCTGCCGAGAGATACCTACAGTTCTGTCTGGGCTGGTTTGCCAACCCCATTTATGCC
GGTGACTACCCCAAGTCATGAAGGACTACATTGGAAGAAAGAGTGCAGAGCAAGGCCTG
GAGATGTCGAGGTTACCGGTGTTCTCACTCCAGGAGAAGAGCTACATTAAAGGCACATCC
GATTTCTTGGGATTAGGTCATTTTACTACTCGGTACATCACGGAAGGAACTACCCCTCC
CGCCAGGGGGCCAGCTACCAGAACGATCGTGACTTGATAGAGCTGGTTGACCCAACTGG
CCAGATCTGGGGTCTAAATGGCTATATTCTGTGCCATGGGGATTTAGGAGGCTCCTTAAC
TTTGCTCAGACTCAATACGGTGATCCTCCCATATATGTGATGGAAAATGGAGCATCTCAA
AAATTCCACTGTACTCAATTATGTGATGAGTGGAGAATTCAATACCTTAAAGGATACATA
AATGAAATGCTAAAAGCTATAAAAGATGGTGCTAATATAAAGGGGTATACTTCCTGGTCT
CTGTTGGATAAGTTTGAATGGGAGAAAGGATACTCAGATAGATATGGATTCTACTATGTT
GAATTTAACGACAGAAATAAGCCTCGCTATCCAAAGGCTTCAGTTCAATATTACAAGAAG
ATTATCATTGCCAATGGGTTTCCCAATCCAAGAGAGGTGGAAAGTTGGTACCTCAAAGCT
TTGGAAACTTGCTCTATCAACAATCAGATGCTTGCTGCAGAGCCTTTGCTAAGTCACATG
CAAATGGTTACGGAGATCGTGGTACCCACTGTCTGCTCCCTCTGTGCTCATCACTGCT
GTTCTACTAATGCTCCTCCTGAGGAGGCAGAGCTTGAGACAGGATTATCAATTTTGGAGCT
TCATAAGAGAATCTTCAGGATCTTCCTCCCTTTTCTGCTTTGAGGGTTTCCATACATTGC
TGTTTTTCAGGTTCTACAATAATTACCTTTTTTTCTCTTTCTCTTTTGGCTTGTGCTGGG
ATTTAAGAATTAGAAAATAAAAATAAGCAGAAATTA

363/392

FIGURE 348

MKPVWVATLLWMLLLVPRLGAARKGSPEEASFYYGTFPLGFSWGVGSSAYQTEGAWDQDG
KGPSIWDVFTHSKGKVLGNETADVACDGYKVEDIILLRELHVNHYRFSLSWPRLLPT
GIRAEQVNKKGIEFYSDLIDALLSSNITPIVTLHHWDLPQLLQVKYGGWQNVSMANYFRD
YANLCFEAFGDRVKHWITFSDPRAMAEGYETGHHAPGLKLRGTGLYKAAHHIIKAHAKT
WHSYNTTWRSKQQGLVGISLNCDWGEPVDISNPKDLEAAERYLQFCLGWFANPIYAGDYP
QVMKDYIGRKSAREQGLEMSRLPVFSLQEKSYIKGTSDFLGLGHFTTRYITERNYPSRQGP
SYQNRDLIELVDPNWPDLGSKWLYSVPWGFRRLLNFAQTQYGDPIYVMENGASQKFHC
TQLCDEWRIQYLKGYINEMLKAIKDGANIKGYTSWSLLDKFEWEKGYSDRYGFYYVEFND
RNKPRYPKASVQYYKKIIANGFPNPREVESWYLKALETCSINNQMLAAEPLLSHMQMVT
EIVVPTVCSLCVLITAVLLMLLLRRQS

Important features:**Signal peptide:**

amino acids 1-21

Transmembrane domain:

amino acids 541-558

N-glycosylation sites:

amino acids 80-84, 171-175, 245-249

Glycosaminoglycan attachment site:

amino acids 72-76

cAMP- and cGMP-dependent protein kinase phosphorylation sites:

amino acids 23-27, 564-568

Tyrosine kinase phosphorylation sites:

amino acids 203-211, 347-355, 460-468, 507-514

N-myristoylation sites:

amino acids 44-50, 79-85, 167-173, 225-231, 257-263, 315-321

Amidation site:

amino acids 307-311

Glycosyl hydrolases family 1 active site:

amino acids 407-416

Glycosyl hydrolases family 1 N-terminal signature:

amino acids 41-56

Motif name Glycosyl hydrolases family:

amino acids 37- 67

364/392

FIGURE 349

CGCAAAGCCGCCCTCGGGGCGCTCATGGCGGGACGCCTCCTGGGAAAGGCTTTAGCCGCG
GTGTCTCTCTCTCTGGCCTTGGCCTCTGTGACTATCAGGTCCTCGCGCTGCCGCGGCATC
CAGGCGTTTCAGAACTCGTTTTTCATCTTCTTGGTTTTTCATCTTAATACCAACGTCATGTCT
GGTTCTAATGGTTCCAAAGAAAATTCTCACAATAAGGCTCGGACGTCTCCTTACCCAGGT
TCAAAAGTTGAACGAAGCCAGGTTCTAATGAGAAAAGTGGGCTGGCTTGTTGAGTGGCAA
GACTATAAGCCTGTGGAATACACTGCAGTCTCTGTCTTGGCTGGACCCAGGTGGGCAGAT
CCTCAGATCAGTGAAAGTAATTTTTCTCCCAAGTTTAACGAAAAGGATGGGCATGTTGAG
AGAAAGAGCAAGAATGGCCTGTATGAGATTGAAAATGGAAGACCGAGAAATCCTGCAGGA
CGGACTGGACTGGTGGGCCGGGGGCTTTTGGGGCGATGGGGCCCAAATCACGCTGCAGAT
CCCATTATAACCAGATGGAAAAGGGATAGCAGTGGAAATAAAATCATGCATCCTGTTTTCT
GGGAAGCATATCTTACAATTTGTTGCAATAAAAAGGAAAGACTGTGGAGAATGGGCAATC
CCAGGGGGGATGGTGGATCCAGGAGAGAAGATTAGTGCCCACTGAAAAGAGAATTTGGT
GAGGAAGCTCTCAACTCCTTACAGAAAACCAGTGCTGAGAAGAGAGAAATAGAGGAAAAG
TTGCACAACTCTTCAGCCAAGACCACCTAGTGATATATAAGGGATATGTTGATGATCCT
CGAAACACTGATAATGCATGGATGGAGACAGAAGCTGTGAACTACCATGACGAAACAGGT
GAGATAATGGATAATCTTATGCTAGAAGCTGGAGATGATGCTGGAAAAGTGAAATGGGTG
GACATCAATGATAAACTGAAGCTTTATGCCAGTCACTCTCAATTCATCAAACCTTGTGGCT
GAGAAACGAGATGCACACTGGAGCGAGGACTCTGAAGCTGACTGCCATGCGTTGTAGCTG
ATGGTCTCCGTGTAAGCCAAAGGCCACAGAGGAGCATATACTGAAAAGAAGGCAGTATC
ACAGAATTTATACTATAAAAAGGGCAGGGTAGGCCACTTGGCCTATTTACTTTCAAAACA
ATTTGCATTTAGAGTGTTTTCGCATCAGAATAACATGAGTAAGATGAACTGGAACACAAAA
TTTTTCAGCTCTTTGGTCAAAAGGAATATAAGTAATCATATTTTGTATGTATTCGATTTAA
GCATGGCTTAAATTAAATTTAAACAATAATGCTCTTTGAAGAATCATAATCAGAATAAA
GATAAATTCTTGATCAGCTATA

365/392

FIGURE 350

MAGRLLGKALAAVSLSLALASVTIRSSRCRGIQAFRNSFSSSWFHLNTNVMMSGNNGSKEN
SHNKARTSPYPGSKVERSQVPNEKVGWLVWQDYKPVEYTAVSVLAGPRWADPQISESNF
SPKFNEKDGHVERKSKNGLYEIENGRPRNPAGRTGLVGRGLLGRWGNHAADPIITRWKR
DSSGNKIMHPVSGKHILQFVAIKRKDCGEWAI PGGMVDPGEKISATLKREFGEEALNSLQ
KTSAEKREIEEKLHKLFSQDHLVIYKGYVDDPRNTDNAWMETEAVNYHDETGEIMDNLM
EAGDDAGKVKWVDINDKLKLYASHSQFIKLVAEKRDAHWSDEDSEADCHAL

Important features of the protein:**Signal peptide:**

1-20

Transmembrane domain:

None

N-glycosylation site:

55-59

cAMP- and cGMP-dependent protein kinase phosphorylation site:

179-183

N-myristoylation site:

53-59, 56-62

mutT domain signature:

215-235

366/392

FIGURE 351

CCTCTGTCTGTGCTCCCATCCCAGGGAGTATAGGTGGAGCCTCCAGAGCCCATGGACAGG
GCATGCTGGGGCTGGGCCAGCCCCAGCGGTGTCTCTAAGGCACCCCTGGGATCCCCACTG
AGCTGGCCTACTTCAGACAGCCAGGGCCCCACCCCTCTGGCCCCCTTAGTGTCCAGCTCGT
GGCCCCCTTGGCATTTCCACAAGACGCCAAGATGGAGATTCCCATGGGGACCCAGGGCTGC
TTCTCAAAGAGCCTCCTGCTCTCAGCCTCAATCCTGGTCCTCTGGATGCTCCAAGGCTCC
CAGGCAGCTCTCTACATCCAGAAGATTCCAGAGCAGCCTCAAAAGAACCAGGACCTTCTC
CTGTCAGTCCAGGGTGTCCCAGACACCTTCCAGGACTTCAACTGGTACCTGGGGGAGGAG
ACGTACGGAGGCACGAGGCTATTTACCTACATCCCTGGGATACAACGGCCTCAGAGGGAT
GGCAGTGCCATGGGACAGCGAGACATCGTGGGCTTCCCCAATGGTTCCATGCTGCTGCGC
CGCGCCCAGCCTACAGACAGTGGCACCTACCAAGTAGCCATTACCATCAACTCTGAATGG
ACTATGAAGGCCAAGACTGAGGTCCAGGTAGCTGAAAAGAATAAGGAGCTGCCCAGTACA
CACCTGCCCACCAACGCTGGGATCCTGGCGGCCACCATCATTTGGATCTCTTGCTGCCGGG
GCCCTTCTCATCAGCTGCATTGCCATCTCTCTGGTGACAAGGAACTGGAGGGGCCAGAGC
CACAGACTGCCTGCTCCGAGGGGCCAGGGATCTCTGTCCATCTTGTGCTCGGCTGTATCC
CCAGTGCCTTCAGTGACGCCCAGCACATGGATGGCGACCACAGAGAAGCCAGAATTGGGC
CCTGCTCATGATGCTGGTGACAACAACATCTATGAAGTGATGCCCTCTCCAGTCTCTTG
GTGTCCCCCATCAGTGACACAAGGTCCATAAACCCAGCCCGGCCCTGCCACACCCCCA
CACCTGCAGGCGGAGCCAGAGAACCACCAGTACCAGCAGGACCTGCTAAACCCCGACCCT
GCCCCCTACTGCCAGCTGGTGCCAACTTCCTGATGGGTCTGGGCCAGGCCAGCCAGGGA
GAAGACAAGGCCCCAGCCCTCCTCTGGGAGCCTCACACCTGAGACCAGCAGGACAAGGCC
ATTGGGGGCTGTGGGGCCGATGAGGTGGACTCAGCCAAAGACTCAGCAGCACATGGGGCA
GGTGTCTTGGCAGGGGGACAGGAGACTGTAACAGGCCCCAGGTCTTGTGCAGCCCCTGAA
TGCACGCCCCGCTTTCGGTCTGTTCTTCAAGCAAGCTGGCCTGGGCCATGTGCCTGTGAA
AGGCAGGCTCTGGCCCCCTTTCCATGCCAAAGTCCCCCAAGATCTGGATATCTGGGGACAA
GATGGTGGCCTCAGGCCTGCCTCCAGGCAGTTGGCTGGGCTCCCAACTGTCTGTCTCTCA
ATGCCCTACCCCAACTCCACTAGTGACCCTCAGAGTCTTCTCCCCTTAGGACAAGGCAGA
CACCCCACCATGCGGGCCTCAGGTGGCAGAGAGGCCAGCCTCACAGGCCTGTGGCCCCA
CACACCAGTCCCAGCAAGGTGACCACGGCTGCTGGACCCCTTCCCTGTTTCAGGCAGGCCC
AGCCCCCTCTCAGAACCTGCTGCCAGCTGCTGGTCTTGGCCCCCACCCCTGAATCTTACTGA
GTCCCTCTGGGCAGCAGCTCCCTTCTCCACCCACCCAGCACCCGTCCCAAATGTGGCC
TCAGCTTGTCCTCCCCTTCCCCAACTATGCATTCAATTCAGCAATAAATGAGCCTTTGCT
GCA

367/392

FIGURE 352

MEIPMGQTQGCFSKSLLLSASILVLWMLQGSQAALYIQKIPEQPQKNQDLLLSVQGVPDF
QDFNWYLGEETYGGTRLFTYIPGIQRPQRDGSAMGQRDIVGFPNGSM LLRRAQPTDSGTY
QVAITINSEWTMKAKTEVQVAEKNKELPSTHLPTNAGILAATIIGSLAAGALLISCIAYL
LVTRNWRGQSHRLPAPRGQGSLSILCSAVSPVPSVTPSTWMATTEKPELGPAGDAGDNNI
YEVMPSPVLLVSPISDTRSINPARPLPTPHLQAEPEHQQD LLNPD PAPYCQLVPTS

Important features of the protein:**Signal peptide:**

Amino acids 1-32

Transmembrane domain:

Amino acids 159-178

N-glycosylation site:

Amino acids 104-108

N-myristoylation sites:

Amino acids 6-12; 29-35; 55-61; 91-97; 157-163; 165-171

368/392

FIGURE 353

CTTCAGAACAGGTTCTCCTTCCCCAGTCACCAGTTGCTCGAGTTAGAATTGTCTGCAATG
GCCGCCCTGCAGAAATCTGTGAGCTCTTTCCTTATGGGGACCCTGGCCACCAGCTGCCTC
CTTCTCTTGGCCCTCTTGGTACAGGGAGGAGCAGCTGCGCCCATCAGCTCCCACTGCAGG
CTTGACAAGTCCAACCTTCCAGCAGCCCTATATCACCAACCGCACCTTCATGCTGGCTAAG
GAGGCTAGCTTGGCTGATAACAACACAGACGTTTCGTCTCATTGGGGAGAACTGTTCCAC
GGAGTCAGTATGAGTGAGCGCTGCTATCTGATGAAGCAGGTGCTGAACCTTCACCCTTGAA
GAAGTGCTGTTCCCTCAATCTGATAGGTTCCAGCCTTATATGCAGGAGGTGGTGGCCCTTC
CTGGCCAGGCTCAGCAACAGGCTAAGCACATGTCATATTGAAGGTGATGACCTGCATATC
CAGAGGAATGTGCAAAAGCTGAAGGACACAGTGAAAAAGCTTGGAGAGAGTGAGAGATC
AAAGCAATTGGAGAACTGGATTTGCTGTTTATGTCTCTGAGAAATGCCTGCATTTGACCA
GAGCAAAGCTGAAAAATGAATAACTAACCCCTTTCCTGCTAGAAATAACAATTAGATG
CCCCAAAGCGATTTTTTTTAACCAAAAGGAAGATGGGAAGCCAACTCCATCATGATGGG
TGGATTCCAAATGAACCCCTGCGTTAGTTACAAAGGAAACCAATGCCACTTTTGTTTATA
AGACCAGAAGGTAGACTTTCTAAGCATAGATATTTATTGATAACATTTTCATTGTAAGTGG
TGTTCTATACACAGAAAACAATTTATTTTTTAAATAATTGTCTTTTCCATAAAAAAGAT
TACTTTCCATTCCCTTAGGGGAAAAAACCCCTAAATAGCTTCATGTTTCCATAATCAGTA
CTTTATATTTATAAATGTATTTATTATTATTATAAGACTGCATTTTATTTATATCATTTT
ATTAATATGGATTTATTTATAGAAACATCATTCGATATTGCTACTTGAGTGTAAGGCTAA
TATTGATATTTATGACAATAATTATAGAGCTATAACATGTTTATTTGACCTCAATAAACA
CTTGGATATCCC

369/392

FIGURE 354

MAALQKSVSSFLMGTLATSCLLLLLALLVQGGAAAPISSHCRLDKSNFQQPYITNRTFMLA
KEASLADNNTDVRLIGEKLFGVSMSERCYLMKQVLNFTLEEVLFPPQSDRFQPYMQEVVP
FLARLSNRLSTCHIEGDDLHIQRNVQKLKDTVKKLGESGEIKAIGELDLLFMSLRNACI

Important features of the protein:

Signal peptide:

amino acids 1-33

N-glycosylation sites:

amino acids 54-58, 68-72, 97-101

N-myristoylation sites:

amino acids 14-20, 82-88

Prokaryotic membrane lipoprotein lipid attachment site:

amino acids 10-21

370/392

FIGURE 355

TGGCCTACTGGAAAAAAAAAAAAAAAAAAAAAAAAAGTCACCCGGGCCCCGCGGTGGCCACAA
CATGGCTGCGGCGCCGGGGCTGCTCTTCTGGCTGTTTCGTGCTGGGGGCGCTCTGGTGGGT
CCCGGGCCAGTCGGATCTCAGCCACGGACGGCGTTTCTCGGACCTCAAAGTGTGCGGGGA
CGAAGAGTGCAGCATGTTAATGTACCGTGGGAAAGCTCTTGAAGACTTCACGGGCCCTGA
TTGTCGTTTTGTGAATTTTAAAAAAGGTGACGATGTATATGTCTACTACAACTGGCAGG
GGGATCCCTTGAACTTTGGGCTGGAAGTGTTGAACACAGTTTTTGGATATTTTCCAAAAGA
TTTGATCAAGGTACTTCATAAATACACGGAAGAAGAGCTACATATTCCAGCAGATGAGAC
AGACTTTGTCTGCTTTGAAGGAGGAAGAGATGATTTTAATAGTTATAATGTAGAAGAGCT
TTTAGGATCTTTGGAACCTGGAGGACTCTGTACCTGAAGAGTCGAAGAAAGCTGAAGAAGT
TTCTCAGCACAGAGAGAAATCTCCTGAGGAGTCTCGGGGGCGTGAACTTGACCCTGTGCC
TGAGCCCGAGGCATTTCAGAGCTGATTTCAGAGGATGGAGAAGGTGCTTCTCAGAGAGCAC
CGAGGGGCTGCAGGGACAGCCCTCAGCTCAGGAGAGCCACCCTCACACCAGCGGTCTTGC
GGCTAACGCTCAGGGAGTGCAGTCTTCGTTGGACACTTTTGAAGAAATTCTGCACGATAA
ATTGAAAGTGCCGGGAAGCGAAAGCAGAACTGGCAATAGTTCTCCTGCCTCGGTGGAGCG
GGAGAAGACAGATGCTTACAAAGTCCTGAAAACAGAAATGAGTCAGAGAGGAAGTGGACA
GTGCGTTATTTCATTACAGCAAAGGATTTTCGTTGGCATCAAAATCTAAGTTTGTTTTACAA
AGATTGTTTT**TA**GTACTAAGCTGCCTTGGCAGTTTGCATTTTGTAGCCAAACAAAAATAT
ATTATTTTCCCTTCTAAGTAAAAAAAAAAAAAAAAAAAAA

371/392

FIGURE 356

MAAAPGLLFWLFLVGLALWWVPGQSDLSHGRRFSDLKVCGDEECSSMLMYRGKALEDFTGPD
CRFVNFKKGDDVYVYYKLAGGSLELWAGSVEHSFGYFPKDLIKVLHKEYTEELHIPADET
DFVCFEGGRDDFNSYNVEELLGSLELEDSVPEESKKAEEVSQHREKSPPEESRGRELDVPV
EPEAFRADSEEDGEGAFSESTEGGLQGQPSAQESHPTSGPAANAQGVQSSLDTFEEILHDK
LKVPGESERTGNSSPASVEREKTDAYKVLKTEMSQRGSGQCVIHYSKGFRWHQNLSLFYK
DCF

Important features of the protein:

Signal peptide:

amino acids 1-22

N-glycosylation site:

amino acids 294-298

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 30-34

Tyrosine kinase phosphorylation site:

amino acids 67-76

N-myristoylation sites:

amino acids 205-211, 225-231, 277-283

Amidation site:

amino acids 28-32

372/392

FIGURE 357

[illegible]

373/392

FIGURE 358

MEAPGPRLRTALCGGCCCLLLCAQLAVAGKGARGFGRGALIRLNIWPAVQGACKQLEVC
EHCVEGDRARNLSSCMWEQCRPEEPGHCVAQSEVVKEGCSIYNRSEACPAHHHPTYPEPK
TVTTGSPPVPEAHSPGFDGASFIGGVVLVLSLQAVAFFVLHFLKAKDSTYQTL

Important features of the protein:

Signal peptide:

1-29

Transmembrane domain:

141-160

N-glycosylation site:

71-75, 103-107

Tyrosine kinase phosphorylation site:

164-171

N-myristoylation site:

15-21

374/392

FIGURE 359

TTCCAGTCAGAGTTAAGTTAAACAGAAAAAGGAAGATGGCAAGAATATTGTTACTTTT
CCTCCCGGGTCTTGTGGCTGTATGTGCTGTGCATGGAATATTTATGGACCGTCTAGCTTC
CAAGAAGCTCTGTGCAGATGATGAGTGTGTCTATACTATTTCTCTGGCTAGTGCTCAAGA
AGATTATAATGCCCCGGACTGTAGATTCATTAACGTTAAAAAAGGGCAGCAGATCTATGT
GTA¹CTCAAAGCTGGTAAAAGAAAATGGAGCTGGAGAATTTTGGGCTGGCAGTGTTTATGG
TGATGGCCAGGACGAGATGGGAGTCGTGGGTATTTCCCCAGGAACCTGGTCAAGGAACA
GCGTGTGTACCAGGAAGCTACCAAGGAAGTTCCCACCACGGATATTGACTTCTTCTGCGA
GTAATAAATTAGTTAA²AACTGCAAATAGAAAGAAAACACCAAAAATAAAGAAAAGAGCAA
AAGTGGCCAAAAAATGCATGTCTGTAATTTTGGACTGACGT

375/392

FIGURE 360

MARILLFLPGLVAVCAVHGIFMDRLASKKLCADDECVTISLASAQEDYNAPDCRFINV
KKGQQIYVYSKLVKENGAGEFWAGSVYGDGQDEMGVVGYPFRNLVKEQRVYQEATKEVPT
TDIDFFCE

Important features of the protein:

Signal peptide:
1-14

Transmembrane domain:
None

N-myristoylation site:
84-90

376/392

FIGURE 361

GGCACGAGCCACCACTTACAACCACACAGCCTATCCAGAAACATGAAGATAAGAAATGCT
TGTGCTGTCCCTTATTGAAGTACTCCTGTTTATACTTGAAGGAGTTACAGGAGCTCGAAAA
ATTTCAACTTTCTCAGGCCCTGGCTCATGGCCGTGCAATCCCAAGTGTGATGGCAGAACT
TACAACCCCTCAGAGGAGTGTGTGTTTCATGACACCATCCTGCCCTTTAAGCGGATTAAC
CTCTGTGGCCCTAGCTGCACCTACAGGCCCTGCTTTGAGCTCTGCTGTCTGAGTCCTAT
AGCCCCAAGAAGAAATTTATTGTCAAGCTTAAAGTTCATGGAGAGAGATCCCATTGCAGT
TCATCCCCTATCTCCAGGAAGTGTAAAAGCAACAAGATTTTTTCATGGAGAAGATATTGAA
GACAACCAACTTTCTCTTAGGAAAAAAAGTGGTGACCAGCCTTGAGAGTCTGCTTTCTTC
CTGCAAGCACCAGTTCCTGAATGTTCTTACTTGAAGAATGGATACCTGAAGCATTGGGGT
GCAGTGATATATGTGTCTCATTACAATGCTCCTTTGGATATTGTTTTCTAAGCATGTGT
TGGAATGTTCCCCCATAACTTTCTAAAATTATCCTATTTCAATGCAACTAAAGATAAATG
TATTCAGCCAGAGTCCACAGAGAAGGCAAGTTATGCAAGGCAGGCATGGGGCCCTCACA
AAATTTCAAGCTGTGCGACTTATGTAGTAATTTTCTACAAACAATCCCTCCTGGATATCC
AGGAGGCTCCAGACCTGAATAAAAACCACATGTCTGTCTAGAAAAAGGGAATGAATCAAG
ATCCACAGGACCTTTTCAAGATTTTAGAAGCAGCAAACTATGGCTGAGAGAAAAGACTCT
CTGACCAGGCAAATTGTTCTGCAGTATTCTCGGGCGTGTAGCTCCCTGAGTAGTCGCC
AGGCTGGTCTTGGCTTTGTAATAATACAGCTGCCTTTGAGTCCTCCCTACCCTGTTAGTA
ACCCCTTGCCTGCACTGTTGTCCTTACAACCGAAATAAACTGATTAGTTG

377/392

FIGURE 362

MKIRNACAVLIEVLLFILEGVTGARKISTFSGPGSWPCNPKCDGRTYNPSEECCVHDTIL
PFKRINLCGPSCTYRPCFELCCPESYSPKKKFIVKLKVHGERSHCSSSPISRNCKSNKIF
HGEDIEDNQLSLRKKSGDQP

Important features of the protein:

Signal peptide:

1-23

Transmembrane domain:

None

Glycosaminoglycan attachment site:

31-35

N-myristoylation site:

20-26, 34-40

378/392

FIGURE 363

ACACTGGCCAAACAAAAACGAAAGCACTCCGTGCTGGAAGTAGGAGGAGAGTCAGGACTC
CCAGGACAGAGAGTGCACAACTACCCAGCACAGCCCCCTCCGCCCCCTCTGGAGGCTGA
AGAGGGATTCAGCCCCCTGCCACCCACAGACACGGGCTGACTGGGGTGTCTGCCCCCTT
GGGGGGGGGAGCAGCACAGGGCCTCAGGCCTGGGTGCCACCTGGCACCTAGAAAGATGCCTGT
GCCCTGGTTCTTGCTGTCTTGGCACTGGGCCGAAGCCCAGTGGTCTTTCTCTGGAGAG
GCTTGTGGGGCCTCAGGACGCTACCCACTGCTCTCCGGGCCTCTCCTGCCGCCTCTGGGA
CAGTGACATACTCTGCCTGCCTGGGGACATCGTGCTGCTCCGGGCCCCGTGCTGGCGCC
TACGCACCTGCAGACAGAGCTGGTGCTGAGGTGCCAGAAGGAGACCGACTGTGACCTCTG
TCTGCGTGTGGCTGTCCACTTGGCCGTGCATGGGCACTGGGAAGAGCCTGAAGATGAGGA
AAAGTTTGGAGGAGCAGCTGACTCAGGGGTGGAGGAGCCTAGGAATGCCTCTCTCCAGGC
CCAAGTCGTGCTCTCCTTCCAGGCCTACCCACTGCCCCGCTGCGTCTTGCTGGAGGTGCA
AGTGCTGCTGCCCTTGTGCAGTTTGGTCAGTCTGTGGGCTCTGTGGTATATGACTGCTT
CGAGGCTGCCCTAGGGAGTGAGGTACGAATCTGGTCTTATACTCAGCCCAGGTACGAGAA
GGAACCTCAACCACACACAGCAGCTGCCTGCCCTGCCCTGGCTCAACGTGTCAGCAGATGG
TGACAACGTGCATCTGGTTCTGAATGTCTCTGAGGAGCAGCACTTCGGCCTCTCCCTGTA
CTGGAATCAGGTCCAGGGCCCCCCTAAACCCCGGTGGCACAAAAACCTGACTGGACCGCA
GATCATTACCTTGAACCACACAGACCTGGTTCCCTGCCTCTGTATTAGGTGTGGCCTCT
GGAACCTGACTCCGTTAGGACGAACATCTGCCCCCTCAGGGAGGACCCCCGCGCACACCA
GAACCTCTGGCAAGCCGCCCCGACTGCGACTGCTGACCCTGCAGAGCTGGCTGCTGGACGC
ACCGTGCTCGCTGCCCGCAGAAGCGGCACTGTGCTGGCGGGCTCCGGGTGGGGACCCCTG
CCAGCCACTGGTCCCACCGCTTTCTGGGAGAACGTCACTGTGGACAAGGTTCTCGAGTT
CCCATTGCTGAAAGGCCACCCTAACCTCTGTGTTAGGTGAACAGCTCGGAGAAGCTGCA
GCTGCAGGAGTGCTTGTGGGCTGACTCCCTGGGGCCTCTCAAAGACGATGTGCTACTGTT
GGAGACACGAGGCCCCCAGGACAACAGATCCCTCTGTGCCTTGAACCCAGTGGCTGTAC
TTCACTACCCAGCAAAGCCTCCACGAGGGCAGCTCGCCTTGGAGAGTACTTACTACAAGA
CCTGCAGTCAGGCCAGTGTCTGCAGCTATGGGACGATGACTTGGGAGCGCTATGGGCCTG
CCCCATGGACAAATACATCCACAAGCGCTGGGCCCTCGTGTGGCTGGCCTGCCTACTCTT
TGCCGCTGCGCTTTCCCTCATCCTCCTTCTCAAAAAGGATCACGCGAAAGGGTGGCTGAG
GCTCTTGAAACAGGACGTCCGCTCGGGGGCGGCCCGCCAGGGGCCGCGCGGCTCTGCTCCT
CTACTCAGCCGATGACTCGGGTTTCGAGCGCCTGGTGGGCGCCCTGGCGTCCGCCCTGTG
CCAGCTGCCGCTGCGCGTGGCCGTAGACCTGTGGAGCCGTCTGTAAGTGAAGCGCGCAGGG
GCCCCGTGGCTTGGTTTCACGCGCAGCGGCCAGACCCTGCAGGAGGGCGGCGTGGTGGT
CTTGCTCTTCTCTCCCGGTGCGGTGGCGCTGTGCAGCGAGTGGCTACAGGATGGGGTGTC
CGGGCCCCGGGGCGCACGGCCCCGCACGACGCCTTCCGCGCCTCGCTCAGCTGCGTGCTGCC
CGACTTCTTGCAAGGCCGGGCGCCCCGCGAGCTACGTGGGGGCTGCTTCGACAGGCTGCT
CCACCCGGACGCCGTACCCGCCCTTTTCCGCACCGTGCCCGTCTTCACTGCCCCTCCA
ACTGCCAGACTTCTGGGGGCCCTGCAGCAGCCTCGCGCCCCGCGTTCGGGGCGGCTCCA
AGAGAGAGCGGAGCAAGTGTCCCGGGCCCTTCAAGCCAGCCCTGGATAGCTACTTCCATCC
CCCGGGGACTCCCGCGCCGGGACGCGGGGTGGGACCAGGGGCGGGACCTGGGGCGGGGGA
CGGGACTTAAATAAAGGCAGACGCTGTTTTTCTAAAAAA

379/392

FIGURE 364

MPVPWFLLSLALGRSPVVLSELERLVGPQDATHCSPGLSCRLWDS DILCLPGDIVPAPGPV
LAPTHLQTELVLRCQKETDCDLCLRVAVHLAVHGHWEPEDEEKFGGAADSGVEEPRNAS
LQAQVVL SFQAYPTARCVLLEVQVPAALVQFGQSVGSVVYDCFEAALGSEVRIWSYTQPR
YEKELNHTQQLPALPWLNV SADGDNVHLVLNVSEEQHFGLSLYWNQVQGP PKPRWHKNLT
GPQIITLNHTDLVPCLCIQVWPLEPDSVRTNICPFREDPRAHQNLWQAARLRLTLQSWL
LDAPCSLPAAEALCWRAPGGDPCQPLVPPLSWENVTVDKVLEFP LLKGHPNLCVQVNSSE
KLQLQECLWADSLGPLKDDVLLLETRGPQDNRS LCALEPSGCTSLPSKASTRAARLGEYL
LQDLQSGQCLQLWDDDLGALWACPMKYIHKRWALVWLACLLFAAALSLILLKKDHAKG
WLRLKQDVRS GAAARGRAALLYSADDSGFERLVGALASALCQLPLRVAVDLWSRRELS
AQGPVAFWFAQRRTLQEGGVVLLFSPGAVALCSEWLQDGVSGPGAHGPHDAFRASLSC
VLPDFLQGRAPGSYVGACFDRLHPDAVPALFRTVPVFTLPSQLPDLGALQQPRAPRSG
RLQERAEQVSRALQPALDSYFHPPGTPAPGRGVGPGAGPGAGDGT

Signal sequence:

amino acids 1-20

Transmembrane domain:

amino acids 453-475

N-glycosylation sites:amino acids 118-121, 186-189, 198-201, 211-214, 238-241,
248-251, 334-337, 357-360, 391-394**Glycosaminoglycan attachment site:**

amino acids 583-586

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 552-555

N-myristoylation sites:amino acids 107-112, 152-157, 319-324, 438-443, 516-521,
612-617, 692-697, 696-701, 700-705

380/392

FIGURE 365

AATAGAAGTCCTCAGGACGGAGCAGAGGTGGCCGGCGGGCCCGGCTGACTGCGCCTCTGC
TTTCTTTCCATAACCTTTTCTTTTCGGA CT CGAATCACGGCTGCTGCGAAGGGTCTAGTTC
CGGACACTAGGGTGCCCGAACGCGCTGATGCCCCGAGTGCTCGCAGGGCTTCCCGCTAAC
CATGCTGCCGCCGCCGCGCGGCCCGCAGCTGCCTTGCGCTGCCTGTGCTCCTGCTACTGCT
GGTGGTGCTGACGCCGCCCCGACCGGCGCAAGGCCATCCCCAGGCCCAGATTACCTGCG
GCGCGGCTGGATGCGGCTGCTAGCGGAGGGCGAGGGCTGCGCTCCCTGCCGGCCAGAAGA
GTGCGCCGCGCCGCGGGGCTGCCTGGCGGGCAGGGTGCGCGACGCGTGCGGCTGCTGCTG
GGAATGCGCCAACCTCGAGGGCCAGCTCTGCGACCTGGACCCAGTGCTCACTTCTACGG
GCACTGCGGCGAGCAGCTTGAGTGCCGGCTGGACACAGGCGGCGACCTGAGCCGCGGAGA
GGTGCCGGAACCTCTGTGTGCCTGTGCTTCGCAGAGTCCGCTCTGCGGGTCCGACGGTCA
CACCTACTCCAGATCTGCCGCCTGCAGGAGGCGGCCCGCGCTCGGCCCGATGCCAACCT
CACTGTGGCACACCCGGGGCCCTGCGAATCGGGGCCCCAGATCGTGTCACATCCATATGA
CACTTGGAATGTGACAGGGCAGGATGTGATCTTTGGCTGTGAAGTGTTCCTACCCCAT
GGCCTCCATCGAGTGGAGGAAGGATGGCTTGACATCCAGCTGCCAGGGGATGACCCCCA
CATCTCTGTGCAGTTTAGGGGTGGACCCAGAGGTTTGAGGTGACTGGCTGGCTGCAGAT
CCAGGCTGTGCGTCCAGTGATGAGGGCACTTACCGCTGCCTTGGCCGCAATGCCCTGGG
TCAAGTGGAGGCCCCTGCTAGCTTGACAGTGCTCACACCTGACCAGCTGAACTCTACAGG
CATCCCCCAGCTGCGATCACTAAACCTGGTTCCCTGAGGAGGAGGCTGAGAGTGAAGAGAA
TGACGATTACTACTAGGTCCAGAGCTCTGGCCCATGGGGGTGGGTGAGCGGCTATAGTGT
TCATCCCTGCTCTTGAAAAGACCTGGAAAGGGGAGCAGGGTCCCTTCATCGACTGCTTTC
ATGCTGTCAGTAGGGATGATCATGGGAGGCCTATTTGACTCCAAGGTAGCAGTGTGGTAG
GATAGAGACAAAAGCTGGAGGAGGGTAGGGAGAGAAGCTGAGACCAGGACCGGTGGGGTA
CAAAGGGGCCCATGCAGGAGATGCCCTGGCCAGTAGGACCTCCAACAGGTTGTTTCCCAG
GCTGGGGTGGGGGCCTGAGCAGACACAGAGGTGCAGGCACCAGGATTCTCCACTTCTTCC
AGCCCTGCTGGGCCACAGTTCTAACTGCCCTTCCCTCCAGGCCCTGGTTCTTGCTATTTC
CTGGTCCCCAACGTTTATCTAGCTTGTTTGCCCTTTCCCCAAACTCATCTTCCAGAACTT
TTCCCTCTCTCCTAAGCCCCAGTTGCACCTACTAACTGCAGTCCCTTTTGCTGTCTGCCG
TCTTTTGTACAAGAGAGAGAACAGCGGAGCATGACTTAGTTCAGTGCAGAGAGATT

381/392

FIGURE 366

MLPPPRPAAALALPVLLLLLVLTTPPTGARPSGPDYLRRGWMRLLAEGEGCAPCRPEE
CAAPRGCLAGRVRDACGCCWECANLEGQLCDLPSAHFYGHCGEQLECRDGTGGDLRGE
VPEPLCACRSQSPLCGSDGHTYSQICRLQEAARARPDANLTVAHPGPCESGPQIVSHPYD
TWNVTGQDVIFGCEVFAYPMASIEWRKDGLDIQLPGDDPHISVQFRGGPQRF EVTGWLQI
QAVRPSDEGTYRCLGRNALGQVEAPASLTVLTPDQLNSTGIPQLRSLNLVPEEEAESEEN
DDYY

Important features of the protein:

Signal peptide:

1-30

Transmembrane domain:

None

N-glycosylation site:

159-163, 183-187, 277-281

Tyrosine kinase phosphorylation site:

244-252

N-myristoylation site:

52-58, 66-72, 113-119, 249-255

Kazal-type serine protease inhibitor domain:

121-168

Immunoglobulin domain:

186-255

Insulin-like growth factor binding proteins:

53-90

382/392

FIGURE 367

AGACGCTACAGGATGGAGCGGGGCGCAGGAGCCAAGCTGCTGCCGCTGCTGCTGCTTCTG
CGGGCGACTGGTTTTCACATGTGCACAGACAGATGGCCGGAACGGCTACACGGCGGTCATC
GAAGTGACCAGCGGGGGTCCCTGGGGCGACTGGGCCTGGCCTGAGATGTGTCCCGATGGA
TTCTTCGCCAGCGGGTTCTCGCTCAAGGTGGAGCCTCCCCAAGGCATTCTGGCGACGAC
ACTGCACTGAATGGGATCAGGCTGCACTGCGCGCGCGGGAACGTCCTAGGCAATACGCAC
GTGGTAGAGTCCAGTCTGGAAGCTGGGGCGAATGGAGTGAGCCGCTGTGGTGTGCGGGC
GGCGCCTACCTAGTGGCTTTCTCGCTTCGCGTGGAGGCACCCACGACCCTCGGTGACAAC
ACAGCAGCGAACAACGTGCGCTTCCGCTGTTCAGACGGCGAGGAACTGCAGGGGCCTGGG
CTGAGCTGGGGAGACTTTGGAGACTGGAGTGACCATTGCCCCAAGGGCGCGTGCGGCCTG
CAGACCAAGATCCAGGGACCTAGAGGCCTCGGCGATGACACTGCGCTGAACGACGCGCGC
TTATTCTGCTGCCGCAGTGAACGGCGCCGCCGCCGCGCTCTCTCCCGGGCCAGGAGGC
TAGTCCCACCTCTTGCTATTAAAGCTTCTCTGAGTTG

383/392

FIGURE 368

MERGAGAKLLPLLLLLLRATGFTCAQTDGRNGYTAVIEVTSGGPWGDWAWPEMCPDGFFAS
GFSLKVEPPQGI PGDDTALNGIRLHCARGNVLGNTHVVESQSGSWGGEWSEPLWCRGGAYL
VAFSLRVEAPTTLGDNNTAANNVRFRCS DGEELQGPGLSWGDFGDWSDHCPKGACGLQTKI
QGPRGLGDDTALNDARLFCCRS

Important features of the protein:

Signal peptide:

1-24

Transmembrane domain:

None

N-myristoylation site:

41-47

89-95

156-162

Growth factor and cytokines receptors family signature 2:

103-110

384/392

FIGURE 369

GCCAACTGGCCAAACCTCGGAGACCGTCCTGCGCTCTCTGGAGACGCGCTGTCCGCGC
CCAGGGTGGTGCCATGTGGGGCGCTCGCCGCTCGTCCGTCTCCTCATCCTGGAACGCCGC
TTCGCTCCTGCAGCTGCTGCTGGCTGCGCTGCTGGCGGCGGGGGCGAGGGCCAGCGGCGA
GTACTGCCACGGCTGGCTGGACGCGCAGGGCGTCTGGCGCATCGGCTTCCAGTGTCCCGA
GCGCTTCGACGGCGGCGACGCCACCATCTGCTGCGGCAGCTGCGCGTTGCGCTACTGCTG
CTCCAGCGCCGAGGCGCGCCTGGACCAGGGCGGCTGCGACAATGACCGCCAGCAGGGCGC
TGGCGAGCCTGGCCGGGCGGACAAAGACGGCCCCGACGGCTCGGCAGTGCCCATCTACGT
GCCGTTCTCATTGTTGGCTCCGTGTTTGTGCGCTTTATCATCTTGGGGTCCCTGGTGGC
AGCCTGTTGCTGCAGATGTCTCCGGCCTAAGCAGGATCCCCAGCAGAGCCGAGCCCCAGG
GGGTAAACCGCTTGATGGAGACCATCCCCATGATCCCCAGTGCCAGCACCTCCCGGGGGTC
GTCCTCACGCCAGTCCAGCACAGCTGCCAGTTCAGCTCCAGCGCCAACTCAGGGGCCCCG
GGCGCCCCCAACAAGGTCACAGACCAACTGTTGCTTGCCGGAAGGGACCATGAACAACGT
GTATGTCAACATGCCCACGAATTTCTCTGTGCTGAACTGTCAGCAGGCCACCCAGATTGT
GCCACATCAAGGGCAGTATCTGCATCCCCATACGTGGGGTACACGGTGCAGCACGACTC
TGTGCCCATGACAGCTGTGCCACCTTTCATGGACGGCCTGCAGCCTGGCTACAGGCAGAT
TCAGTCCCCCTTCCCTCACACCAACAGTGAACAGAAGATGTACCCAGCGGTGACTGTATA
ACCGAGAGTCACTGGTGGGTTCCTTTACTGAAGGGAGACGAAGGCAGGGGTGGATTTTCG
AGGTGGAAGT

385/392

FIGURE 370

MWGARRSSVSSSWNAASLLQLLLAALLAAGARASGEYCHGWLDAQGVWRIGFQCPERFDG
GDATICCGSCALRYCCSSAEARLDQGGCDNDRQQGAGEPGRADKDGPDGSAPVIYVPFLI
VGSVFVAFIILGSLVAACCCRCLRPKQDPQQSRAPGGNRLMETIPMIPSASTSRGSSSRQ
SSTAASSSSSANS GARAPPTRSQTNCCLPEGTMMNVYVNMPTNFSVLNCQQATQIVPHQG
QYLHPPYVGyTVQHDSVPMTAVPPFMDGLQPGYRQIQSPFPHTNSEQKMPAVTV

Important features of the protein:**Signal peptide:**

1-33

Transmembrane domain:

54-78

N-glycosylation site:

223-226

cAMP- and cGMP-dependent protein kinase phosphorylation site:

5-8

N-myristoylation site:

3-8, 30-35, 60-65, 86-91, 132-137, 211-216, 268-273

Prokaryotic membrane lipoprotein lipid attachment site:

128-138

386/392

FIGURE 371

CACCAGACAGCACTCCAGCACTCTGTTTGGGGGGCATTGAAACAGCAAAATCACTCATA
AAAGGCCAAAAAATTGCAAAAAAATAGTAATAACCAGCATGGCACTAAATAGACCATGA
AAAGACATGTGTGTGCAGTATGAAAATTGAGACAGGAAGGCAGAGTGTCTGCTGTGAA
CCTCAGCTGGGAATGTGCATCAGGCAACTCAAGTTTTTTCACCACGGCATGTGTCTGTGAA
TGTCCGCAAAACATTCTCTCTCCCCAGCCTTCATGTGTAAACCTGGGGATGATGTGGACC
TGGGCACTGTGGATGCTCCCTTCACTCTGCAAATTCAGCCTGGCAGCTCTGCCAGCTAAG
CCTGAGAACATTTCTGTGTCTACTACTATAGGAAAAATTTAACCTGCACTTGGAGTCCA
GGAAAGGAAACCAGTTATACCCAGTACACAGTTAAGAGAACTTACGCTTTTGGAGAAAAA
CATGATAATTGTACAACCAATAGTTCTACAAGTGAAAATCGTGCTTCGTGCTCTTTTTTC
CTTCCAAGAATAACGATCCCAGATAATTATACCATTTGAGGTGGAAGCTGAAAATGGAGAT
GGTGTAAATTAATCTCATATGACATACTGGAGATTAGAGAACATAGCGAAAACCTGAACCA
CCTAAGATTTTCCGTGTGAAACCAGTTTTTGGGCATCAAACGAATGATTCAAATTGAATGG
ATAAAGCCTGAGTTGGCGCCTGTTTCATCTGATTTAAAAATACACACTTCGATTCAGGACA
GTCAACAGTACCAGCTGGATGGAAGTCAACTTCGCTAAGAACCGTAAGGATAAAAAACCA
ACGTACAACCTCACGGGGCTGCAGCCTTTTACAGAATATGTCATAGCTCTGCGATGTGCG
GTCAAGGAGTCAAAGTTCTGGAGTGAAGTGGAGCCAAGAAAAAATGGGAATGACTGAGGAA
GAAGCTCCATGTGGCCTGGAAGTGTGGAGAGTCTGAAACCAGCTGAGGCGGATGGAAGA
AGGCCAGTGGGTTGTTATGGAAGAAGGCAAGAGGAGCCCCAGTCCTAGAGAAAAACACTT
GGCTACAACATATGGTACTATCCAGAAAGCAACACTAACCTCACAGAAACAATGAACACT
ACTAACCAGCAGCTTGAAGTGCATCTGGGAGGCGAGAGCTTTTGGGTGTCTATGATTTCT
TATAATTCTCTTGGGAAGTCTCCAGTGGCCACCCTGAGGATTCCAGCTATTCAAGAAAAA
TCATTTTCAGTGCATTGAGGTCATGCAGGCCTGCGTTGCTGAGGACCAGCTAGTGGTGAAG
TGGCAAAGCTCTGCTCTAGACGTGAACACTTGGATGATTGAATGGTTTCCGGATGTGGAC
TCAGAGCCCCACCACCCTTTTCTGGGAATCTGTGTCTCAGGCCACGAACCTGGACGATCCAG
CAAGATAAATTAAAAACCTTTCTGGTGCTATAACATCTCTGTGTATCCAATGTTGCATGAC
AAAGTTGGCGAGCCATATTCCATCCAGGCTTATGCCAAAGAAGGCGTTCCATCAGAAGGT
CCTGAGACCAAGGTGGAGAACATTTGGCGTGAAGACGGTCACGATCACATGGAAGAGATT
CCCAAGAGTGAGAGAAAGGGTATCATCTGCAACTACACCATCTTTTACCAAGCTGAAGGT
GGAAAAGGATTCTGTAAGCACGCCCATAGCGAAGTGGAAAAAACCCTCAAGCCCCAGATA
GATGCTATGGATAGACCTGTTGTAGGCATGGCTCCCCATCTCATTTGTGACTTGCAACCT
GGCATGAATCACTTAGCTTCTTTAAATCTCTCTGAAAATGGGGCCAAGAGCACCCACCTT
TTGGGGTTTTTGGGGTTAAATGAGAGTGAAGTGACAGTACCTGAGAGGAGAGTCTTGAGG
AAATGGAAGGAGTTGTTATAATTTTGTCCTGGTTAGGCCCTGAATTGACCTCCCGGGAGCT
CCCCGACCATCATTCCCAGGAATGGCGTGCCTGGCTTAAAGAGTGAGGAGGAACAGACCC
TGTCACCATGACTTCTACTGCCCCTGCCAAATCATGCTTTTGTTTTTTCAGTCCACCTTAT
CTCCTGACATCTTAAATACTGGGCAAGGCTTGGATTCTTGCTTAGGCTAAATAATTTTTT
CTTATGGTAAAAATACACGTAAAAATTTTTTCCAGTTTAAACATTTGAAAGTGTACAATTT
AGTGGCATTAGAAGCATTCACAATATTGTGCAACCATCACCCTATTTCCAGAACTCTTC
TATTTCTGCCCAAATAGAAGCCCTATACCATTCATTAGTCACTCCCCATTCTCTCTC
CCACAGCCCCCTGGCAACTACCAAAGTCTTTGTGTCTCTATGGATTGCCTATTTTGGATA
TTTCATATACATAGAATCATAAANTAAAAA

387/392

FIGURE 372

MCIRQLKFFTTACVCECPQNILSPQPSCVNLGMMWTWALWMLPSLCKFSLAALPAKPENI
SCVYYYRKNLTCTWSPGKETSYTQYTVKRTYAFGEKHDNCTTNSSTSENRASCSFFLPRI
TIPDNYTIEVEAENG DGVIKSHMTYWRL ENIAKTEPPKIFRVKPV LGIKRMIQIEWIKPE
LAPVSSDLKYTLRFRTVNSTSWMEVNF AKNRKDKNQTYNLTGLQPFTEYVIALRC AVKES
KFWSDWSQEKMGMT EEEAPCGLELWRVLKPAEADGRRPVRL LWKKARGAPVLEKTLGYNI
WYYPESNTNLTETMNTTNQQLELHLGGESFWVSMISYNSLGKSPVATLRIPAIQEKSFQC
IEVMQACVAEDQLVVKWQSSALDVNTWMI EWFPD VDSEPTTLSWESVSQATNWTIQQDKL
KPFWCYNISVYPMLHDKVGEPYSIQAYAKEGVPSEGPETKVENIGVKT V TITWKEIPKSE
RKGII CNYTIFYQAEGGKGFC KHAHSEVEKNPKPQIDAMDRPVVGMAPP SHCDLQPGMNH
LASLNLS ENGA KSTHLLGFWGLNESEVTVPERRVLRKWKELL

Important features of the protein:

Signal peptide:

1-46

Transmembrane domain:

None

N-glycosylation site:

59-63, 69-73, 99-103, 103-107, 125-129, 198-202, 215-219, 219-223, 309-313, 315-319, 412-416, 427-431, 487-491, 545-549, 563-567

N-myristoylation site:

32-38, 137-143, 483-489, 550-556, 561-567

Amidation site:

274-278

Growth factor and cytokines receptors family signature 1:

62-75

Fibronectin type III domain:

54-144

154-247

388/392

FIGURE 373

CCAGGTCCAACCTGCACCTCGGTTCTATCGATTGAATTCCTCGGGGATCCTCTAGAGATCC
CTCGACCTCGACCCACGCGTCCGCCAAGCTGGCCCTGCACGGCTGCAAGGGAGGCTCCTG
TGGACAGGCCAGGCAGGTGGGCCTCAGGAGGTGCCTCCAGGCGGCCAGTGGGCCTGAGGC
CCCAGCAAGGGCTAGGGTCCATCTCCAGTCCCAGGACACAGCAGCGGCCACCATGGCCAC
GCCTGGGCTCCAGCAGCATCAGCAGCCCCCAGGACCGGGGGAGGCACAGGTGGCCCCCAC
CACCCGGAGGAGCAGCTCCTGCCCCCTGTCCGGGGGATGACTGATTCTCCTCCGCCAGGCC
ACCCAGAGGAGAAGGCCACCCCGCCTGGAGGCACAGGCCATGAGGGGCTCTCAGGAGGTG
CTGCTGATGTGGCTTCTGGTGTTGGCAGTGGGCGGCACAGAGCACGCCTACCGGCCCGGC
CGTTAGGGTGTGTGCTGTCCCGGGCTCACGGGGACCCTGTCTCCGAGTCGTTCTGTGCAGC
GTGTGTACCAGCCCTTCTCACCACCTGCGACGGGCACCGGGCCTGCAGCACCTACCGAA
CCATTTATAGGACCGCTACCGCCGAGCCCTGGGCTGGCCCTGCCAGGCCTCGCTACG
CGTGCTGCCCCGGCTGGAAGAGGACCAGCGGGCTTCTGGGGCCTGTGGAGCAGCAATAT
GCCAGCCGCCATGCCGAACGGAGGGAGCTGTGTCCAGCCTGGCCGCTGCCGCTGCCCTG
CAGGATGGCGGGGTGACACTTGCCAGTCAGATGTGGATGAATGCAGTGCTAGGAGGGGCG
GCTGTCCCCAGCGCTGCATCAACACCGCCGGCAGTTACTGGTGCCAGTGTTGGGAGGGGC
ACAGCCTGTCTGCAGACGGTACACTCTGTGTGCCCAAGGGAGGGCCCCCAGGGTGGCCC
CCAACCCGACAGGAGTGGACAGTGCAATGAAGGAAGAAGTGCAGAGGCTGCAGTCCAGGG
TGGACCTGCTGGAGGAGAAGCTGCAGCTGGTGCTGGCCCCACTGCACAGCCTGGCCTCGC
AGGCACTGGAGCATGGGCTCCCGGACCCCGGCAGCCTCCTGGTGCACTCCTTCCAGCAGC
TCGGCCGCATCGACTCCCTGAGCGAGCAGATTTCTTCTTCTGGAGGAGCAGCTGGGGTCCT
GCTCCTGCAAGAAAGACTCGTGACTGCCCAGCGCCCCAGGCTGGACTGAGCCCCTCACGC
CGCCCTGCAGCCCCCATGCCCTGCCCAACATGCTGGGGGTCCAGAAGCCACCTCGGGGT
GACTGAGCGGAAGGCCAGGCAGGGCCTTCTCCTTTTCTCCTCCCCTTCCCTCGGGAGG
GTCCCCAGACCTGGCATGGGATGGGCTGGGATTTTTTTTGTGAATCCACCCCTGGCTAC
CCCCACCCTGGTTACCCCAACGGCATCCCAAGGCCAGGTGGGCCCTCAGCTGAGGGAAGG
TACGAGTTCCCTGCTGGAGCCTGGGACCCATGGCACAGGCCAGGCAGCCCGGAGGCTGG
GTGGGGCCTCAGTGGGGGCTGCTGCCTGACCCCCAGCACATAAAAATGAAACGTGAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAGGGCGGCCGCGACTCTAGAGTC
GACCTGCAGAAGCTTGGCCGCCATGGCCCAACTTGTTTATTGCAGCTTATAATGGTTACAAAT

389/392

FIGURE 374

MTDSPPPGHPPEEKATPPGGTGHEGLSGGAADVASGVGSGRHRARLPARPLGCVLSRAHGD
PVSESFVQRVYQPFLTTCDGHRACSTYRTIYRTAYRRSPGLAPARPRYACCPGWKRTSGL
PGACGAAICQPPCRNGGSCVQPGRRCPCAGWRGDTQCQSDVDECSARRGGCPQRCINTAGS
YWCQCWEGHSLSADGTL CVPKGGPPRVAPNPTGVDSAMKEEVQRLQSRVDLLEEKQLVL
APLHSLASQALEHGLPDPGSLLVHSFQQLGRIDSLSEQISFLEEQLGSCSCKKDS

390/392

FIGURE 375

Wholemount In Situ with PRO1449 Orthologue





FIGURE 376


PRO1449 is expressed in vasculature of many inflamed and diseased tissues

Human tumor tissue (BF)



Human tumor tissue (DF)


DNA64907 in situ



392/392

FIGURE 377

Mouse orthologue of PR01449 induces ectopic vessels in the eyes of chicken embryos



(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
3 January 2002 (03.01.2002)

PCT

(10) International Publication Number
WO 02/000690 A3

(51) International Patent Classification ⁷ : C07K 14/47, C12N 15/12, A61K 38/17, C07K 16/18		09/808,689	14 March 2001 (14.03.2001)	US
		09/816,744	22 March 2001 (22.03.2001)	US
		09/828,366	5 April 2001 (05.04.2001)	US
(21) International Application Number: PCT/US01/19692		09/854,208	10 May 2001 (10.05.2001)	US
		09/854,280	10 May 2001 (10.05.2001)	US
(22) International Filing Date: 20 June 2001 (20.06.2001)		09/866,028	25 May 2001 (25.05.2001)	US
		09/866,034	25 May 2001 (25.05.2001)	US
(25) Filing Language: English		PCT/US01/17092	25 May 2001 (25.05.2001)	US
		09/870,574	30 May 2001 (30.05.2001)	US
(26) Publication Language: English		PCT/US01/17443	30 May 2001 (30.05.2001)	US
		PCT/US01/17800	1 June 2001 (01.06.2001)	US
(30) Priority Data:				
60/213,637	23 June 2000 (23.06.2000)	US	(71) Applicant (for all designated States except US): GENEN- TECH, INC. [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US).	
60/219,556	20 July 2000 (20.07.2000)	US		
60/220,624	25 July 2000 (25.07.2000)	US		
60/220,664	25 July 2000 (25.07.2000)	US		
PCT/US00/20710	28 July 2000 (28.07.2000)	US		
60/222,695	2 August 2000 (02.08.2000)	US		
09/643,657	17 August 2000 (17.08.2000)	US		
PCT/US00/23522	23 August 2000 (23.08.2000)	US		
PCT/US00/23328	24 August 2000 (24.08.2000)	US		
60/230,978	7 September 2000 (07.09.2000)	US		
60/000,000	15 September 2000 (15.09.2000)	US	(72) Inventors; and (75) Inventors/Applicants (for US only): BAKER, Kevin, P. [GB/US]; 14006 Indian Run Drive, Darnestown, MD 20878 (US). FERRARA, Napoleone [US/US]; #704, 2090 Pacific Avenue, San Francisco, CA 94109 (US). GERBER, Hanspeter [CH/US]; #5, 1121 Tennessee Street, San Francisco, CA 94107 (US). GERRITSEN, Mary, E. [CA/US]; 541 Parrott Drive, San Mateo, CA 94402 (US). GODDARD, Audrey [CA/US]; 110 Congo Street, San Francisco, CA 94131 (US). GODOWSKI, Paul, J. [US/US]; 25 Orange Court, Hillsborough, CA 94010 (US). GURNEY, Austin, L. [US/US]; 1 Debbie Lane, Belmont, CA 94002 (US). HILLAN, Kenneth, J. [GB/US]; 64 Seward Street, San Francisco, CA 94114 (US). MARSTERS, Scot, A. [US/US]; 990 Cherry Street, San Carlos, CA 94070 (US). PAN, James [CA/US]; 2705 Coronet Boulevard, Belmont, CA 94002 (US). PAONI, Nicholas, F. [US/US]; 1756 Terrace Drive, Belmont, CA 94002 (US). STEPHAN, Jean-Philippe, F. [FR/US]; 320 C Lansdale Avenue, Millbrae, CA 94030 (US). WATAN- ABE, Colin, K. [US/US]; 128 Corliss Drive, Moraga, CA 94556 (US). WILLIAMS, P., Mickey [US/US]; 509 Alto Avenue, Half Moon Bay, CA 94019 (US). WOOD, William, I. [US/US]; 35 Southdown Court, Hillsborough, CA 94010 (US). YE, Weilan [CN/US]; 119 Barkentine Street, Foster City, CA 94404 (US).	
09/664,610	18 September 2000 (18.09.2000)	US		
09/665,350	18 September 2000 (18.09.2000)	US		
60/242,922	24 October 2000 (24.10.2000)	US		
09/709,238	8 November 2000 (08.11.2000)	US		
PCT/US00/30952	8 November 2000 (08.11.2000)	US		
PCT/US00/30873	10 November 2000 (10.11.2000)	US		
PCT/US00/32678	1 December 2000 (01.12.2000)	US		
09/747,259	20 December 2000 (20.12.2000)	US		
PCT/US00/34956	20 December 2000 (20.12.2000)	US		
09/767,609	22 January 2001 (22.01.2001)	US	(73) Title: COMPOSITIONS AND METHODS FOR THE DIAGNOSIS AND TREATMENT OF DISORDERS INVOLVING AN- GIOGENESIS	
09/796,498	28 February 2001 (28.02.2001)	US		
PCT/US01/06520	28 February 2001 (28.02.2001)	US		
PCT/US01/06666	1 March 2001 (01.03.2001)	US	(57) Abstract: Compositions and methods are disclosed for stimulating or inhibiting angiogenesis and/or cardiovascularization in mammals, including humans. Pharmaceutical compositions are based on polypeptides or antagonists thereto that have been identified for one or more of these uses. Disorders that can be diagnosed, prevented, or treated by the compositions herein include trauma such as wounds, various cancers, and disorders of the vessels including atherosclerosis and cardiac hypertrophy. In addition, the present invention is directed to novel polypeptides and to nucleic acid molecules encoding those polypeptides. Also provided herein are vectors and host cells comprising those nucleic acid sequences, chimeric polypeptide molecules comprising the polypeptides of the present invention fused to heterologous polypeptide sequences, antibodies which bind to the polypeptides of the present invention and to methods for producing the polypeptides of the present invention.	
09/802,706	9 March 2001 (09.03.2001)	US		

[Continued on next page]

WO 02/000690 A3



(74) Agents: AGARWAL, Atulya, R. et al.; c/o GENENTECH, INC., MS49, 1 DNA Way, South San Francisco, CA 94080-4990 (US).

patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

Published:

— with international search report

(88) Date of publication of the international search report:
13 March 2003

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/19692

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/47 A61K38/17 C07K16/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EP0-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,L	WO 99 46281 A (BAKER KEVIN P ;CHEN JIAN (US); GENENTECH INC (US); GURNEY AUSTIN () 16 September 1999 (1999-09-16) see PR0181	1-19
X,L	US 5 968 744 A (CORLEY-NEIL C ET AL) 19 October 1999 (1999-10-19) L: priority. the whole document	1-19
X	WO 99 50405 A (GENETICS INST) 7 October 1999 (1999-10-07) see pk65_4 and seq.ID's 11 and 12	1-19
X	WO 99 43802 A (KATO SEISHI ;KIMURA TOMOKO (JP); PROTEGENE INC (JP); SEKINE SHINGO) 2 September 1999-(1999-09-02)- see clone HP02239.	1-19
-/--		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

21 August 2002

Date of mailing of the international search report

29. 11. 2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Smalt, R

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/19692

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 10363 A (WEI YING FEI ;HUMAN GENOME SCIENCES INC (US); ROSEN CRAIG A (US);) 4 March 1999 (1999-03-04) see secreted protein 24 ---	1-19
X	WO 99 33979 A (CHIRON CORP) 8 July 1999 (1999-07-08) the whole document ---	1-19
A	WO 95 29242 A (CHILDRENS MEDICAL CENTER) 2 November 1995 (1995-11-02) the whole document ---	
A	EP 0 335 243 A (HARVARD COLLEGE) 4 October 1989 (1989-10-04) the whole document ---	
A	FOLKMAN J: "CLINICAL APPLICATIONS OF RESEARCH ON ANGIOGENESIS" NEW ENGLAND JOURNAL OF MEDICINE, THE, MASSACHUSETTS MEDICAL SOCIETY, WALTHAM, MA, US, vol. 333, no. 26, 28 December 1995 (1995-12-28), pages 1757-1763, XP000654557 ISSN: 0028-4793 the whole document ---	
A	KLGSBRUN M: "REGULATORS OF ANGIOGENESIS: STIMULATORS, INHIBITORS, AND EXTRACELLULAR MATRIX" JOURNAL OF CELLULAR BIOCHEMISTRY, LISS, NEW YORK, NY, US, vol. 47, no. 3, 1 November 1991 (1991-11-01), pages 199-200; XP000616588 ISSN: 0730-2312 the whole document ---	
P,X, L	WO 00 53756 A (BOTSTEIN DAVID;BAKER KEVIN P ; GENENTECH INC (US); ASHKENAZI AVI J) 14 September 2000 (2000-09-14) see seq.ID.322; L: priority. ---	1-19
P,X, L	WO 00 53751 A (BAKER KEVIN P ;GENENTECH INC (US); NAPIER MARY A (US); ASHKENAZI A) 14 September 2000 (2000-09-14) the whole document ---	1-21,23, 27-29, 31,32, 34,35
P,X	WO 01 00828 A (CORIXA CORP ;FANGER GARY R (US); RETTER MARC W (US); WANG TONGTONG) 4 January 2001 (2001-01-04) see seq.ID.327 ---	1-21,23, 27-29, 31,32, 34,35

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/19692

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E,L	WO 02 08284 A (FERRARA NAPOLEONE ;STEPHAN JEAN PHILIPPE F (US); WILLIAMS P MICKEY) 31 January 2002 (2002-01-31) See seq.ID's 1 and 120. L: priority. -----	1-35,37, 41

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 01/19692

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 20-34, 37 and 41 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-35, 37, 41 all partially

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Present claims 20-28,31,34,35,37, and 41 relate to products defined by reference to a desirable characteristic or property, namely having (ant)agonistic activity towards the protein(s) of claim 1. The application provides no support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for any such products. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the product by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search impossible. Consequently, the search of said claims, in as far as the (ant)agonists are concerned, has been carried out for those aspects which appear to be clear, supported and disclosed, namely those parts relating to antibodies directed against the protein(s) of claim 1.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

Invention 1: 1-35, 37, 41 all partially

Nucleic acid with at least 80% identity to seq.ID.1 and protein encoded thereby, vector, host cell, method for producing the protein, chimeric protein, antibody, and pharmaceutical compositions.

Inventions 2-187: 1-42, all partially,
and as far as applicable

Subject matter as defined for invention 1. above, but limited to the respective nucleic acid sequences 3-373 (odd numbers) and the polypeptides encoded thereby (seq.ID's 4-374, even numbers).

For the sake of conciseness, the first subject matter is explicitly defined, the other subject matters are defined by analogy thereto.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/19692

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9946281	A	16-09-1999	US 6391311 B1	21-05-2002
			AU 3072199 A	27-09-1999
			CA 2321677 A1	16-09-1999
			US 2002127584 A1	12-09-2002
			US 2002094416 A1	18-07-2002
			WO 9946281 A2	16-09-1999
			US 2002156006 A1	24-10-2002
			US 2002169284 A1	14-11-2002
			ZA 9901937 A	17-04-2000
			EP 1241187 A2	18-09-2002
			AU 3075099 A	11-10-1999
			CA 2322792 A1	23-09-1999
			EP 1064382 A2	03-01-2001
			WO 9947677 A2	23-09-1999
			US 6455283 B1	24-09-2002
			EP 1241183 A2	18-09-2002
			US 2002072496 A1	13-06-2002
			US 2002142961 A1	03-10-2002
			US 2002072067 A1	13-06-2002
			US 2002072092 A1	13-06-2002
			US 2002072497 A1	13-06-2002
			US 2002103125 A1	01-08-2002
			US 2002123463 A1	05-09-2002
			US 2002132252 A1	19-09-2002
			US 2002127576 A1	12-09-2002
			US 2002132253 A1	19-09-2002
			US 2002160384 A1	31-10-2002
			AU 735081 B2	28-06-2001
			AU 1532499 A	15-06-1999
			CA 2309358 A1	03-06-1999
			EP 1241180 A2	18-09-2002
			EP 1032667 A2	06-09-2000
			WO 9927098 A2	03-06-1999
			US 6410708 B1	25-06-2002
			EP 1241185 A2	18-09-2002
			EP 1249491 A2	16-10-2002
			EP 1251139 A2	23-10-2002
			US 2002161199 A1	31-10-2002
			EP 1241179 A2	18-09-2002
			EP 1241186 A2	18-09-2002
			AU 741133 B2	22-11-2001
			AU 3757099 A	08-11-1999
			CA 2324297 A1	28-10-1999
			EP 1071773 A1	31-01-2001
			JP 2002512032 T	23-04-2002
			WO 9954467 A1	28-10-1999
			US 2002147322 A1	10-10-2002
			EP 1241251 A2	18-09-2002
			US 2002119118 A1	29-08-2002
US 5968744	A	19-10-1999	US 2002103342 A1	01-08-2002
			US 6348576 B1	19-02-2002
WO 9950405	A	07-10-1999	AU 3457299 A	18-10-1999
			CA 2324492 A1	07-10-1999
			EP 1068307 A1	17-01-2001
			JP 2002509722 T	02-04-2002
			WO 9950405 A1	07-10-1999

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/19692

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9943802	A	02-09-1999	AU 2549399 A	15-09-1999
			CA 2322711 A1	02-09-1999
			EP 1056847 A2	06-12-2000
			WO 9943802 A2	02-09-1999
			JP 2002506615 T	05-03-2002

WO 9910363	A	04-03-1999	AU 8921598 A	16-03-1999
			CA 2302387 A1	04-03-1999
			EP 1017707 A1	12-07-2000
			JP 2002532054 T	02-10-2002
			WO 9910363 A1	04-03-1999
			US 6410709 B1	25-06-2002
			US 2001016647 A1	23-08-2001

WO 9933979	A	08-07-1999	AU 1929599 A	19-07-1999
			EP 1042470 A2	11-10-2000
			JP 2002511231 T	16-04-2002
			WO 9933979 A2	08-07-1999
			US 2002034800 A1	21-03-2002

WO 9529242	A	02-11-1995	US 5639725 A	17-06-1997
			US 5792845 A	11-08-1998
			AU 692865 B2	18-06-1998
			AU 2461795 A	16-11-1995
			BR 9507479 A	16-09-1997
			CA 2188813 A1	02-11-1995
			CN 1149319 A	07-05-1997
			CZ 9603122 A3	14-05-1997
			EP 0758390 A1	19-02-1997
			HU 76095 A2	30-06-1997
			JP 9512173 T	09-12-1997
			NZ 285501 A	26-02-1998
			WO 9529242 A1	02-11-1995
			US 5885795 A	23-03-1999
			US 5733876 A	31-03-1998
			US 5776704 A	07-07-1998
			ZA 9503419 A	11-01-1996

EP 0335243	A	04-10-1989	US 4900673 A	13-02-1990
			AT 96844 T	15-11-1993
			AU 621358 B2	12-03-1992
			AU 3433189 A	16-10-1989
			CA 1331356 A1	09-08-1994
			DE 68910354 D1	09-12-1993
			DE 68910354 T2	03-03-1994
			DK 233390 A	26-11-1990
			EP 0335243 A2	04-10-1989
			IE 890954 L	28-09-1989
			JP 3503641 T	15-08-1991
			PT 90129 A	10-11-1989
			WO 8909277 A1	05-10-1989
			ZA 8902270 A	24-12-1991

WO 0053756	A	14-09-2000	AU 1748200 A	19-06-2000
			AU 1749900 A	12-07-2000
			AU 2192800 A	12-07-2000
			AU 2224800 A	28-09-2000
			AU 2399300 A	28-09-2000

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/19692

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0053756	A		AU 2495200 A	28-09-2000
			AU 2596700 A	28-09-2000
			AU 2600800 A	28-09-2000
			AU 2879400 A	31-07-2001
			AU 2883600 A	28-09-2000
			AU 3072199 A	27-09-1999
			AU 3107700 A	28-09-2000
			AU 3381600 A	28-09-2000
			AU 3514400 A	28-09-2000
			CA 2321677 A1	16-09-1999
			EP 1173563 A1	23-01-2002
			EP 1159419 A1	05-12-2001
			EP 1220905 A2	10-07-2002
			EP 1135485 A2	26-09-2001
			EP 1141284 A2	10-10-2001
			EP 1159422 A1	05-12-2001
			EP 1141289 A2	10-10-2001
			US 2002119130 A1	29-08-2002
			WO 0053753 A2	14-09-2000
			WO 0053754 A1	14-09-2000
			WO 0053755 A2	14-09-2000
			WO 0153486 A1	26-07-2001
			WO 0053756 A2	14-09-2000
			WO 0053757 A2	14-09-2000
			WO 0053758 A2	14-09-2000
			US 2002127584 A1	12-09-2002
			US 2002094416 A1	18-07-2002
			WO 0032221 A2	08-06-2000
			WO 0053750 A1	14-09-2000
			WO 0037638 A2	29-06-2000
			WO 0037640 A2	29-06-2000
			WO 0053751 A1	14-09-2000
			WO 0053752 A2	14-09-2000
			US 2002119118 A1	29-08-2002
			US 2002161199 A1	31-10-2002
			US 2002156006 A1	24-10-2002
			US 2002169284 A1	14-11-2002
			US 2002072496 A1	13-06-2002
			US 2002142961 A1	03-10-2002
			US 2002072067 A1	13-06-2002
			US 2002072092 A1	13-06-2002
			US 2002072497 A1	13-06-2002
			US 2002103125 A1	01-08-2002
WO 0053751	A	14-09-2000	AU 1748200 A	19-06-2000
			AU 1749900 A	12-07-2000
			AU 2192800 A	12-07-2000
			AU 2224800 A	28-09-2000
			AU 2399300 A	28-09-2000
			AU 2495200 A	28-09-2000
			AU 2596700 A	28-09-2000
			AU 2600800 A	28-09-2000
			AU 2879400 A	31-07-2001
			AU 2883600 A	28-09-2000
			AU 3072199 A	27-09-1999
			AU 3107700 A	28-09-2000
			AU 3381600 A	28-09-2000
			AU 3514400 A	28-09-2000

INTERNATIONAL SEARCH REPORT
Information on patent family members

International Application No

PCT/US 01/19692

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0053751	A	CA 2321677 A1	16-09-1999
		EP 1173563 A1	23-01-2002
		EP 1159419 A1	05-12-2001
		EP 1220905 A2	10-07-2002
		EP 1135485 A2	26-09-2001
		EP 1141284 A2	10-10-2001
		EP 1159422 A1	05-12-2001
		EP 1141289 A2	10-10-2001
		US 2002119130 A1	29-08-2002
		WO 0053753 A2	14-09-2000
		WO 0053754 A1	14-09-2000
		WO 0053755 A2	14-09-2000
		WO 0153486 A1	26-07-2001
		WO 0053756 A2	14-09-2000
		WO 0053757 A2	14-09-2000
		WO 0053758 A2	14-09-2000
		US 2002127584 A1	12-09-2002
		US 2002094416 A1	18-07-2002
		WO 0032221 A2	08-06-2000
		WO 0053750 A1	14-09-2000
		WO 0037638 A2	29-06-2000
		WO 0037640 A2	29-06-2000
		WO 0053751 A1	14-09-2000
		WO 0053752 A2	14-09-2000
		US 2002119118 A1	29-08-2002
		US 2002161199 A1	31-10-2002
		US 2002156006 A1	24-10-2002
		US 2002169284 A1	14-11-2002
		US 2002072496 A1	13-06-2002
		US 2002142961 A1	03-10-2002
		US 2002072067 A1	13-06-2002
		US 2002072092 A1	13-06-2002
		US 2002072497 A1	13-06-2002
		US 2002103125 A1	01-08-2002
WO 0100828	A	04-01-2001	
		AU 5902700 A	31-01-2001
		BR 0012069 A	27-08-2002
		CN 1367829 T	04-09-2002
		EP 1196565 A2	17-04-2002
		HU 3201757 A2	28-09-2002
		NO 20016427 A	25-02-2002
		WO 0100828 A2	04-01-2001
		US 2002168637 A1	14-11-2002
		TR 200200691 T2	21-10-2002
WO 0208284	A	31-01-2002	
		US 2002042367 A1	11-04-2002
		US 2002010137 A1	24-01-2002
		US 2002106743 A1	08-08-2002
		US 2002052027 A1	02-05-2002
		US 2002058309 A1	16-05-2002
		US 2002102723 A1	01-08-2002
		AU 6531101 A	17-12-2001
		AU 7197301 A	05-02-2002
		AU 7315001 A	30-01-2002
		AU 7885201 A	08-01-2002
		WO 0193983 A1	13-12-2001
		WO 0200690 A2	03-01-2002
		WO 0208288 A2	31-01-2002

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/19692

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0208284	A	WO 0208284 A2	31-01-2002
		AU 1916701 A	16-07-2001
		AU 2055401 A	12-06-2001
		AU 6391000 A	19-02-2001
		AU 6802801 A	24-09-2001
		AU 6901800 A	12-06-2001
		EP 1244784 A2	02-10-2002
		EP 1250426 A2	23-10-2002
		WO 0140464 A1	07-06-2001
		WO 0149715 A2	12-07-2001
		WO 0140466 A2	07-06-2001
		WO 0168848 A2	20-09-2001
		US 2002127584 A1	12-09-2002
		US 2002094416 A1	18-07-2002
		US 2002160374 A1	31-10-2002
		US 2002146709 A1	10-10-2002
		US 2002132240 A1	19-09-2002
		US 2002142958 A1	03-10-2002
		US 2002150976 A1	17-10-2002
		US 2002132981 A1	19-09-2002
		US 2002165143 A1	07-11-2002
		US 2002156004 A1	24-10-2002
		US 2002142419 A1	03-10-2002
		US 2002102647 A1	01-08-2002
		US 2002110859 A1	15-08-2002
		US 2002142959 A1	03-10-2002
		US 2002115145 A1	22-08-2002
		US 2002132768 A1	19-09-2002
		US 2002127643 A1	12-09-2002
		US 2002156006 A1	24-10-2002
		US 2002169284 A1	14-11-2002
		US 2002072496 A1	13-06-2002
		US 2002142961 A1	03-10-2002
		US 2002072067 A1	13-06-2002
		US 2002072092 A1	13-06-2002
		US 2002072497 A1	13-06-2002
		US 2002103125 A1	01-08-2002

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.